

Match Overview

1	Internet 35 words crawled on 30-Jul-2010 jasn.asnjournals.org	<1%
2	Internet 27 words crawled on 27-Jan-2015 www.science.gov	<1%
3	Internet 24 words crawled on 11-Feb-2015 www.jimmunol.org	<1%
4	CrossCheck 19 words "Australian and New Zealand Society of Nephrology 38 ... Annual Scientific Meeting Sydney, September 2-4, 2002".	<1%
5	Internet 18 words crawled on 25-Jul-2010 www.nature.com	<1%
6	Internet 18 words crawled on 15-Apr-2014 www.naika.or.jp	<1%

Name of Journal: *World Journal of Nephrology*

ESPS Manuscript NO: 23164

Manuscript Type: ORIGINAL ARTICLE

Basic Study

²² C5b-9 does not mediate tubulointerstitial injury in experimental acute glomerular disease characterized by selective proteinuria

Gopala K. Rangan^{1,2}

学术搜索

获得 32 条结果 (用时0.09秒)

文章

我的图书馆

时间不限

2016以来

2015以来

2012以来

自定义范围...

按相关性排序

按日期排序

搜索所有网页

中文网页

简体中文网页

☒ 包括专利☐ 包含引用

创建快讯

小提示: 只搜索中文(简体)结果, 可在 学术搜索设置 指定搜索语言

How does proteinuria cause progressive renal damage?

[M Abbate](#), [C Zoja](#), [G Remuzzi](#) - Journal of the American Society of ..., 2006 - Am Soc Nephrol
... C3 and other complement factors (77) in ways that may have critical importance in disease, as ...
A direct role for protein overload as a stimulus was indicated by findings that the ... This approach
would obviate the need to inhibit complement systemically and its function in innate ...
被引用次数: 466 相关文章 所有 6 个版本 引用 保存

asnjournals.org 中的 [HTML]

[HTML] Proteinuria and interstitial injury

[AA Eddy](#) - Nephrology Dialysis Transplantation, 2004 - ERA-EDTA
... treated animals developed less interstitial inflammation and fibrosis and had better renal function
than control ... Both of these receptors were shown to modify specific macrophage phagocytic
functions that may serve ... The role of proteinuria in the progression of chronic renal failure ...
被引用次数: 146 相关文章 所有 6 个版本 引用 保存

oxfordjournals.org 中的 [HTML]

Cellular responses to protein overload: key event in renal disease progression

[C Zoja](#), [A Benigni](#), [G Remuzzi](#) - Current opinion in nephrology and ..., 2004 - journals.lww.com
... Studies in the early 1980s by Brenner's group [1] highlighted the central role of changes ...
inflammation and fibrosis, which are ultimately responsible for renal scarring and loss of function. ...
expressed at the brush border of proximal tubular cells, interact to mediate the reabsorption ...
被引用次数: 142 相关文章 所有 5 个版本 引用 保存

Complement-mediated dysfunction of glomerular filtration barrier accelerates progressive renal injury

[M Abbate](#), [C Zoja](#), [D Corna](#), [D Rottoli](#)... - Journal of the ..., 2008 - Am Soc Nephrol
... the lack of C6 was not protective in nonproteinuric models, 22 indirectly suggesting the importance
of apical ... Our data are fully consistent with a role of protein overload of glomerular epithelial cells
in ... Animals without graft function died within 36 h of the contralateral nephrectomy ...

asnjournals.org 中的 [HTML]

Scholar

About 276 results (0.08 sec)

Articles

Case law

My library

Any time

Since 2016

Since 2015

Since 2012

Custom range...

Sort by relevance

Sort by date

☒ include patents☒ include citations Create alert**How does proteinuria cause progressive renal damage?**

[M Abbate](#), [C Zoja](#), [G Remuzzi](#) - Journal of the American Society of ..., 2006 - Am Soc Nephrol
 ... How **Does Proteinuria** Cause Progressive Renal **Damage** ... heat-shock protein 47 also was upregulated in primary cultures of mouse proximal tubular epithelial cells in response to **C5b-9**, further suggesting that complement activation on tubular cells **can** stimulate the ...
 Cited by 457 Related articles All 5 versions Web of Science: 290 Cite Save

[HTML] from asnjournals.org
 @ University of Sydney

[HTML] Understanding the nature of renal disease progression

[G Remuzzi](#), [P Ruggenenti](#), [A Benigni](#) - Kidney international, 1997 - nature.com
 ... role of filtered complement might provide a partial explanation as to why **selective proteinuria does not** necessarily lead to ... Hence, the **C5b-9** complex is found in the urine of a number of renal conditions ... with those reported in IDD [149] and show that ACE inhibition **can** lower the ...
 Cited by 703 Related articles All 5 versions Web of Science: 494 Cite Save

[HTML] from nature.com
 @ University of Sydney

Role of the microvascular endothelium in progressive renal disease

[DH Kang](#), [J Kanellis](#), [C Hugo](#), [L Truong](#) ... - Journal of the ..., 2002 - Am Soc Nephrol
 ... As discussed above, TSP-1 is a potent inhibitor of endothelial cell proliferation and **will** also directly initiate ... Tumor necrosis factor- α (TNF- α) **does not** decrease VEGF secretion at the concentration of 10 ng/ml ... **Can** Administration of Angiogenic Agents Prevent Renal Progression ...
 Cited by 433 Related articles All 5 versions Web of Science: 271 Cite Save

[HTML] from asnjournals.org
 @ University of Sydney

Interstitial nephritis induced by protein-overload proteinuria

[AA Eddy](#) - The American journal of pathology, 1989 - ricbi.nlm.nih.gov
 ... 35'1 Monoclonal antibodies reactive with subsets of lym- phohemopoietic cells **can** be used ... rabbit antiserum to rat albumin as pre- viously described. 12 This antiserum **does not** cross react ... Tissue was stained for rat **C5b-9** neoantigens using monoclonal antibody 2A1 (a gift of Dr ...
 Cited by 303 Related articles All 7 versions Web of Science: 262 Cite Save

[PDF] from nih.gov
 @ University of Sydney

Complement inhibitors targeted to the proximal tubule prevent injury in experimental nephrotic syndrome and demonstrate a key role for C5b-9

[C He](#), [M Imai](#), [H Song](#), [RJ Quigg](#) ... - The Journal of ..., 2005 - Am Assoc Immunol
 ... 7% and 8%). We **did not** measure creatinine clearances or perform survival renal biopsies to ... the inaccessibility of inflammatory cells bearing receptors for the latter, whereas **C5b-9 can** act directly on ... A high dose of enalapril **can** reduce **proteinuria** in the early phase of **disease** ...
 Cited by 71 Related articles All 4 versions Web of Science: 47 Cite Save

[HTML] from jimmunol.org
 @ University of Sydney

Pathophysiology of proteinuria

[HTML] from nature.com

[全部](#)[图片](#)[新闻](#)[视频](#)[更多 ▾](#)[搜索工具](#)

找到约 3,830 条结果 (用时 0.87 秒)

Leaked protein and interstitial damage in the kidney: is ...

www.ncbi.nlm.nih.gov > ... > PubMed Central (PMC) ▾ [翻译此页](#)

作者: NS SHEERIN - 2002 - 被引用次数: 25 - 相关文章

High urinary protein levels in patients with glomerular disease predicts an ... However, C5b-9 can be found in the urine from proteinuric patients who do not have C5b-9 ... falling well within the range of proteins filtered in non-selective proteinuria. ... role in mediating proteinuria-associated tubulointerstitial injury, complement ...

Complement-Mediated Dysfunction of Glomerular Filtration ...

www.ncbi.nlm.nih.gov > ... > PubMed Central (PMC) ▾ [翻译此页](#)

作者: M Abbate - 2008 - 被引用次数: 38 - 相关文章

Protein overload led to the development of glomerular injury, accumulation of ... first characterized the renal disease phenotype of protein overload proteinuria in tubule cell origin is a crucial mediator of injury in experimental acute allograft Couser WG: C5b-9 does not mediate chronic tubulointerstitial disease in the ...

Proteinuria and interstitial injury

ndt.oxfordjournals.org/content/19/2/277.full ▾ [翻译此页](#)

作者: AA Eddy - 2004 - 被引用次数: 146 - 相关文章

Although there does not appear to be an immunological response to the albumin, ... (1) In glomerular diseases associated with proteinuria, proximal tubular cells may be The recruitment of macrophages to sites of acute injury represents a (C5b-9) mediates interstitial disease in experimental nephrotic syndrome.

Proteinuria and Tubulointerstitial Injury: The ... - Karger

<https://www.karger.com/Article/Pdf/71734> - [翻译此页](#)

glomerular injury, but also the independent factor to promote tubulointerstitial ... diseases that persistent proteinuria stimulates proximal tubular epithelial ... In adriamycin