

Name of Journal: *World Journal of Nephrology*

ESPS Manuscript NO: 25036

Manuscript Type: Case Control Study

Diabetes mellitus increases the prevalence of anemia in patients with chronic kidney disease: a nested case-control study.

Charalampos Loutradis, Alexandra Skodra, Panagiotis Georgianos, Panagiota Tolika, Dimitris Alexandrou, Afroditi Avdelidou, Pantelis A. Sarafidis

Match Overview

1	CrossCheck 146 words Moschopoulou, Maria, Foteini Ch. Ampatzidou, Charalampos Loutradis, Afroditi Boutou, Charilaos-Panagiotis K	3%
2	Internet 45 words crawled on 11-Dec-2014 www.ncbi.nlm.nih.gov	1%
3	CrossCheck 31 words Meininger, Gary, William Canovatchel, David Polidori, et al., and Norm Rosenthal. "Canagliflozin for the treatment of ad	1%
4	Internet 28 words crawled on 07-Jan-2016 www.healthyliving2015.nl	1%
5	CrossCheck 26 words "Saturday, June 26, 2010: Poster Session 1", NDT Plus, 2010.	1%
6	CrossCheck 23 words Speeckaert, M., W. Van Biesen, J. Delanghe, R. Slingerland, A. Wiecek, J. Heaf, C. Drechsler, R. Lacatus, R. Vanh	1%

学术搜索

找到约 1,870 条结果 (用时0.09秒)

小提示: 只搜索中文(简体)结果, 可在 学术搜索设置 指定搜索语言

时间不限

2016以来

2015以来

2012以来

自定义范围...

按相关性排序

按日期排序

搜索所有网页

中文网页

简体中文网页

☒ 包括专利☒ 包含引用

Relationship between heart failure treatment and development of worsening renal function among hospitalized patients

J Butler, DE Forman, WT Abraham, SS Gottlieb... - American heart ..., 2004 - Elsevier

... Net increase in dose from admission to the day before WRF was 92 mg in patients and 64 mg in control subjects. ... A history of diabetes mellitus or HF, systolic blood pressure >160 mm Hg or creatinine level >221.0 μ mol/L (>2.5 mg/dL ... History of diabetes, 5.85, 1.74, 1.11-2.73, .02. ...

被引用次数: 422 相关文章 所有 8 个版本 引用

[HTML] The risk of acute renal failure in patients with chronic kidney disease

CY Hsu, JD Ordonez, GM Chertow, D Fan... - Kidney international, 2008 - Elsevier

... The risk then increases sharply for increasing severity of baseline kidney disease (Table 3 ... GFR 45-59 ml per min per 1.73 m² had a twofold increase in adjusted ... 20 These included age, sex, race/ethnicity, diabetes mellitus, 20, 29 diagnosed hypertension (by ICD-9 codes 401 ...

被引用次数: 264 相关文章 所有 7 个版本 引用

[HTML] The clinical epidemiology of cardiac disease in chronic renal failure

PS Parfrey, RN Foley - Journal of the American Society of ..., 1999 - Am Soc Nephrol

... These data are compatible with the belief that systolic hypertension causes concentric hypertrophy, and that anemia and hemodynamic overload cause an increase in LV volume, with further ... Diabetes Mellitus. ... Outcomes of diabetic relative to nondiabetic dialysis patients (23). ...

被引用次数: 641 相关文章 所有 6 个版本 引用

Vitamin D levels and early mortality among incident hemodialysis patients

M Wolf, A Shah, O Gutierrez, E Ankers, M Monroy... - Kidney ..., 2007 - nature.com

... Although this high prevalence may relate in part to the large representation of black and diabetic subjects, which is typical ... Finally, active vitamin D use increases over time on hemodialysis, a phenomenon that would limit the statistical power of ... Diabetes Care 2006; 29: 650-656 ...

被引用次数: 645 相关文章 所有 7 个版本 引用

sciencedirect.com

asnjournals.org 中

nature.com 中的 [

学术搜索

找到约 1,900 条结果 (用时0.12秒)

小提示: 只搜索中文(简体)结果, 可在 学术搜索设置 指定搜索语言

时间不限

2016以来

2015以来

2012以来

自定义范围...

按相关性排序

按日期排序

搜索所有网页

中文网页

简体中文网页

☒ 包括专利

☒ 包含引用

Relationship between heart failure treatment and development of worsening renal function among hospitalized patients

J Butler, DE Forman, WT Abraham, SS Gottlieb... - American heart ..., 2004 - Elsevier

... Net increase in dose from admission to the day before WRF was 92 mg in patients and 64 mg in control subjects. ... A history of diabetes mellitus or HF, systolic blood pressure >160 mm Hg or creatinine level >221.0 μmol/L (>2.5 mg/dL ... History of diabetes, 5.85, 1.74, 1.11-2.73, .02 ...

被引用次数: 424 相关文章 所有 8 个版本 引用

[HTML] The risk of acute renal failure in patients with chronic kidney disease

CY Hsu, JD Ordonez, GM Chertow, D Fan... - Kidney international, 2008 - Elsevier

... The risk then increases sharply for increasing severity of baseline kidney disease (Table 3 ... GFR 45-59 ml per min per 1.73 m² had a twofold increase in adjusted ... 20 These included age, sex, race/ethnicity, diabetes mellitus, 20, 29 diagnosed hypertension (by ICD-9 codes 401 ...

被引用次数: 272 相关文章 所有 7 个版本 引用

[HTML] The clinical epidemiology of cardiac disease in chronic renal failure

PS Parfrey, RN Foley - Journal of the American Society of ..., 1999 - Am Soc Nephrol

... These data are compatible with the belief that systolic hypertension causes concentric hypertrophy, and that anemia and hemodynamic overload cause an increase in LV volume, with further ... Diabetes Mellitus. ... Outcomes of diabetic relative to nondiabetic dialysis patients (23). ...

被引用次数: 647 相关文章 所有 6 个版本 引用

Vitamin D levels and early mortality among incident hemodialysis patients

M Wolf, A Shah, O Gutierrez, E Ankers, M Monroy... - Kidney ..., 2007 - nature.com

... Although this high prevalence may relate in part to the large representation of black and diabetic subjects, which is typical ... Finally, active vitamin D use increases over time on hemodialysis, a phenomenon that would limit the statistical power of ... Diabetes Care 2006; 29: 650-656 ...

被引用次数: 651 相关文章 所有 7 个版本 引用

[全部](#)[新闻](#)[图片](#)[视频](#)[更多 ▾](#)[搜索工具](#)

找到约 55,500 条结果 (用时 0.28 秒)

[PDF] Prevalence of Cardiovascular Risk Factors in Patients wit...

medind.nic.in/jac/t09/i1/jact09i1p23.pdf ▾ [翻译此页](#)

作者: S Kalra - [相关文章](#)

Results: The mean age for **cases** and **controls** was 53.3 (SD \pm 13.7) and 51.5 (SD ... The **prevalence** of **diabetes mellitus**, hypertension, and smoking was higher ... Conclusion: **Patients** with **CKD** have higher **prevalence** of risk factors for CVD. ... have shown that this **increased** cardiovascular risk begins ... It was a **nested**.

Revascularization Options in Patients with Chronic Kidney Dis...

www.ncbi.nlm.nih.gov > ... > [PubMed Central \(PMC\)](#) ▾ [翻译此页](#)

作者: G Ashrith - 2010 - 被引用次数: 24 - [相关文章](#)

The **prevalence** of end-stage renal disease (ESRD), defined as renal dysfunction ... Studies of revascularization in **chronic kidney disease patients** who do not require ... as in **patients** with **diabetes mellitus**, use of the internal mammary artery bypass graft end stage renal disease: a prospective, **nested case-control study**.

Stroke in patients with type 2 diabetes mellitus, chronic kidney...

www.ncbi.nlm.nih.gov/pubmed/22104547 ▾ [翻译此页](#)

作者: H Skali - 2011 - 被引用次数: 55 - [相关文章](#)

2011年11月21日 - Stroke in **patients** with type 2 **diabetes mellitus**, **chronic kidney disease**, ... **increased** risk related to darbepoetin alfa, we performed a **nested case-control** analysis (1:10 matching) identifying nonstroke **controls** ... Aged; **Anemia**/drug therapy*; **Anemia/epidemiology***; **Case-Control Studies**; Darbepoetin alfa ...

The risk of acute renal failure in patients with chronic kidney di...

www.ncbi.nlm.nih.gov > ... > [PubMed Central \(PMC\)](#) ▾ [翻译此页](#)

作者: CY Hsu - 2008 - 被引用次数: 273 - [相关文章](#)

2008年4月2日 - Pre-admission baseline **diabetes mellitus**, diagnosed hypertension and