

Name of Journal: *World Journal of Nephrology*

ESPS Manuscript NO: 25453

Manuscript Type: ORIGINAL ARTICLE

Observational Study

Is overhydration in Peritoneal Dialysis patients associated with cardiac mortality that might be reversible?

Elizabeth Oei, Klara Paudel, Annemarie Visser, Hazel Finney (MSc), Stanley L Fan (FRCP)

Overhydration; a modifiable cardiac risk?

Match Overview

1

Internet 160 words
crawled on 11-Jun-2014
www.bcm-fresenius.com

4%

2

CrossCheck 158 words
O'Lone, E. L., A. Visser, H. Finney, and S. L. Fan. "Clinical significance of multi-frequency bioimpedance spectroscopy ..."

4%

学术搜索

找到约 879 条结果 (用时0.10秒)

小提示: 只搜索中文(简体)结果, 可在 学术搜索设置 指定搜索语言

时间不限

2016以来

2015以来

2012以来

自定义范围...

按相关性排序

按日期排序

搜索所有网页

中文网页

简体中文网页

☒ 包括专利☒ 包含引用**Cardiovascular risk in the peritoneal dialysis patient**

RT Krediet, O Balafa - Nature Reviews Nephrology, 2010 - nature.com

... patients on hemodialysis after 1 year of therapy, 128 suggesting that the dialysate glucose concentration does not have ... Residual renal function is related to all-cause mortality and risk of cardiovascular death in patients on hemodialysis or ... Ultrafiltration failure and overhydration. ...

被引用次数: 92 相关文章 所有 5 个版本 引用

nature.com 中的

Crush injuries with impairment of renal function

EGL Bywaters, D Beall - Br Med J, 1941 - Springer

... invented a plastic conduit for placement into the peritoneum for repeated peritoneal lavage. ... F. Arteriovenous hemofiltration; a new simple method for treatment of overhydrated patients resistant to ... Weaknesses There is some concern that the two patient groups may not have ...

被引用次数: 874 相关文章 所有 15 个版本 引用

nih.gov 中的 [PDF]

[HTML] Pathophysiology of cardiovascular disease in hemodialysis patients

F Meeus, O Kourilsky, AP Guerin, C Gaudry... - Kidney ..., 2000 - nature.com

... tightly related to hyperkinetic circulation caused by anemia, arteriovenous fistula, or overhydration and is ... in ESRD were the elastic modulus and dilation of arteries, that is, generalized arterial ... Dialysis by itself does not improve arterial distensibility 78, and some studies indicate ...

被引用次数: 135 相关文章 所有 5 个版本 引用

nature.com 中的 Full View

Altered morphologic properties of large arteries in children with chronic renal failure and after renal transplantation

M Litwin, E Wühl, C Jourdan, J Trelewicz... - Journal of the ..., 2005 - Am Soc Nephrol

... overt inflammatory disease at the time of investigation, clinically significant overhydration or dehydration ... already in children and adolescents is in keeping with but does not prove the ... In conclusion, we demonstrated that CKD is associated with morphologic alterations of the large ...

被引用次数: 170 相关文章 所有 5 个版本 引用

asnjournals.org 中

学术搜索

找到约 823 条结果 (用时0.10秒)

文章

我的图书馆

时间不限

2016以来

2015以来

2012以来

自定义范围...

按相关性排序

按日期排序

搜索所有网页

中文网页

简体中文网页

☒ 包括专利☐ 包含引用☒ 创建快讯

小提示: 只搜索中文(简体)结果, 可在 学术搜索设置 指定搜索语言

Cardiovascular risk in the peritoneal dialysis patient

RT Krediet, O Balafa - Nature Reviews Nephrology, 2010 - nature.com

... patients on hemodialysis after 1 year of therapy, 128 suggesting that the dialysate glucose concentration does not have ... Residual renal function is related to all-cause mortality and risk of cardiovascular death in patients on hemodialysis or ... Ultrafiltration failure and overhydration. ...

被引用次数: 92 相关文章 所有 5 个版本 引用 保存

Crush injuries with impairment of renal function

EGL Bywaters, D Beall - Br Med J, 1941 - Springer

... invented a plastic conduit for placement into the peritoneum for repeated peritoneal lavage. ... F. Arteriovenous hemofiltration; a new simple method for treatment of overhydrated patients resistant to ... Weaknesses There is some concern that the two patient groups may not have ...

被引用次数: 876 相关文章 所有 15 个版本 引用 保存

[HTML] Pathophysiology of cardiovascular disease in hemodialysis patients

F Meeus, O Kourilsky, AP Guerin, C Gaudry... - Kidney ..., 2000 - nature.com

... tightly related to hyperkinetic circulation caused by anemia, arteriovenous fistula, or overhydration and is ... in ESRD were the elastic modulus and dilation of arteries, that is, generalized arterial ... Dialysis by itself does not improve arterial distensibility 78, and some studies indicate ...

被引用次数: 136 相关文章 所有 5 个版本 引用 保存

Altered morphologic properties of large arteries in children with chronic renal failure and after renal transplantation

M Litwin, E Wühl, C Jourdan, J Trelewicz... - Journal of the ..., 2005 - Am Soc Nephrol

... overt inflammatory disease at the time of investigation, clinically significant overhydration or dehydration ... already in children and adolescents is in keeping with but does not prove the ... In conclusion, we demonstrated that CKD is associated with morphologic alterations of the large ...

[全部](#)[视频](#)[新闻](#)[图片](#)[更多 ▾](#)[搜索工具](#)

找到约 47,400 条结果 (用时 0.58 秒)

Google 学术: Is overhydration in Peritoneal Dialysis patients associated with cardiac mortality that might be reversible?

... mortality risk of overhydration in haemodialysis patients - [Wizemann](#) - 被引用次数: 322

Cardiovascular risk in the peritoneal dialysis patient - [Krediet](#) - 被引用次数: 96

... of cardiovascular disease in hemodialysis patients - [Meeus](#) - 被引用次数: 139

Volume overhydration is related to endothelial dysfunction in ... - NCBI

www.ncbi.nlm.nih.gov/pubmed/18556383 ▾ [翻译此页](#)

作者: LT Cheng - 2008 - 被引用次数: 36 - [相关文章](#)

OBJECTIVE: In dialysis patients, volume overhydration is common and is related to increased risk of cardiovascular morbidity and mortality. ... 81 stable patients on continuous ambulatory peritoneal dialysis in a single center were recruited. ... The results of this study may help us understand the underlying mechanism of ...

Fluid Status in Peritoneal Dialysis Patients: The European Body ...

journals.plos.org/plosone/article?id=10.1371/journal.pone.0017148 ▾ [翻译此页](#)

作者: W Van Biesen - 2011 - 被引用次数: 115 - [相关文章](#)

2011年2月24日 - Close monitoring of volume status, a correct dialysis prescription adapted to ... peritoneal dialysis (PD) patients, as [1], [2] volume overload is related to cardiac It can be hypothesized that fluid overload is induced by not adapting the (2009) The mortality risk of overhydration in haemodialysis patients.

Overhydration prevalence in peritoneal dialysis – A 2 year longitudinal ...

www.revistanefrologia.com/en-publicacion-nefrologia-articulo-overhydratio... ▾ [翻译此页](#)

作者: PV Aguiar - 2015 - [相关文章](#)