

Name of Journal: *World Journal of Nephrology*

ESPS Manuscript: 27079

Manuscript Type: Prospective Study

What is the optimal level of vitamin D in non-dialysis chronic kidney disease population?

Pablo Molina, José L Górriz, Mariola D Molina, Sandra Beltrán, Belén Vizcaíno, Verónica Escudero, Julia Kanter, Ana I Ávila, Jordi Bover, Elvira Fernández, Javier Nieto, Secundino Cigarrán, Enrique Gruss, Gema Fernández-Juárez, Alberto Martínez-Castelao, Juan F Navarro-González, Ramón Romero, Luis M Pallardó

Match Overview

| | | |
|---|--|-----|
| 1 | Internet 37 words crawled on 11-Nov-2008 www2.kemc.co.uk | 1% |
| 2 | CrossCheck 24 words Claudia Boucher-Berry. "Vitamin D, osteocalcin, and ri... for adiposity as comorbidities in middle school childre | <1% |
| 3 | Internet 24 words crawled on 06-Sep-2014 run.unl.pt | <1% |
| 4 | Internet 21 words crawled on 10-Jan-2016 www.vitamindwiki.com | <1% |
| 5 | Internet 21 words crawled on 26-May-2016 jasn.asnjournals.org | <1% |
| 6 | Internet 20 words crawled on 12-Mar-2016 www.dovepress.com | <1% |

学术搜索

获得 4 条结果 (用时0.11秒)

小提示: 只搜索中文(简体)结果, 可在 学术搜索设置 指定搜索语言

时间不限

2016以来

2015以来

2012以来

自定义范围...

按相关性排序

按日期排序

搜索所有网页

中文网页

简体中文网页

☒ 包括专利

☒ 包含引用

Vitamin D deficiency

MF Holick - New England Journal of Medicine, 2007 - Mass Medical Soc

... kidney disease, 25-hydroxyvitamin D should be measured annually, and the level should be ... Sensible sun exposure can provide an adequate amount of vitamin D 3, which is stored ... E, Willett WC, Dietrich T, Dawson-Hughes B. Estimation of optimal serum concentrations of 25 ...

被引用次数: 8613 相关文章 所有 78 个版本 引用

The effects of normal as compared with low hematocrit values in patients with cardiac disease who are receiving hemodialysis and epoetin

A Besarab, WK Bolton, JK Browne... - ... England Journal of ..., 1998 - Mass Medical Soc

... The interim results of the study did not reach the prespecified stopping boundary corresponding to an overall 5 percent level of significance, complicating the ... 3. Tajinderpal Saraon, Stuart D. Katz. ... (2015) High Dose ESAs Are Associated with High iPTH Levels in Hemodialysis ...

被引用次数: 1781 相关文章 所有 5 个版本 引用

Clinical Practice Guideline for the Management of Chronic Kidney Disease - Mineral and Bone Disorder

M Fukagawa, K Yokoyama, F Koiwa... - Therapeutic ..., 2013 - Wiley Online Library

... been adopted as a treatment tool for keeping serum levels of serum phosphorus and calcium optimal. ... has different properties, there is no consensus regarding which phosphorus binder is best for improving ... 2.5 mEq/L or 3.0 mEq/L KDIGO states that a calcium level between 2.5 ...

被引用次数: 76 相关文章 所有 7 个版本 引用

MOON JU-YOUNG1, KIM SU-MI1, JEONG KYUNG-HWAN1, LEE SANG-HO1, LEE TAE-WON1, IHM CHUN-GYOO1, LIM SUNG-JIG2 1Division of Nephrology, ...

H KAZUHIRO, W SHU, H KOICHI, I HIROSHI - 2014 - Wiley Online Library

... Conclusion: Optimal time period to diagnose resistance to immunosuppressive therapy is 1-2 months in ... higher than that in control ($7.2 \pm 6.81\%$, $P = 0.0005$) and correlated with amount of ...



Scholar

3 results (0.10 sec)

Articles

Case law

My library

Any time

Since 2016

Since 2015

Since 2012

Custom range...

Sort by relevance

Sort by date

☒ include patents

☒ include citations

☐ Create alert

[HTML] The effects of normal as **compared** with low hematocrit values in patients with cardiac disease who are receiving hemodialysis and epoetin

A Besarab, WK Bolton, JK Browne... - ... England Journal of ..., 1998 - Mass Medical Soc

... not reach the prespecified stopping boundary corresponding to an overall 5 percent **level** of significance ... and hypertension as a cause of **renal** failure) than the general US **population** of patients ... Braumann KM , Nonnast-Daniel B , Boning D , Bocker A , Frei U . Improved physical ...

Cited by 1813 Related articles All 7 versions Cite Save

Clinical practice guideline for the management of **chronic kidney disease**-mineral and bone disorder

M Fukagawa, K Yokoyama, F Koiwa... - Therapeutic ..., 2013 - Wiley Online Library

... been adopted as a treatment tool for keeping **serum levels** of **serum** phosphorus and calcium **optimal**. ... has different properties, there is no consensus regarding which phosphorus binder is **best** for improving ... 2.5 mEq/L or 3.0 mEq/L. KDIGO states that a calcium **level** between 2.5 ...

Cited by 80 Related articles All 8 versions Cite Save

MOON JU-YOUNG¹, KIM SU-MI¹, JEONG KYUNG-HWAN¹, LEE SANG-HO¹, LEE TAE-WON¹, IHM CHUN-GYOO¹, LIM SUNG-JIG² ¹Division of Nephrology, ...

H KAZUHIRO, W SHU, H KOICHI, I HIROSHI - 2014 - Wiley Online Library

... Conclusion: **Optimal** time period to diagnose resistance to immunosuppressive therapy is 1–2 months in ... higher than that in control ($7.2 \pm 6.81\%$, $P = 0.0005$) and correlated with **amount** of proteinuria ... of (P)RR [s(P)RR] is secreted into the extracellular space and **serum level** of s ...

All 2 versions Cite Save

全部

图片

新闻

视频

更多 ▾

搜索工具

找到约 1,220,000 条结果 (用时 0.53 秒)

NKF KDOQI Guidelines - National Kidney Foundation

www2.kidney.org/professionals/kdoqi/guidelines_bone/guide7.htm ▾ 翻译此页

Serum **levels** of 25-hydroxyvitamin D (not the **levels** of 1,25-dihydroxyvitamin D) are ... and GFR of 20 to 60 mL/min/1.73 m², and in **CKD** patients undergoing **dialysis**. The prevention and treatment of **vitamin D** insufficiency in patients with **CKD** ... in the normal **population**, where 1,25(OH)₂D **levels** are not dependent on the ...

Vitamin D Therapy in Chronic Kidney Disease and End Stage Renal ...

www.ncbi.nlm.nih.gov ▾ NCBI ▾ Literature ▾ PubMed Central (PMC) ▾ 翻译此页

作者: ML Melamed - 2012 - 被引用次数: 48 - 相关文章

Low **Vitamin D Levels** in **CKD** and ESRD and Outcomes. Multiple **Vitamin D** compounds for people with **chronic kidney disease** not requiring **dialysis**.

Vitamin D and Chronic Kidney Disease

www.ncbi.nlm.nih.gov ▾ NCBI ▾ Literature ▾ PubMed Central (PMC) ▾ 翻译此页

作者: S Williams - 2009 - 被引用次数: 48 - 相关文章

In patients with **CKD**, the new **non-classical** role of **vitamin D** also may be required to maintain optimum levels of **vitamin D** in normal **populations**, and it is ...

Vitamin D Metabolism and Treatment in Chronic Kidney Disease ...

www.medscape.org/viewarticle/571558_2 ▾ 翻译此页

Biologic motivation for minimizing **vitamin D** activity in **CKD** is not entirely clear ... the **prevalence** of overt **vitamin D** deficiency (25-hydroxy **vitamin D levels** < 15 Controversy exists as to **optimal** 25-hydroxy **vitamin D levels** with respect to health. ... of activated **vitamin D** therapy in dialysis and **nondialysis** patients with **CKD** ...

Vitamin D resistance in chronic kidney disease (CKD) - BMC Nephrology

bmopen.bmc-hjmedcentral.com/articles/10.1186/1471-2260-15-17 ▾ 翻译此页