



Name of Journal: *World Journal of Nephrology*

ESPS Manuscript NO: 27100

Manuscript Type: Original Article

*Case Control Study*

Any Link of Gout Disease Control Among Hypertensive Patients and Onset of End-stage Renal Disease? Results from a Population-Based Study

Sylvie Perreault, Javier Nuevo, Scott Baumgartner, and Robert Morlock

## Match Overview

1	Crossref 164 words Miyaoka, Tokiko, Toshio Mochizuki, Takashi Takei, Ken Tsuchiya, and Kosaku Nitta. "Serum uric acid levels and ..."	3%
2	Crossref 139 words S. Perreault. "Better adherence to antihypertensive agents and risk reduction of chronic heart failure", Journal of I	3%
3	Internet 125 words crawled on 06-Sep-2016 <a href="http://acrabstracts.org">acrabstracts.org</a>	3%
4	Crossref 97 words De Vera, Mary A., Greg Marcotte, Sharan Rai, Jessica S. Galo, and Vidula Bhole. "Medication adherence in gout"	2%
5	Internet 55 words crawled on 28-Nov-2014 <a href="http://www.ncbi.nlm.nih.gov">www.ncbi.nlm.nih.gov</a>	1%
6	Crossref 42 words	1%

学术搜索

找到约 1,100 条结果 (用时0.13秒)

时间不限

2016以来

2015以来

2012以来

自定义范围...

按相关性排序

按日期排序

搜索所有网页

中文网页

简体中文网页

☒ 包括专利☒ 包含引用

小提示: 只搜索中文(简体)结果, 可在 学术搜索设置 指定搜索语言

### Diabetes, hypertension, and cardiovascular disease an update

JR Sowers, M Epstein, ED Frohlich - *Hypertension*, 2001 - Am Heart Assoc

... a major pathway for insulin signal transduction 23 24 25 (Figure 2). The relationship between the RAS ... 26 This pathway has also been implicated in mediating the activation of NO synthase and ... 19 Finally, the use of ACE inhibitors or Ang II antagonists in any pregnant woman ...

被引用次数: 1184 相关文章 所有 9 个版本 引用

### Essential hypertension, progressive renal disease, and uric acid: a pathogenetic link?

RJ Johnson, MS Segal, T Srinivas, A Ejaz... - *Journal of the ...*, 2005 - Am Soc Nephrol

... Essential Hypertension, Progressive Renal Disease, and Uric Acid: A Pathogenetic Link? ... ESRD is correct for the large number of cases reported (6,7). There is no doubt that ... Younger subjects with short-duration (mean, 3 yr) hypertension also do not show any increase in ...

被引用次数: 231 相关文章 所有 7 个版本 引用

### Potential role of sugar (fructose) in the epidemic of hypertension, obesity and the metabolic syndrome, diabetes, kidney disease, and cardiovascular disease

RJ Johnson, MS Segal, Y Sautin... - *The American journal ...*, 2007 - Am Soc Nutrition

... Although these epidemiologic associations suggest a potential causal role, are there any direct experimental ... with the consumption of ATP; unlike in glucose metabolism, there is no negative regulatory ... been viewed as simply a potential risk factor for inducing gout, recent studies ...

被引用次数: 603 相关文章 所有 14 个版本 引用

### Relationship of uric acid with progression of kidney disease

M Chonchol, MG Shlipak, R Katz, MJ Sarnak... - ... of *Kidney Diseases*, 2007 - Elsevier

... Any major ECG abnormality, 260 (25), 329 (25), 271 (30), 364 (31), 477 (41), <0.001. ... Cross-Sectional Relationship Between Quintiles of Uric Acid and CKD. ... 95% CI, 0.86 to 1.76) for quintiles 1 through 5, respectively (Table 3). Similar to our previous results, no association was ...

被引用次数: 233 相关文章 所有 7 个版本 引用



学术搜索

找到约 854 条结果 (用时0.14秒)

文章

我的图书馆

时间不限

2016以来

2015以来

2012以来

自定义范围...

按相关性排序

按日期排序

搜索所有网页

中文网页

简体中文网页

☒ 包括专利

☐ 包含引用

创建快讯

小提示: 只搜索中文(简体)结果, 可在 学术搜索设置 指定搜索语言

## Diabetes, hypertension, and cardiovascular disease an update

JR Sowers, M Epstein, ED Frohlich - *Hypertension*, 2001 - Am Heart Assoc

... a major pathway for insulin signal transduction 23 24 25 (Figure 2f). The relationship between the RAS ... 26 This pathway has also been implicated in mediating the activation of NO synthase and ... 19 Finally, the use of ACE inhibitors or Ang II antagonists in any pregnant woman ...

被引用次数: 1231 相关文章 所有 9 个版本 引用 保存

## Essential hypertension, progressive renal disease, and uric acid: a pathogenetic link?

RJ Johnson, MS Segal, T Srinivas, A Ejaz... - *Journal of the ...*, 2005 - Am Soc Nephrol

... Essential Hypertension, Progressive Renal Disease, and Uric Acid: A Pathogenetic Link? ... ESRD is correct for the large number of cases reported (6,7). There is no doubt that ... Younger subjects with short-duration (mean, 3 yr) hypertension also do not show any increase in ...

被引用次数: 238 相关文章 所有 7 个版本 引用 保存

## Potential role of sugar (fructose) in the epidemic of hypertension, obesity and the metabolic syndrome, diabetes, kidney disease, and cardiovascular disease

RJ Johnson, MS Segal, Y Sautin... - *The American journal ...*, 2007 - Am Soc Nutrition

... Although these epidemiologic associations suggest a potential causal role, are there any direct experimental ... with the consumption of ATP; unlike in glucose metabolism, there is no negative regulatory ... been viewed as simply a potential risk factor for inducing gout, recent studies ...

被引用次数: 656 相关文章 所有 17 个版本 引用 保存

## Hyperuricemia in primary and renal hypertension

PJ Cannon, WB Stason, FE Demartini... - *England Journal of ...*, 1966 - Mass Medical Soc

... had elevations of serum uric acid (Table 4). Antihypertensive drug therapy of any variety increased this ... per 100 ml.); there was no statistically significant elevation of the pyruvate level of the ...

[全部](#)[新闻](#)[图片](#)[地图](#)[更多 ▾](#)[搜索工具](#)

找到约 1,140,000 条结果 (用时 1.04 秒)

### Risk of end-stage renal disease associated with gout: a ... - NCBI

[www.ncbi.nlm.nih.gov](http://www.ncbi.nlm.nih.gov) > NCBI > Literature > PubMed Central (PMC) ▾ [翻译此页](#)

作者: KH Yu - 2012 - 被引用次数: 37 - 相关文章

2012年4月18日 - A recent **study based** on data from the National Dialysis Registry of Taiwan ... to a prevalence of 2,288 **ESRD** cases per **population** of 1,000,000 [4]. ... Since **gout** frequently coexists with DM [8] and **hypertension** [9], it is .... Risk for **end-stage renal disease among patients** with **gout**, stratified by age group.

### Management of Chronic Kidney Disease: The Relationship Between ...

<https://www.ncbi.nlm.nih.gov> > NCBI > Literature > PubMed Central (PMC) - [翻译此页](#)

作者: C Mende - 2015 - 被引用次数: 4 - 相关文章

2015年12月9日 - **Chronic kidney disease** (CKD) is increasingly recognized as a global .... CV and renal **disease**, metabolic syndrome, **diabetes**, and **hypertension** [30]. ... In a large **population-based study**, increased SUA levels were shown to be .... **patients** with stage 3 CKD and hyperuricemia with or without **gout** showed ...

### Chronic kidney disease and heart failure—Bidirectional close link and ...

[www.sciencedirect.com/science/article/pii/S0914508710001802](http://www.sciencedirect.com/science/article/pii/S0914508710001802) ▾ [翻译此页](#)

作者: N Shiba - 2011 - 被引用次数: 34 - 相关文章

**Chronic kidney disease** (CKD) is common and the estimated prevalence is ... **result** from **any** structural or functional cardiac disorder and the prevalence is ... More than 40% of HF **patients** have CKD and the close **relationship between** ..... Outcome of heart failure with preserved ejection fraction in a **population-based study**.

### Hyperuricemia, Gout, and Cardiovascular Disease: An Update - Springer

[link.springer.com/article/10.1007/s11926-015-0495-2](http://link.springer.com/article/10.1007/s11926-015-0495-2) - [翻译此页](#)

作者: AM Abeles - 2015 - 被引用次数: 9 - 相关文章

2015年3月5日 **Gout**Hyperuricemia**Hypertension**Cardiovascular diseaseAllerurinal Although the link