

Name of Journal: *World Journal of Nephrology*

ESPS Manuscript NO: 31293

Manuscript Type: Original Article

Retrospective Study

Advanced wasting in peritoneal dialysis patients

Xu Zhi, Glen H Murata, Robert H Glew, Yijuan Sun, Darlene Vigil, Karen S Servilla,
Antonios H Tzamaloukas

Abstract

AIM: To identify ⁵patients with end-stage renal disease (ESRD) treated by peritoneal dialysis (PD) who had zero body fat (BF) as determined by analysis of body composition using ²anthropometric formulas estimating body water (V) and to compare nutritional parameters between these patients and PD patients whose BF was above zero.

Match Overview

1	Crossref 40 words Gupta, N.. "Chronic ocular hypertension induces dend ... e pathology in the lateral geniculate nucleus of the brai	1%
2	Crossref 37 words Antonios H Tzamaloukas, "Estimates of body water, fat-f ree mass, and body fat in patients on peritoneal dialys ...	1%
3	Crossref 21 words Nutrition in Kidney Disease, 2014.	1%
4	Crossref 19 words Schnabel, R.. "Lack of effect of histological lesions on th e phenytoin and phenobarbital concentrations in the b ...	1%
5	Internet 19 words crawled on 24-Mar-2010 www.homedialysis.org	1%
6	Internet 18 words crawled on 17-Jun-2016 spandidos-publications.com	1%

[All](#)[Shopping](#)[News](#)[Images](#)[Videos](#)[More ▾](#)[Search tools](#)

About 720,000 results (0.65 seconds)

Scholarly articles for Advanced Wasting in Peritoneal Dialysis Patients

... of continuous ambulatory **peritoneal dialysis patients**: ... - [Young](#) - Cited by 584

... **wasting** and inflammation in moderate and **advanced** ... - [Kovesdy](#) - Cited by 107

... criteria for protein–energy **wasting** in acute and ... - [Fouque](#) - Cited by 804

Wasting in chronic kidney disease - NCBI - National Institutes of Health

<https://www.ncbi.nlm.nih.gov> › [NCBI](#) › [Literature](#) › [PubMed Central \(PMC\)](#)

by RH Mak - 2011 - Cited by 92 - [Related articles](#)

Mar 16, 2011 - A more recent study showed that higher fat mass in dialysis **patients** Pathophysiology of cachexia/PEW syndrome in **advanced** CKD ... In addition, nutrients may be introduced during **peritoneal dialysis** in CKD **patients**.

Protein energy wasting in chronic kidney disease: An update with ...

<https://www.ncbi.nlm.nih.gov> › [NCBI](#) › [Literature](#) › [PubMed Central \(PMC\)](#)

by YP Jadeja - 2012 - Cited by 31 - [Related articles](#)

Protein-energy **wasting** (PEW) is common in **patients** with chronic kidney disease ... atherosclerosis as predictors of a 10-year mortality in **hemodialysis patients**. **wasting** and inflammation in moderate and **advanced** chronic kidney disease.

Frailty and Protein-Energy Wasting in Elderly Patients with End Stage ...

jasn.asnjournals.org/content/24/3/337.full ▾

by JC Kim - 2013 - Cited by 52 - [Related articles](#)

Frailty and Protein-Energy **Wasting** in Elderly **Patients** with End Stage Kidney Disease ... with **advanced** CKD, including those **patients** undergoing maintenance dialysis ... **patients** undergoing maintenance **hemodialysis** (MHD) or chronic peritoneal ... PEW is found in 18%–75% of maintenance dialysis **patients** in different ...

Counteracting Muscle Wasting in Dialysis Patients - Renal and ...

www.renalandurologynews.com › [Departments](#) › [Commentary](#) › [Expert Reviews](#) ▾

Jan 22, 2010 - The mechanisms leading to muscle **wasting** in dialysis **patients** are still being ... in

[全部](#)[图片](#)[新闻](#)[视频](#)[购物](#)[更多](#)[设置](#)[工具](#)

找到约 919,000 条结果 (用时 0.66 秒)

Google 学术: Advanced wasting in peritoneal dialysis patients

... of continuous ambulatory peritoneal dialysis patients: ... - Young - 被引用次数: 594

... wasting and inflammation in moderate and advanced ... - Kovesdy - 被引用次数: 110

... criteria for protein-energy wasting in acute and ... - Fouque - 被引用次数: 835

Wasting in chronic kidney disease - NCBI - National Institutes of Health

<https://www.ncbi.nlm.nih.gov> > NCBI > Literature > PubMed Central (PMC) - [翻译此页](#)

作者: RH Mak - 2011 - 被引用次数: 98 - [相关文章](#)

2011年3月16日 - A more recent study showed that higher fat mass in dialysis patients Pathophysiology of cachexia/PEW syndrome in advanced CKD ... In addition, nutrients may be introduced during peritoneal dialysis in CKD patients.

Protein energy wasting in chronic kidney disease: An update with ...

<https://www.ncbi.nlm.nih.gov> > NCBI > Literature > PubMed Central (PMC) - [翻译此页](#)

作者: YP Jadeja - 2012 - 被引用次数: 33 - [相关文章](#)

Protein-energy wasting (PEW) is common in patients with chronic kidney disease ... atherosclerosis as predictors of a 10-year mortality in hemodialysis patients. wasting and inflammation in moderate and advanced chronic kidney disease.

Pathophysiology of protein-energy wasting in chronic renal failure.

<https://www.ncbi.nlm.nih.gov/pubmed/9915908> - [翻译此页](#)

全部

图片

新闻

视频

购物

更多

设置

工具

找到约 901,000 条结果 (用时 0.57 秒)

Google 学术: Advanced wasting in peritoneal dialysis patients

... of continuous ambulatory peritoneal dialysis patients: ... - Young - 被引用次数: 595

... wasting and inflammation in moderate and advanced ... - Kovesdy - 被引用次数: 112

... criteria for protein-energy wasting in acute and ... - Fouque - 被引用次数: 848

Counteracting Muscle Wasting in Dialysis Patients - Renal and ...

www.renalandurologynews.com > ... > Commentary > Expert Reviews ▾ 翻译此页

2010年1月22日 - The mechanisms leading to muscle wasting in dialysis patients are still being ... even in patients with advanced CKD, nitrogen balance studies show a ... is detectable in maintenance hemodialysis patients—both obese and

Recent advances in the pathophysiology and management of protein ...

<https://rrtjournal.biomedcentral.com/articles/10.1186/s41100-016-0015-5> ▾ 翻译此页

作者: K Nitta - 2016 - 被引用次数: 1 - 相关文章

2016年1月30日 - Malnutrition Protein-energy wasting Mortality Chronic kidney disease similar between healthy participants and hemodialysis patients [96].

Wasting in chronic kidney disease - NCBI - National Institutes of Health

<https://www.ncbi.nlm.nih.gov> > NCBI > Literature > PubMed Central (PMC) - 翻译此页

作者: RH Mak - 2011 - 被引用次数: 100 - 相关文章

2011年3月16日 - A more recent study showed that higher fat mass in dialysis patients Pathophysiology of cachexia/PEW syndrome in advanced CKD ... In addition, nutrients may be introduced during peritoneal dialysis in CKD patients.

Protein-Energy Wasting and Mortality in Chronic Kidney Disease - NCBI

<https://www.ncbi.nlm.nih.gov> > NCBI > Literature > PubMed Central (PMC) - 翻译此页