



26935-Review

 Quotes Excluded
 Bibliography Excluded

 14%
 SIMILAR

Name of Journal: *World Journal of Methodology*
ESPS Manuscript NO: 26935

Manuscript Type: Therapeutics Advances

Modified skin bridge technique for ilio-inguinal lymph node dissection: A forgotten technique revisited
Mukur D Ray, Pankaj K Garg, Ashish Jakhetiya, Sunil Kumar, Durgatosh Pandey

Abstract

Ilio-inguinal lymph node dissection (IILD) is a commonly performed surgical procedure for a number of malignant conditions involving mainly the male and female genitalia, and skin; however the perioperative morbidity of IILD, due to high frequency of flap necrosis, wound infection and seroma formation, has always been a major concern for the surgeons. The aim of the paper is to propose a modified skin bridge technique of IILD using two parallel curvilinear incisions to minimize perioperative morbidity, classically associated with the

Match Overview

1	CrossCheck 78 words &NA, "15th Biennial Meeting of the International Gynecologic Cancer Society", International Journal of Gynecological	7%
2	CrossCheck 17 words R. RAVI. British Journal of Urology, 12/1993	1%
3	Internet 14 words crawled on 19-Jun-2011 www.humangrossanatomy.us	1%
4	CrossCheck 14 words John Spratt. "Groin dissection", Journal of Surgical Oncology, 04/2000	1%
5	CrossCheck 10 words Skandalakis, Lee J., and John E. Skandalakis. "Abdominal Wall and Hernias", Surgical Anatomy and Technique, 201	1%
6	CrossCheck 10 words Wolfla, Christopher E., and Daniel K. Resnick. "62 Surgical Treatment of Meralgia Paresthetica", Spine and Peripheral	1%

[全部](#)[视频](#)[图片](#)[新闻](#)[更多 ▾](#)[搜索工具](#)

找到约 1,940 条结果 (用时 0.86 秒)

您是不是要找: **Modified skin bridge technique for
ILIOINGUINAL LYMPH NODE DISSECTION: A forgotten
technique revisited**

[PDF] **Sentinel Lymph Node Biopsy for Melanoma and Surgical**

cdn.intechopen.com/pdfs-wm/42185.pdf ▾ [翻译此页](#)

作者: Y Nakamura - 2013 - 被引用次数: 2 - [相关文章](#)

and therapeutic **lymph node dissection** (TLND) only when clinical nodal disease is present. The **technique** of lymphatic mapping and sentinel **lymph node** biopsy (SLNB). 16 patients), whereas for those who underwent **ilioinguinal** LND, it was 0% (0 of 25 patients) ... sion below and a **skin bridge** between[73, 84, 100].

[PDF] **download the pdf version - Lymphology2013**

www.lymphology2013.com/Abstract_Book/files/abstract_book.pdf ▾

2013年9月16日 - New functional lymphatic imaging **techniques** involving the use of SLN biopsy may avoid a morbid bilateral radical **ilioinguinal** lymph ... However, the extent of **lymph node dissection** remains the same. tissue fluid and lymph, location sites of edema fluid in **skin**, subcutis and Sant'Angelo **Bridge** ...

[PDF] **Gastrointestinal Radiology - RSNA 2014**

rsna2014.rsna.org/pdf/Gastrointestinal%20Radiology.pdf ▾

Normal anatomy of bowel wall Sonographic **techniques** Review of imaging findings vascular anatomy (2) Tumor location (3) **Lymph node dissection** level (4) Abdominal and **Pelvic** Imaging of Complications to Systemic Cancer Therapy system, GI and GU tract, and **skin** Inflammation of small blood vessels of the gut ...

[PDF] 2015 - American College of Physicians

全部

视频

图片

新闻

更多 ▾

搜索工具

找到约 1,890 条结果 (用时 0.66 秒)

您是不是要找: **Modified skin bridge technique for *ilioinguinal* lymph node dissection: A forgotten technique revisited**

[PDF] **download the pdf version - Lymphology2013**

www.lymphology2013.com/Abstract_Book/files/abstract_book.pdf ▾

2013年9月16日 - New functional lymphatic imaging **techniques** involving the use of SLN biopsy may avoid a morbid bilateral radical **ilioinguinal** lymph ... However, the extent of **lymph node dissection** remains the same. tissue fluid and lymph, location sites of edema fluid in **skin**, subcutis and Sant'Angelo **Bridge**

[PDF] **Annual Report 1988 - NKI**

research.nki.nl/library/jaarverslagen/AR1988.pdf ▾

this **technique** to plastic embedded sections, a method which preserves the However, the binding sites on **lymph node** cells of one patient were saturated af ...

[PDF] **2015 - American College of Physicians**

https://www.acponline.org/system/files/documents/about_acp/chapters/.../salud_15.pdf

2015年11月13日 - PGY-2 The **Forgotten** Disease: Lemierre's Syndrome. 93. CR Mohit. Pahuja **lymph node** revealed classic HL with mixed cellularity. Currently ...

[PDF] **contents - American College of Physicians**

<https://www.acponline.org/system/files/documents/.../residents/.../rf-winners-2016.pdf>

Does mammographic density **modify** the characteristics of the breast cancer Diffuse Papulonodular **Skin** Lesions in a Renal Transplant Patient ... exists on the integration of bedside ultrasound **techniques** into resuscitation of cardiac arrest. for diagnostic laparoscopy and mesenteric and right **inguinal lymph node** ...

[全部](#)[视频](#)[图片](#)[新闻](#)[更多 ▾](#)[搜索工具](#)

找到约 1,820 条结果 (用时 0.90 秒)

Modified technique of radical inguinal lymphadenectomy for ... - NCBI

www.ncbi.nlm.nih.gov/pubmed/20620415 ▾ [翻译此页](#)

作者: K Yao - 2010 - 被引用次数: 37 - 相关文章

2010年6月17日 - Modified technique of radical inguinal lymphadenectomy for penile carcinoma: ...

Modified inguinal lymphadenectomy has been used to decrease the ... wound infection (1.4%), skin necrosis (4.7%), lymphedema (13.9%), ...

缺少字词: bridge ilio forgotten revisited

Modified radical inguinal lymphadenectomy for carcinoma of the penis ...

www.ncbi.nlm.nih.gov/pubmed/12629358 ▾ [翻译此页](#)

作者: U Jacobellis - 2003 - 被引用次数: 66 - 相关文章

Modified radical inguinal lymphadenectomy for carcinoma of the penis; technique and results. ... with leiomyosarcoma of the penis underwent bilateral modified inguinal lymphadenectomy, including removal of the ... RESULTS: At a followup of 6 to 104 months no skin necrosis, infection or deep venous thrombosis occurred.

缺少字词: bridge ilio forgotten revisited

[PDF] Modified Technique of Radical Inguinal Lymphadenectomy for Penile ...

www.urologicaltumor.com/uploadfiles/file/.../20121031021708_73377.pdf ▾ [翻译此页](#)

作者: K Yao - 1842 - 被引用次数: 37 - 相关文章

2012年10月31日 - Modified inguinal lymphadenectomy has been used to decrease the complication ... we modified the technique based on the boundaries Skin flaps were separated in plane. c, tissues between superficial layer of Camper's.

缺少字词: bridge forgotten revisited

[PDF] download the pdf version - Lymphology2013

www.lymphology2013.com/Abstract_Book/files/abstract_book.pdf ▾

2013年9月16日 - New functional lymphatic imaging techniques involving the use of SLN biopsy may