

BIostatISTICS STATEMENT

Name of Journal: *World Journal of Gastrointestinal Oncology*

Manuscript No: 39357

Manuscript Type: Retrospective study

IS C-REACTIVE PROTEIN A PROGNOSTIC FACTOR FOR THE WHOLE GEP-NET GROUP?

Ömer KOMA, Göksel BENİ, Özgül SAĞOL, Mesut AKARSU

Correspondence to: Göksel Bengi, MD, PhD, Assis. Prof., Department of Gastroenterology, Dokuz Eylül University Hospital, Inciraltı, 35000, İzmir, Turkey. goksel.bengi@deu.edu.tr

Telephone: +90-532-4626972 +90-232-4123713

Fax: +90-232-4123799

Biostatistics statement:

Statistical analysis was performed using the SPSS 19.0 for Windows (SPSS, Inc.; Chicago, USA) package program. Descriptive values were given as number (n), percentage (%), mean, standard deviation (SD) and median. Pearson chi-square test was used for the comparison of categorical variables. Continuous variables were compared with non-parametric tests (Mann-Whitney U test and Kruskal-Wallis test) because they did not have a normal distribution when evaluated for normality of distribution with Kolmogorov-Smirnov and Shapiro-Wilk tests. Level of statistical significance was $p < 0.05$.

Name: Göksel Bengi

Signature



20/06/2018