

Biostatistics statement

Name of Journal: World Journal of Gastrointestinal Endoscopy

ESPS Manuscript NO: 24729

Title: Prior minimal endoscopic sphincterotomy to prevent pancreatitis related to endoscopic balloon sphincteroplasty

Author List: Ryo Kanazawa,M.D., Jin Kan Sai,M.D., Tomoyasu Ito,M.D., Hiroko Miura,M.D.,Shigeto Ishii,M.D., Hiroaki Saito,M.D., Ko Tomishima,M.D., Ryo Shimizu,M.D., Koki Sato,M.D., Manabu Hayashi,M.D., Sumio Watanabe,M.D., Shuichiro Shiina,M.D.

Correspondence to: Jin Kan Sai, MD

In this manuscript, statistical analysis was performed with SPSS version 17.0. Data were compared using paired t-test. Mann-Whitney *U* test was used for comparing continuous data with skewed distribution in the two groups. A Chi-square test with Yate's correction was used to analyze gender. The difference in serum amylase level after the procedure and the incidence of post-procedure pancreatitis were compared using Wilcoxon signed-rank test.

Here I confirm that the statistical analyses of this study are performed in an appropriate manner. If you have any question concerning the statistical analyses, please do not hesitate to contact me.

Jin Kan Sai, MD,

Associate Professor

Department of Gastroenterology,

Juntendo University

2-1-1 Hongo, Bunkyo-ku, Tokyo 113-8421, Japan.

Telephone: +81-3-5802-1061; **Fax:** +81-3-5684-5960

E-mail: jinkans@juntendo.ac.jp