

70186_Auto_Edited.docx

2

Name of Journal: *World Journal of Clinical Cases*

Manuscript NO: 70186

Manuscript Type: CASE REPORT

Primary central nervous lymphoma presenting as a single choroidal lesion mimicking metastasis : A case report

Jang HR *et al.* Primary central nervous system lymphoma

Hee Ryeong Jang, Kyu-Hyoung Lim, Kyungyul Lee

Hee Ryeong Jang, Kyu-Hyoung Lim, Internal Medicine, Kangwon National University School of Medicine, Chuncheon 24289, South Korea

Kyungyul Lee, Department of Pathology, Kangwon National University School of Medicine, Chuncheon 24289, South Korea

1

Author contributions: All authors contributed to the study conception and design. Material preparation and data collection and analysis were performed by Kyu-Hyoung Lim, Hee Ryeong Jang, and Kyung Yul Lee. The first draft of the manuscript was written by Hee Ryeong Jang, and all authors commented on the previous versions of the manuscript; all authors read and approved the final manuscript.

Corresponding author: **Kyu-Hyoung Lim**, Doctor, MD, PhD, Associate Professor, Internal Medicine, Kangwon National University School of Medicine, 156 Baengnyeong-ro, Chuncheon-si, Gangwon-do 24289, Republic of Korea, Chuncheon 24289, South Korea. kyuhyoung.lim@gmail.com

Received: July 26, 2021

Revised: November 4, 2021

Accepted:

Published online:

Abstract

BACKGROUND

Primary choroidal lymphoma is usually an indolent B-cell lymphoma and rarely progresses to extraocular sites. Herein, we report a case of primary choroidal lymphoma diagnosed as diffuse large B-cell lymphoma (DLBL), which progressed to the brain parenchyma after 4 mo.

CASE SUMMARY

A 78-year-old man presented with diminution of vision in his right eye. A choroidal lesion suspected of metastatic lesion was observed in the right eye by ophthalmologic examination. To discover the primary tumor, imaging investigations were performed but no malignant lesion was detected. After 4 mo, the patient returned to the clinic presenting with neurological symptoms. Brain magnetic resonance imaging revealed an abnormal contrast-enhancing mass in the left cerebellum. A stereotactic biopsy was performed, and DLBL was confirmed. The patient received the high dose methotrexate-based chemotherapy and he achieved complete remission.

CONCLUSION

Primary choroidal lymphoma is usually known to have a benign clinical course without systemic involvement. We present a rare case of primary choroidal lymphoma diagnosed as DLBL that progressed to the brain parenchyma within months.

Key Words: Primary choroidal lymphoma; Diffuse large B-cell lymphoma; Primary central nervous system lymphoma; Case report

Jang HR, Lim KH, Lee K. Primary central nervous lymphoma presenting as a single choroidal lesion mimicking metastasis : A case report . *World J Clin Cases* 2021; In press

Core Tip: Primary choroidal lymphoma is a rare subset of primary intraocular lymphoma and shows a benign clinical course with no systemic involvement. Our case report involves a primary choroidal lymphoma demonstrating rare extraocular progression within months.

6

INTRODUCTION

Primary intraocular lymphoma (PIOL) is a rare subset of primary central nervous system lymphoma (PCNSL) and includes vitreoretinal, choroidal, and iridal lymphomas^[1]. Among PIOLs, primary choroidal lymphoma is a very rare disease, and only a small number of cases have been reported in the literature. Previously reported primary choroidal lymphomas were usually low-grade B-cell lymphomas, which rarely progressed to the central nervous system^[2,3]. Herein, we report a very rare case of primary choroidal lymphoma diagnosed as diffuse large B-cell lymphoma (DLBL), which initially presented as a unilateral choroidal lesion mimicking metastasis.

11

CASE PRESENTATION

Chief complaints

A 78-year-old male came to the hospital presenting with a diminution of vision in his right eye.

History of present illness

His blurred vision had been aggravated for several weeks.

History of past illness

The patient had no relevant medical history, such as that of diabetes mellitus or hypertension. In the past, the patient had undergone short-term treatment for occupational pneumoconiosis resulting from his work as a coal miner for > 10 years.

9

Personal and family history

No special history of personal and family.

Physical examination

An ophthalmological examination revealed exudative subretinal detachment with a choroidal lesion suspicious of metastasis in the right eye (Figure 1A).

Laboratory examinations

Lactate dehydrogenase was moderately elevated at 387 U/L (normal range: 119–229 U/L).

Imaging examinations

To identify the primary tumor, the patient was referred to the Department of Hemato-oncology at our institute. Brain ¹³magnetic resonance imaging (MRI) and computed tomography (CT) of the neck, chest, and abdomen-pelvis showed no malignant lesions.

FINAL DIAGNOSIS

Four months later, the patient revisited the clinic because of dysarthria, headache, and right-sided weakness. No apparent aggravation of visual acuity was observed. The same findings of ophthalmoscopy as at the first visit were confirmed (Figure 2B).

Brain MRI showed a single, irregular, contrast-enhancing mass in the left cerebellum (Figure 2A). The patient underwent a stereotactic biopsy of the mass and was diagnosed with DLBL (Figure 3). Further imaging, including CT and positron emission tomography/CT, revealed no systemic involvement. There was no evidence of lymphoma involvement in the cerebrospinal fluid and bone marrow.

TREATMENT

The patient received high-dose methotrexate (HD-MTX)-based chemotherapy, which comprised methotrexate (3.5 g/m²), vincristine (1.4 mg/m², capped at 2 mg), and prednisolone (100 mg/d), with the addition of procarbazine (100 mg/m²) in the first, third, and fifth cycles. Intrathecal injection was not given in this patient because there was no evidence of lymphoma involvement in the CSF test before and after the first cycle of HD-MTX based chemotherapy.

OUTCOME AND FOLLOW-UP

After completing six cycles of HD-MTX-based chemotherapy, the patient's neurological symptoms, such as right-sided weakness, dysarthria, and blurred vision, markedly improved. The abnormal, contrast-enhancing mass in the left cerebellum disappeared on brain MRI (Figure 2B), and the creamy yellow, subretinal infiltrates were markedly decreased (Figure 1C); these findings were compatible with complete remission. Maintenance therapy was recommended, but the patient and caregiver refused the treatment due to concerns about the side effects of additional chemotherapy and radiotherapy. The patient and his main caregiver were seriously concerned about the risk of cognitive decline that can be caused by additional treatments. Complete remission was maintained for approximately 17 mo without the maintenance chemotherapy.

DISCUSSION

Intraocular lymphoma is extremely rare and accounts for approximately 1.86% of intraocular malignancies^[4]. Intraocular lymphoma is a heterogeneous group of malignancies located in different tissues, including the vitreous, retina, choroid, ciliary body, and iris, within the eye. It also refers to forms of primary or secondary to central nervous system lymphoma or disseminated systemic disease^[3, 5].

Primary intraocular lymphoma is considered a subset of PCNSL, and it progresses to the central nervous system in 15%-25% of PCNSL cases^[3]. Primary vitreoretinal lymphoma is the most common intraocular lymphoma, followed by uveal lymphoma.⁴

Choroidal lymphoma is a subset of uveal lymphoma. It can be subdivided into primary and secondary lymphoma based on the presence of systemic lymphoma at the time of ocular presentation. Primary choroidal lymphomas are defined as the absence of prior systemic lymphomas or concurrent extraocular lymphomas^[6]. Several studies have reported clinical differences between primary and secondary choroidal lymphomas^[2,7].

⁸ Primary choroidal lymphomas are mainly low grade B-cell lymphomas such as extranodal marginal zone B-cell lymphoma, are usually unilateral, and typically do not progress to the central nervous system parenchyma. Secondary choroidal lymphoma is characterized by the presence of previously known cancer or concurrent systemic lymphomas at the initial ocular presentation. In contrast to primary choroidal lymphomas, secondary choroidal lymphomas are more likely to demonstrate bilateral involvement and preexistent lymphomas. More than half of secondary choroidal lymphomas have been confirmed as high-grade B-cell lymphomas, such as DLBL^[2,8].

Unlike the previously reported cases of primary choroidal lymphoma, this case was characterized by the pathological findings of DLBL and disease progression to the brain parenchyma within a few months. In most of the previous cases of primary choroidal lymphoma, management involved local treatment or observation, whereas in our case, HD-MTX-based chemotherapy was administered, and the treatment response was complete remission.

CONCLUSION

Primary choroidal lymphoma is generally known to have a benign clinical course without systemic involvement. We reported a rare case of primary choroidal lymphoma diagnosed as DLBL, characterized by an aggressive clinical course that progressed to the brain parenchyma within a few months.

ACKNOWLEDGEMENTS

The authors thank the Department of Radiology, Kangwon National University School of Medicine.

REFERENCES

- 1 **Tang LJ**, Gu CL, Zhang P. Intraocular lymphoma. *Int J Ophthalmol* 2017; **10**: 1301-1307 [PMID: 28861359 DOI: 10.18240/ijo.2017.08.19]
- 2 **Mashayekhi A**, Shukla SY, Shields JA, Shields CL. Choroidal lymphoma: clinical features and association with systemic lymphoma. *Ophthalmology* 2014; **121**: 342-351 [PMID: 23978622 DOI: 10.1016/j.opthta.2013.06.046]
- 3 **Chan CC**, Rubenstein JL, Coupland SE, Davis JL, Harbour JW, Johnston PB, Cassoux N, Touitou V, Smith JR, Batchelor TT, Pulido JS. Primary vitreoretinal lymphoma: a report from an International Primary Central Nervous System Lymphoma Collaborative Group symposium. *Oncologist* 2011; **16**: 1589-1599 [PMID: 22045784 DOI: 10.1634/theoncologist.2011-0210]
- 4 **Reddy EK**, Bhatia P, Evans RG. Primary orbital lymphomas. *Int J Radiat Oncol Biol Phys* 1988; **15**: 1239-1241 [PMID: 3053542 DOI: 10.1016/0360-3016(88)90210-6]
- 5 **Konstantinidis L**, Damato B. Intraocular Metastases--A Review. *Asia Pac J Ophthalmol (Phila)* 2017; **6**: 208-214 [PMID: 28399345 DOI: 10.22608/APO.201712]
- 6 **Coupland SE**, Foss HD, Hidayat AA, Cockerham GC, Hummel M, Stein H. Extranodal marginal zone B cell lymphomas of the uvea: an analysis of 13 cases. *J Pathol* 2002; **197**: 333-340 [PMID: 12115879 DOI: 10.1002/path.1130]
- 7 **Coupland SE**, Damato B. Understanding intraocular lymphomas. *Clin Exp Ophthalmol* 2008; **36**: 564-578 [PMID: 18954321 DOI: 10.1111/j.1442-9071.2008.01843.x]
- 8 **Doycheva D**, Zierhut M, Süsskind D, Bartz-Schmidt KU, Deuter C. [Diagnostics and treatment of choroidal lymphoma]. *Ophthalmologe* 2015; **112**: 217-222 [PMID: 25693876 DOI: 10.1007/s00347-014-3206-x]

Footnotes

Informed consent statement: The need for patient consent for this report was waived by the institutional review board.

Conflict-of-interest statement: Kyu-Hyoung Lim, Hee Ryeong Jang, and Kyung Yul Lee declare that there is no conflict of interest.

CARE Checklist (2016) statement: The authors have read the CARE Checklist (2016), and the manuscript was prepared and revised according to the CARE Checklist (2016).

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, ¹ which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <https://creativecommons.org/licenses/by-nc/4.0/>

¹⁰ **Provenance and peer review:** Unsolicited Manuscript; Externally peer reviewed.

Peer-review model: Single blind

Peer-review started: July 26, 2021

First decision: October 22, 2021

² **Article in press:**

Specialty type: Oncology

Country/Territory of origin: South Korea

Peer-review report's scientific quality classification

Grade A (Excellent): 0

Grade B (Very good): 0

Grade C (Good): C

Grade D (Fair): D

Grade E (Poor): 0

P-Reviewer: Dou AX, Montemurro N **S-Editor:** Xing YX **L-Editor:** **P-Editor:**

Figure Legends




Figure 1 Course of ophthalmoscopy examinations. A: Image showing multifocal, creamy yellow, subretinal infiltrates during the first clinical visit; B: Image showing the spread of the creamy yellow, subretinal infiltrates toward the mid-periphery of the fundus. The image was taken 4 mo after the first clinical visit; C: Subretinal infiltrates appear to be markedly decreased after six cycles of systemic chemotherapy.




Figure 2 Findings of brain magnetic resonance imaging. A: Image showing no evidence of space-occupying intracranial or diffusion-restricted lesions. The image was taken at the first clinical visit; B: Image showing an irregular contrast-enhancing lesion, measuring approximately 3.8 cm × 3.8 cm in size, with diffusion restriction in the left cerebellum. The image was taken 4 mo after the first clinical visit; C: Image showing no evidence of abnormal contrast-enhancing lesion in the cerebellum after six cycles of systemic chemotherapy.




Figure 3 Representative microscopic images of diffuse large B-cell lymphoma in the brain. A: Infiltration of large, atypical cells with prominent nucleoli around the granular layer of the cerebellum (hematoxylin and eosin stain, magnification: 200×); B: Infiltration of large, atypical cells with prominent nucleoli around the granular layer of the cerebellum (hematoxylin and eosin stain, magnification: 400×); C: Tumor cells show strong positivity for the B-cell surface marker CD20 on immunohistochemistry (Immunohistochemical stain, magnification: 200×).

16%

SIMILARITY INDEX

PRIMARY SOURCES

1	www.wjgnet.com Internet	63 words — 4%
2	f6publishing.blob.core.windows.net Internet	35 words — 2%
3	link.springer.com Internet	29 words — 2%
4	Arman Mashayekhi, Shripaad Y. Shukla, Jerry A. Shields, Carol L. Shields. "Choroidal Lymphoma", Ophthalmology, 2014 Crossref	17 words — 1%
5	www.cambridge.org Internet	17 words — 1%
6	Shouvik Saha, Elizabeth M. Graham. "Chapter 10 Masquerade Syndromes", Springer Science and Business Media LLC, 2009 Crossref	15 words — 1%
7	Catherine Tang, Julian Lindsay, Anthony Gill, Ian Kerridge. "Defibrotide Use in Vincristine-induced Hepatic Sinusoidal Obstruction Syndrome", Clinical Lymphoma Myeloma and Leukemia, 2017 Crossref	14 words — 1%

- 9 Hyeok Gyu Kwon, Jeong-Hee Yang, Jee-Hyun Kwon, Dongseok Yang. "Association between scrub typhus encephalitis and diffusion tensor tractography detection of Papez circuit injury: A case report", World Journal of Clinical Cases, 2021

13 words — 1%

Crossref

- 10 Yuka Ikeda, Nozomi Nagase, Ai Tsuji, Yasuko Kitagishi, Satoru Matsuda. " Neuroprotection by dipeptidyl-peptidase-4 inhibitors and glucagon-like peptide-1 analogs the modulation of AKT-signaling pathway in Alzheimer's disease ", World Journal of Biological Chemistry, 2021

13 words — 1%

Crossref

- 11 Jia-Xi Tang, Ling Wang, Wei-Qi Nian, Wan-Yan Tang, Jing-Yu Xiao, Xi-Xi Tang, Hong-Liang Liu. "Asymptomatic carbon dioxide embolism during transoral vestibular thyroidectomy: A case report", World Journal of Clinical Cases, 2021

12 words — 1%

Crossref

- 12 e-tjo.org

Internet

12 words — 1%

- 13 eprints.lib.hokudai.ac.jp

Internet

12 words — 1%