



Meta-analysis of outcomes from drug-eluting stent implantation in infrapopliteal arteries

Ming-Xuan Li, Hai-Xia Tu, Meng-Chen Yin

Specialty type: Peripheral vascular disease

Provenance and peer review: Unsolicited article; Externally peer reviewed.

Peer-review model: Single blind

Peer-review report's scientific quality classification

Grade A (Excellent): 0
Grade B (Very good): B
Grade C (Good): C
Grade D (Fair): 0
Grade E (Poor): 0

P-Reviewer: Chow WK, Taiwan; Ueda H, Japan

Received: May 5, 2023

Peer-review started: May 5, 2023

First decision: July 3, 2023

Revised: July 7, 2023

Accepted: July 17, 2023

Article in press: July 17, 2023

Published online: August 6, 2023



Ming-Xuan Li, Hai-Xia Tu, Meng-Chen Yin, Department of Vascular Surgery, Beijing Fengtai You'anmen Hospital, Beijing 100069, China

Corresponding author: Hai-Xia Tu, MM, Chief Physician, Department of Vascular Surgery, Beijing Fengtai You'anmen Hospital, No. 199 You'anmenwai Street, Beijing 100069, China. haixia_tu2023@163.com

Abstract

BACKGROUND

Percutaneous drug-eluting stent implantation (DESI) is an emerging and promising treatment modality for infrapopliteal artery diseases (IPADs). This systematic review and meta-analysis summarizes and quantitatively analyzes the outcomes of DESI in IPADs considering the hazard ratio (HR), which is a more accurate and appropriate outcome measure than the more commonly used relative risk and odds ratio.

AIM

To explore the superiority of drug-eluting stents (DESs) *vs* traditional treatment modalities for IPADs.

METHODS

The following postoperative indicators were the outcomes of interest: All-cause death (ACD)-free survival, major amputation (MA)-free survival, target lesion revascularization (TLR)-free survival, adverse event (AE)-free survival, and primary patency (PP) survival. The outcome measures were then compared according to their respective HRs with 95% confidence intervals (CIs). The participants were human IPAD patients who underwent treatments for infrapopliteal lesions. DESI was set as the intervention arm, and traditional percutaneous transluminal angioplasty (PTA) with or without bare metal stent implantation (BMSI) was set as the control arm. A systematic search in the Excerpta Medica Database (Embase), PubMed, Web of Science, and Cochrane Library was performed on November 29, 2022. All controlled studies published in English with sufficient data on outcomes of interest for extraction or conversion were included. When studies did not directly report the HRs but gave a corresponding survival curve, we utilized Engauge Digitizer software and standard formulas to convert the information and derive HRs. Then, meta-analyses were conducted using a random-effects model.

RESULTS

Five randomized controlled trials and three cohort studies involving 2639 participants were included. The ACD-free and MA-free survival HR values for DESI were not statistically significant from those of the control treatment ($P > 0.05$); however, the HR values for TLR-free, AE-free, and PP-survival differed significantly [2.65 (95% CI: 1.56-4.50), 1.57 (95% CI: 1.23-2.01), and 5.67 (95% CI: 3.56-9.03), respectively].

CONCLUSION

Compared with traditional treatment modalities (*i.e.*, PTA with or without BMSI), DESI for IPADs is superior in avoiding TLR and AEs and maintaining PP but shows no superiority or inferiority in avoiding ACD and MA.

Key Words: Infrapopliteal; Drug-eluting stent; Below-the-knee; Meta-analysis; Hazard ratio

©The Author(s) 2023. Published by Baishideng Publishing Group Inc. All rights reserved.

Core Tip: The utility of drug-eluting stents (DESs) for infrapopliteal artery diseases was explored using traditional percutaneous transluminal angioplasty with or without bare stent implantation as control. The results suggest that the DES is superior on multiple outcomes. The hazard ratio, which is most appropriate for various outcomes categorized as time-to-event data by type, was adopted as the outcome measure, rather than the relative risk or the odds ratio.

Citation: Li MX, Tu HX, Yin MC. Meta-analysis of outcomes from drug-eluting stent implantation in infrapopliteal arteries. *World J Clin Cases* 2023; 11(22): 5273-5287

URL: <https://www.wjgnet.com/2307-8960/full/v11/i22/5273.htm>

DOI: <https://dx.doi.org/10.12998/wjcc.v11.i22.5273>

INTRODUCTION

Whether accompanied by femoropopliteal inflow disease or not, infrapopliteal artery disease (IPAD) is the primary cause of critical limb ischemia (CLI)[1,2]. Femoropopliteal-to-distal bypass surgery is considered the traditional treatment option for revascularization in IPADs[3-5]. Over the past few decades, minimally invasive percutaneous transluminal angioplasty (PTA) (*i.e.*, balloon dilatation) with or without bare metal stent implantation (BMSI) has been widely used, especially for patients with physical conditions that make it difficult for them to withstand open surgery or those lacking suitable distal arteries for bypass[6-10]. However, although this modality has a satisfactory technical success rate, it is associated with a significantly high risk of clinical failure due to lesion restenosis, even in the short term[11-13].

Drug-eluting stents (DESs) have demonstrated success in coronary artery diseases and have been widely demonstrated to maintain longer patency in femoropopliteal artery disease[14-16]. DES implantation (DESI) for IPADs was introduced over ten years ago[17-19]. Fusaro *et al*[20] reported the odds ratios (ORs) of some outcomes of DESI for IPADs *vs* control treatments in a meta-analysis published in 2013. Liu *et al*[21] reported the relative risks (RRs) in another meta-analysis published in 2017. However, similar to other intravascular therapies, almost all outcomes of concern after DESI are time-to-event data[22]; thus, the incidences of these outcomes will change significantly over time. In the PADI trial[23], the cumulative mortality at the one-year follow-up was 23.3% for DESI in IPADs and 62.3%[24] and 80.8%[25] at the 5- and 10-year follow-ups, respectively. Gratifyingly, an increasing number of clinical trials[24-27] have reported the outcomes of DESI for IPADs using the hazard ratio (HR), which is more appropriate for analyzing time-to-event data[28]. This makes it possible for us to perform a meta-analysis using this outcome measure.

MATERIALS AND METHODS

Study protocol

This review was registered with PROSPERO (CRD42022377456) in accordance with the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) framework[29]. All data analyses were based on original studies; thus, no additional ethical approvals or participant consent forms were needed.

Search strategy

The Excerpta Medica, PubMed, Web of Science, and Cochrane Library databases were searched on November 29, 2022. We searched without date limits for all relevant articles using “eluting”, “stent”, “limb” and all possible synonyms. All entry terms and search commands can be found in Supplementary material.

Study selection

All references were imported into Endnote X9 for duplicate removal and a brief information review. Then, the full texts of all available articles that passed the preliminary screening were downloaded and read to identify those that could be

included in the final study. At this stage, the bibliographies and citations of the related articles were also screened for other potential articles.

We defined the HRs (DESI *vs* control treatments) with 95% confidence intervals (CIs) of the following five outcomes of IPADs as the primary outcome measures of interest: all-cause death (ACD)-free survival, major amputation (MA)-free survival, target lesion revascularization (TLR)-free survival, adverse event (AE)-free survival, and primary patency (PP) survival. Studies that simultaneously met the following criteria were included: 1) the study design was a randomized controlled trial (RCT) or cohort study; 2) the language of publication was English; 3) the target lesions of the human participants who underwent interventions were in infrapopliteal arteries demonstrating IPAD; 4) the number of participants in each arm was no less than 10; 5) DESI was conducted in one of the arms; and 6) at least one of the primary outcome measures of interest was directly reported, or at least one of the survival curves was provided. IPAD was defined as a disease caused by intraluminal atherosclerotic stenosis or occlusion of the popliteal artery below the tibial plateau level, anterior tibial artery, tibioperoneal trunk artery, posterior tibial artery, or peroneal artery. We did not limit the definition of the above five outcomes. Studies that did not meet the above criteria or only included complete duplicates of the outcome data available for extraction were excluded.

Two authors (Li MX and Tu HX) independently performed the search, title abstract filtering, and full-text review based on the above selection criteria. Any discrepancies were resolved by consensus.

Data extraction

After identifying the studies for inclusion, we extracted the basic study and participant characteristics and the primary outcome measures. Directly reported adjusted HRs derived from multivariate Cox proportional hazard models were preferentially adopted and extracted. If HRs were absent but survival curves were present, we used Engauge Digitizer 11.3, an open-source software that can extract digital data from a graph, to transform the information in the curves and calculate the HRs[28,30,31]. Tierney *et al*[28] comprehensively summarized relevant statistical theories and provided an HR calculation spreadsheet (Excel format) with a preset calculation formula. We used this spreadsheet to calculate HRs instead of a manual calculation process. Data extraction was performed by a pair of independent authors (Li MX and Tu HX). Any discrepancies were resolved by consensus.

Risk of bias assessment

We assessed the bias risk in the RCTs using the Cochrane Risk of Bias 2.0 (RoB 2.0) tool[32]. This tool evaluates 5 domains: The randomization process, deviations from intended interventions, missing outcome data, outcome measurements, and reported result selection. The risk for each of the 5 domains and the overall risk is described as low, some concerns, or high. We assessed the risks of bias in cohort studies using the Cochrane Risk Of Bias In Nonrandomized Studies-of Interventions (ROBINS-I) tool[33]. This tool evaluates 7 domains: Confounding, participant selection, intervention classification, deviations from intended interventions, missing outcome data, outcome measurements, and reported result selection. The level of bias risk in each main study and overall was divided into five levels: low, moderate, serious, critical, and unclear. The highest risk level among all domains was adopted as the overall assessment result separately for each tool.

The risk of bias assessment was performed by a pair of independent authors (Li MX and Yin MC). When the two authors had different opinions on a certain assessment result, the worst opinion was adopted.

Statistical analysis

Stata (Stata Corp, Texas, United States) version 16.0 was used for all statistical analyses. In each meta-analysis, we took the natural logarithms of the extracted HR value and the maximum and minimum 95%CI values per study and then included the three obtained variables in the "metan" command. To reduce error, a random-effects model rather than a fixed-effects model was used, regardless of the degree of heterogeneity among studies[34]. We corrected the degrees of freedom by restricted maximum likelihood estimation[35]. The calculation of the effect size (ES), *i.e.*, the pooled HR, of different study design types (RCTs or cohort studies) on individual outcomes was performed separately.

The formulas for the Cox proportional hazard model[36] (1) and the meta-analysis based on extracted or transformed data[28,30] (2,3) are as follows:

$$h(t, X) = h_0(t) \exp(\beta_1 x_1 + \beta_2 x_2 + \dots + \beta_m x_m) \quad (1)$$

$$\text{pooled } \ln HR = \sum \left(\frac{\ln HR}{v} \right) / \sum \left(\frac{1}{v} \right) \quad (2)$$

$$V = [\ln(\text{upper } 95\%CI) - \ln(\text{lower } 95\%CI)]^2 / (2 * 1.96)^2 \quad (3)$$

Heterogeneity assessment and sensitivity analysis

The heterogeneity across the studies was assessed and reported as a percentage using the I^2 index value[37] and as a P value using the Cochrane Q test of chi-square[38]. $I^2 < 25\%$ suggests low heterogeneity, 25% to 50% suggests moderate heterogeneity, and $\geq 50\%$ suggests high heterogeneity. $P < 0.1$ for the Q test suggests high heterogeneity, and ≥ 0.1 suggests low heterogeneity. Only the models without high heterogeneity in both tests were adopted.

Regardless of the degree of heterogeneity, checking calculations were performed by omitting the included studies one by one after a meta-analysis of at least three studies to analyze the sensitivity of the resulting model. A study was considered to introduce instability when the new pooled HR value obtained from the meta-analysis after its omission was distant from the previously obtained HR or beyond the 95%CI range.

Publication bias assessment

The following methods of publication bias assessment were only performed if at least three studies were included in each meta-analysis. We assessed publication bias using Egger's test[39]. $P < 0.05$ indicated a high publication bias. Funnel plots [40] were also drawn. An asymmetric plot with the pooled HR value as the axis was considered to indicate high publication bias.

Evidence quality grade assessment

After finishing the meta-analyses of all outcomes of interest, we used the Grading of Recommendations Assessment, Development and Evaluation system (GRADE)[41] to evaluate the qualities of evidence and make recommendations. Each result was graded as high, moderate, low, or very low. The results derived from the meta-analyses of RCTs were initially set as high, and the rating was lowered by a corresponding number of levels if the result appeared suspect in terms of overall bias, publication bias, inconsistency, imprecision, or indirectness. Those derived from the meta-analyses of cohort studies were initially set as very low, and the rating was raised by a corresponding number of levels if the result appeared suspect in terms of large magnitude of effect, dose-response gradient, or plausible confounding. The assessment was performed by a pair of independent authors (Li MX and Tu HX). When an assessment result was discordant and consensus could not be reached, the one with the lower grade was adopted. The higher assessment grade was adopted when two meta-analyses according to different study types were performed on the same outcome.

RESULTS

Selected studies and extracted data

We initially identified 1,234 articles by searching the 4 academic databases; 507 articles remained after removing duplicates. Thirteen articles were retained after title abstract filtering. After reviewing the full text, 5 papers[24,25,27,42,43] were retained, and 3 additional papers[26,44,45] obtained from the references in these articles were included in the final study. One study[25] was the continuation of another study[24] from the same RCT (PADI trial) over a longer follow-up period and reported an updated HR of ACD for 10 follow-up years. The PRISMA flowchart of study selection is shown in Figure 1.

All the control treatments adopted in the included studies could be divided into 2 categories: PTA with primary BMSI [26,27,42-44] and PTA with or without provisional BMSI[24,25,45]. In the recruitment or selection process, 2 studies[26,45] included a small number of non-CLIs, while the others included only CLIs. On grouping for outcome analysis, the 8 studies all followed the "intention-to-treat" principle rather than the "as-treated" principle. A total of 1493 patients who underwent DESI and 1146 who underwent control treatments were included. Their mean age exceeded 69 y, and the majority were male. The specific main characteristics of these studies and the baseline patient data are shown in Table 1. Some of the definitions of outcomes (other than ACD) in these studies were inevitably different. The specific definitions are shown in Table 2.

Risks of bias

After assessment with the RoB 2.0 tool, the overall risk of bias was "high" in three[24,25,43] of the 5 included RCTs, and "some concerns" regarding this risk were found for the other two studies[26,45]. The 3 included cohort studies[27,42,44] were all at a "moderate" overall risk level after assessment by the ROBINS-I tool. The detailed final assessment results are shown in Table 3.

ACD-free survival

One RCT[25] that directly reported the adjusted HR and another[43] that provided a K-M survival curve were included in a meta-analysis. The low heterogeneity suggested that there was no significant difference between DESI and control treatments in the risk of postoperative ACD [HR = 0.91 (95%CI: 0.38-2.18)] (Figure 2). A meta-analysis including 2 cohort studies[27,42] that directly reported the adjusted HRs and one[44] that provided a K-M survival curve yielded a similar result [HR = 1.15 (95%CI: 0.68-1.95)] with lower heterogeneity (Figure 3A). The sensitivity analysis suggested that this model had high stability (Figure 3B). The model had a P value of 0.137 by Egger's test, and its funnel plot was roughly symmetrical, indicating low publication bias (Figure 3C).

MA-free survival

The results of a meta-analysis including 2 cohort studies[27,42] that directly reported the adjusted HRs and one[44] that provided a K-M survival curve suggested low heterogeneity and that there was no significant difference between the two arms in the risk of postoperative MA [HR = 1.20 (95%CI: 0.84-1.71)] (Figure 4A). The stability of the model was unsatisfactory (Figure 4B), but the publication bias was low ($P = 0.350$ for Egger's test) (Figure 4C). Only one result[24] was available among the included RCTs, which was a directly reported adjusted HR [1.64 (95%CI: 0.74-3.70)], similar to the above findings.

TLR-free survival

The 3 included cohort studies[27,42,44] all directly reported the adjusted HRs of TLR-free survival. A primary meta-analysis yielded a result favoring DESI [HR = 1.93 (95%CI: 1.16-3.22)] (Figure 5A). However, the model was highly heterogeneous ($I^2 = 56.3\%$) and was not adopted. We conducted meta-analyses of pairwise combinations of the 3 studies

Table 1 Main characteristics and baselines of the included studies

Ref.	Design	Coutry/Registry	Enrollment period	Drug on DES	Control	Optimal stenting ^a	Num of patients	Mean age (yr)	Male (%)	Num of limbs	Num of arteries	Num of lesions	Median RC	Mean LRD (mm)	Mean LL (mm)	CTO (%)	Follow up period (yr) ^b
Siablis <i>et al</i> [42], 2009	PCS	Greece	NA	S	PTA+BMSI	N vs N	62 vs 41	69 vs 72	71 vs 90	75 vs 47	NA	153 vs 77	5 vs 5	NA	55 vs 45	25 vs 35	1
Karnabatidis <i>et al</i> [44], 2011	ACS	Greece	2006-2009	S	PTA+BMSI	Y vs N	47 vs 34	71 vs 71	74 vs 82	51 vs 36	75 vs 57	102 vs 72	4 vs 5	NA	76 vs 77	17 vs 35	1
Scheinert <i>et al</i> [45], 2012	RCT	ACHILLES	2008-2010	S	PTA±BMSI	Y vs N	99 vs 101	72 vs 74	68 vs 75	NA	NA	113 vs 115	4 vs 4	2.6 vs 2.6	27 vs 27	81 vs 75	1
Rastan <i>et al</i> [26], 2012	RCT	Germany	2006-2008	S	PTA+BMSI	Y vs Y	82 vs 79	73 vs 72	68 vs 65	NA	NA	82 vs 79	4 vs 3	3.0 vs 3.0	30 vs 31	23 vs 22	3
Bosiers <i>et al</i> [43], 2012	RCT	DESTINY	2008-2009	E	PTA+BMSI	Y vs Y	74 vs 66	75 vs 76	61 vs 58	78 vs 76	78 vs 76	78 vs 76	4.5 vs 5	3.0 vs 2.9	16 vs 19	15 vs 17	1
Spren <i>et al</i> [24], 2017	RCT	PADI	2007-2013	P	PTA±BMSI	Y vs N	73 vs 64	74 vs 73	67 vs 73	74 vs 66	NA	121 vs 91	5 vs 5	2.9 vs 2.9	21 vs 23	NA	3
Konijn <i>et al</i> [25], 2020	RCT	PADI	2007-2013	P	PTA±BMSI	Y vs N	73 vs 64	74 vs 73	67 vs 73	74 vs 66	NA	121 vs 91	5 vs 5	2.9 vs 2.9	21 vs 23	NA	10
Zuzek <i>et al</i> [27], 2022	RCS	USA	2016-2017	NA	PTA+BMSI	NA	1056 vs 761	72 vs 72	59 vs 64	NA	NA	NA	5 vs 5	NA	NA	NA	0.5

^aDES implantation was performed regardless of whether predilatation was performed or the outcome of predilatation.

^bApproximate mean. The data variables of double arms are presented in the form of "DES vs control".

PCS: Prospective cohort study; ACS: Ambispective cohort study; RCT: Randomized controlled trial; RCS: Retrospective cohort study; NA: Not available; DES: Drug-eluting stent; S: Sirolimus; E: Everolimus; P: Paclitaxel; PTA: Percutaneous transluminal angioplasty; BMSI: Bare metal stent implantation; N: No; Y: Yes; Num: Number; RC: Rutherford classification; LRD: Lesion reference diameter; LL: Lesion length; CTO: Chronic total occlusion.

and obtained a model with low heterogeneity[42,44], which suggested a similar result to that described above [HR = 2.65 (95%CI: 1.56-4.50)] (Figure 5B). Among the RCTs included, only[43] had an available result, *i.e.*, a K-M survival curve. From this curve, we extracted and converted a result [HR = 2.07 (95%CI: 0.78-5.52)], and the findings indicated no significant difference between the two arms.

AE-free survival

The results of a meta-analysis including 2 RCTs[24,26] that directly reported the adjusted HRs and one[45] that provided a K-M survival curve suggested low heterogeneity and that DESI better prevented AEs postoperatively [HR = 1.57 (95%CI: 1.23-2.01)] (Figure 6A). The stability of the model was satisfactory (Figure 6B), and the publication bias was low ($P = 0.917$ for Egger's test) (Figure 6C). Only one result[24] of the included cohort studies was available, which was an adjusted HR [2.19 (95%CI: 1.16-4.13)] that was directly reported and similar to above.

Table 2 Definitions of the outcomes during follow-up extracted from the included studies

Ref.	ACD	MA	TLR	AE	PP
Siablis <i>et al</i> [42], 2009	All-cause death	Amputation above the ankle	Repeated revascularization on the target lesion prompted by deterioration of limb ischemia	NA	Absence of repeated intervention and occlusion detected by angiography in the target lesion
Karnabatidis <i>et al</i> [44], 2011	Same as the top ^a	NI ^a	Same as the top	ACD, or MA, or TLR	Same as the top
Scheinert <i>et al</i> [45], 2012	NA	NA	NA	ACD, or MA, or TLR, or RC ≥ 4 ^a	NA
Rastan <i>et al</i> [26], 2012	NA	NA	NA	ACD, or MA, or target vessel revascularization, or myocardial infarction	NA
Bosiers <i>et al</i> [43], 2012	Same as the top ^a	NA	NI ^a	NA	Absence of ≥ 50% binary ISR detected by angiography ^a
Spren <i>et al</i> [24], 2017	NA	Amputation above the ankle	NA	MA, or target limb revascularization	NA
Konijn <i>et al</i> [25], 2020	Same as the top	NA	NA	NA	NA
Zuzek <i>et al</i> [27], 2022	Same as the top	Amputation above the transmetatarsal	Same as the top	NA	NA

^aData extracted from survival curve presented in the article and thus transformed. ACD: All-cause death; NA: Not available; MA: Major amputation; NI: No information; TLR: Target lesion revascularization; AE: Adverse event; RC: Rutherford classification; PP: Primary patency; ISR: In-stent restenosis.

Table 3 Risk bias assessment results of included studies

The domains in RoB 2.0a for RCTs	Scheinert <i>et al</i> [45], 2012	Rastan <i>et al</i> [26], 2012	Bosiers <i>et al</i> [43], 2012	Spren <i>et al</i> [24], 2017	Konijn <i>et al</i> [25], 2020
1 Randomisation process	Low	Low	Low	Low	Low
2 Deviations from the intended interventions	Low	Low	Low	Low	Low
3 Missing outcome data	Some concerns	Some concerns	High	High	High
4 Measurement of the outcome	Low	Low	Low	Low	Low
5 Selection of the reported result	Low	Low	Low	Low	Low
6 Overall	Some concerns	Some concerns	High	High	High
The domains in ROBINS-Ib for cohort studies	Siablis <i>et al</i> [42], 2009	Karnabatidis <i>et al</i> [44], 2011	Zuzek <i>et al</i> [27], 2022		
1 Confounding	Low	Moderate	Moderate		
2 Selection of participants	Low	Low	Low		
3 Classification of interventions	Low	Low	Low		
4 Deviations from intended interventions	Low	Low	Moderate		
5 Missing outcome data	Moderate	Moderate	Moderate		
6 Measurement of outcomes	Moderate	Moderate	Moderate		
7 Selection of the reported result	Low	Low	Low		
8 Overall	Moderate	Moderate	Moderate		

^aThe risk is graded into 3 levels: low: some concerns: and high.

^bThe risk is graded into 5 levels: low: moderate: serious: critical: and no information. RoB: Risk of bias tool; RCT: Randomized controlled trial; ROBINS-I: Risk of bias in non-randomised studies-of interventions.

PP survival

Two cohort studies[42,44] were included in a meta-analysis, both of which directly reported the adjusted HRs. The results of the analysis suggested that the DESI performed better in maintaining PP postoperatively [HR = 5.67 (95%CI: 3.56-9.03)] (Figure 7). Among the RCTs included, only one[43] had an available result, that is, a K-M survival curve. We extracted and converted a result from this curve [HR = 1.68 (95%CI: 0.88-3.94)], and no significant difference was indicated between the two arms.

Evidence quality grade

Only ACD-free survival among the 5 outcomes of interest necessitated 2 meta-analyses due to different study types. We adopted a higher evidence quality grade, *i.e.*, moderate, after the assessment. The detailed assessment results are shown in Table 4.

DISCUSSION

This meta-analysis including 5 RCTs and 3 cohort studies systematically reviewed and analyzed multiple follow-up outcomes of DESI performed in infrapopliteal arteries. The results revealed that DESI showed no superiority in comparison with the control treatments in terms of ACD-free and MA-free survival; however, DESI demonstrated statistically significant advantages in terms of TLR-free, AE-free, and PP survival. Studies[46-49] have extensively demonstrated the superiority of stents eluted with antiproliferative drugs, such as paclitaxel, sirolimus, and everolimus, in coronary arteries with similar luminal diameters to infrapopliteal arteries. It is unsurprising that such promising stents have gradually been used in lower leg lesions. As traditional revascularization modalities[6-10], BMSI and PTA are currently the most commonly used control treatments in related studies. The PADI trial[25] resulted in survival curves up to the 10-year follow-up of DESI for IPADs with PTA ± BMSI as a control. Zuzek *et al*[27] reported survival curves (DESI *vs* BMSI) for multiple outcomes in a cohort study including 1817 participants with IPAD. In recent years, meta-analyses of relevant controlled studies have also been published. In these analyses, few[21,50] reported different pooled ESs after grouping by follow-up period; most[2,51-53] only reported those at the one-year follow-up, and some[20,54] even directly pooled the ESs among different follow-up periods. The outcomes of interest for these analyses were, without exception, the cumulative postoperative incidences of some events at a given follow-up moment, which were considered static indicators. However, as described in the "Introduction" section, our outcomes of interest are all time-to-event data[22]. For example, the ACD of a PADI patient is almost impossible to determine on the first postoperative day of DESI but will occur over time. Unlike the RR or data [which can only evaluate the cumulative risk at a certain period, the HR derived from a proportional hazards regression model (such as the Cox model) including the time variable is more appropriate for assessing the risks of relevant outcomes[28,36]. However, no relevant meta-analyses using HR as the outcome measure were found. In addition, when using the HR for meta-analysis, ESs from studies with different follow-up periods can be included in the same pooled analysis without grouping, and some rare studies with long follow-up periods can also be included (*e.g.*, the PADI trial[25] seems to be the only control study that has reported the relevant outcomes at the 10-year follow-up). This approach is beneficial to expand the sample size and increase the statistical power.

In addition to the 5 RCTs, we also included 3 cohort studies. The inherent flaws in study design that render cohort studies more at risk of bias than RCTs may call them into question[41]. However, the included 3 cohort studies provided more data that could be used for meta-analysis than the above 5 RCTs. In addition, after assessment, no critical risk of bias was found among these cohort studies, and the evidence qualities of the resulting meta-analyses were mostly satisfactory (2 high, 1 medium, and only 1 Low).

We found that the most prominent risks of all the included studies were all from the domain of missing outcome data. Due to very limited information, we could not determine the specific reasons for missing data in each study. However, we believe that censoring due to ACD, which is impossible to completely avoid, is a common and important reason for this type of bias risk in these studies. In addition to the general doubts in the above domain, the risks in other domains of the included studies were not serious after assessment, which is why we did not exclude any of the 8 studies based on the overall assessment results.

The results of statistical analyses are discussed below. First, the two meta-analyses on ACD-free survival both resulted in the same conclusion: DESI for IPADs has no advantages in avoiding postoperative ACD *vs* the control treatments, increasing persuasiveness. Second, a meta-analysis on MA-free survival resulted in a similar conclusion to that described above: DESI has no advantages in avoiding postoperative MA. The model derived from this analysis was considered less stable after the sensitivity analysis. This instability was caused by a study favoring DES and had a very narrow 95%CI range of HR, accounting for over 80% of the weight individually in the pooled analysis[27]. According to the available data, this study was the only retrospective study among the included studies, and its sample size seemed to be significantly larger than that of other studies. However, this is an insufficient explanation for the differing result. It is also slightly regrettable that because the data were derived from cohort studies rather than RCTs and there was only 1 add-on in the assessment, this conclusion was the only one of the 5 recommended conclusions to be assessed as low-quality. Third, after excluding one study, the meta-analysis on TLR-free survival with lower heterogeneity indicated that DESI has significant advantages in avoiding postoperative TLR [HR = 2.65 (95%CI: 1.56-4.50)]. The study that brought major heterogeneity was again the one mentioned above[27], and its result was also different (*i.e.*, no tendency). Fourth, after the meta-analysis on AE-free survival, we concluded that DESI has significant advantages in avoiding postoperative AE [HR = 1.57 (95%CI: 1.23-2.01)]. Similarly, one study[45] was not statistically consistent with the other two[24,26], and the final

Table 4 Evidence quality grade assessment of pooled outcomes of interest


Outcome	Source of data	Num of participants	Pooled HR (DES vs control)	Certainty of the evidence (GRADE) ^a	Alteration to initial rating
ACD-free survival	2 RCTs	277	0.91 (95%CI: 0.38-2.18)	Moderate	-1 ^b
ACD-free survival	3 cohort studies	2001	1.15 (95%CI: 0.68-1.95)	Low	+1 ^c
MA-free survival	3 cohort studies	2001	1.20 (95%CI: 0.84-1.71)	Low	+1 ^c
TLR-free survival	2 cohort studies	184	2.65 (95%CI: 1.56-4.50)	High	+3 ^d
AE-free survival	3 RCTs	498	1.57 (95%CI: 1.23-2.01)	Moderate	-1 ^b
PP survival	2 cohort studies	184	5.67 (95%CI: 3.56-9.03)	High	+3 ^d

^aHigh certainty: We are very confident that the true effect lies close to that of the estimate of the effect; moderate certainty: We are moderately confident in the effect estimate; low certainty: Our confidence in the effect estimate is limited; very low certainty: We have very little confidence in the effect estimate.

^bThe reason for lowering rating was a suspect of overall bias.

^cThe reason for raising rating was a consideration of plausible confounding.

^dThe reason for raising rating was a consideration of large magnitude of effect, dose-response gradient, and plausible confounding. ACD: All-cause death; MA: major amputation; TLR: Target lesion revascularization; AE: Adverse event; PP: Primary patency; RCT: Randomized controlled trial; Num: number; HR: Hazard ratio; CI: Confidence interval; GRADE: Grading of recommendations assessment: Development and evaluation.

**Figure 1 PRISMA flowchart.** PRISMA: Preferred Reporting Items for Systematic Reviews and Meta-Analyses.

result was “no tendency”. The biggest evident discrepancy in that study is that its follow-up period was only 1 year, while the others involved a 3-year follow-up. Thus, AE-free survival and other IPAD outcomes potentially do not develop linearly, and the advantages of DESI gradually emerge with time. Fifth, the meta-analysis on PP survival revealed that DESI has significant advantages in maintaining postoperative PP survival [HR = 5.67 (95%CI: 3.56-9.03)]. The high HR value reflects the great advantages of DES in this respect.

This study has some limitations. First, the sample size is small, which reduces the persuasiveness of the results. To date, the number of controlled studies on DESI for IPADs is still limited in comparison with those on DESI for femoropopliteal artery diseases[55-66]. DESs specific for femoropopliteal arteries, such as Zilver PTX (Cook Medical, United States) and Eluvia (Boston Scientific, United States). Have been used in many countries. However, a dedicated infrapopliteal artery stent is absent, at least in China, and a coronary stent is used. Second, the definitions of some outcomes of interest slightly differed among the included studies. For example, one study[26] regarded myocardial infarction as an AE, while others with relevant data did not; some studies[43,44] did not even explain the definitions of certain outcomes, increasing the risks of error in the results. Third, to bring as many studies as possible into the analyses, we extracted data from the survival curves given in some of the studies[43-45]. Although sufficient evidence[28,30,31]




Figure 2 Forest plot of all-cause death-free survival derived from pooling 2 randomized controlled trials (random effects model). HR: Hazard ratio; CI: Confidence interval; REML: Restricted maximum likelihood; DES: Drug-eluting stent.




Figure 3 All-cause death-free survival. A: Forest plot derived from pooling 3 cohort studies (random effects model); B: Sensitivity analysis of the model assuming that each study was omitted separately [$\ln(\text{HR})$]; C: Funnel plot with pseudo 95% confidence limits. HR: Hazard ratio; CI: Confidence interval; REML: Restricted maximum likelihood; DES: Drug-eluting stent; ACD: All-cause death.

supports the rationality of this approach, it is, after all, a recalculation and not a direct HR, unlike that in the other studies, which inevitably increases the error.

CONCLUSION

Compared with traditional treatment modalities (PTA with or without BMSI), DESI for IPADs is significantly superior in avoiding TLR and AEs and maintaining PP survival, while showing no superiority or inferiority in terms of ACD-free and MA-free survival. In conclusion, DES is a good option for IPADs to maintain efficacy for long periods.




Figure 4 Major amputation -free survival. A: Forest plot derived from pooling 3 cohort studies (random effects model); B: Sensitivity analysis of the model assuming that each study was omitted separately [$\ln(HR)$]; C: Funnel plot with pseudo 95% confidence limits. HR: Hazard ratio; CI: Confidence interval; REML: Restricted maximum likelihood; DES: Drug-eluting stent.




Figure 5 Target lesion revascularization -free survival. A: Forest plot with high heterogeneity derived from pooling 3 cohort studies (random effects model); B: Forest plot with low heterogeneity derived from pooling 2 cohort studies (random effects model). HR: Hazard ratio; CI: Confidence interval; REML: Restricted maximum likelihood; DES: Drug-eluting stent; TLR: Target lesion revascularization.




Figure 6 Adverse event -free survival. A: Forest plot derived from pooling 3 cohort studies (random effects model); B: Sensitivity analysis of the model assuming that each study was omitted separately [$\ln(HR)$]; C: Funnel plot with pseudo 95% confidence limits. HR: Hazard ratio; CI: Confidence interval; REML: Restricted maximum likelihood; DES: Drug-eluting stent.




Figure 7 Forest plot of primary patency survival derived from pooling 2 cohort studies (random effects model). HR: Hazard ratio; CI: Confidence interval; REML: Restricted maximum likelihood; DES: Drug-eluting stent.

ARTICLE HIGHLIGHTS

Research background

Whether accompanied with femoropopliteal inflow disease or not, infrapopliteal artery disease (IPAD) is the primary cause of critical limb ischemia. In the past few decades, minimally invasive percutaneous transluminal angioplasty (PTA) with or without bare metal stent implantation (BMSI) has been widely used.

Research motivation

However, although this treatment has satisfactory technical success rate, it still has a significantly high risk of clinical failure caused by lesion restenosis even in the short term.

Research objectives

In order to more accurately evaluate the efficacy of drug-eluting stents (DES) implantation for IPADs, we performed this systematic review and meta-analysis.

Research methods

After extensive retrieval of major databases, the hazard ratio (HR) is used as the outcome measure for extraction or conversion, and the meta analyses for multiple outcomes of interest were performed.

Research results

Five randomized controlled trials and three cohort studies involving 2639 participants totally were included. Compared with the control arm (PTA and BMSI), the HR values of the DES implantation on all-cause death-free survival and major amputation-free survival were not statistically significant ($P > 0.05$), but the HR values on target lesion revascularization-free survival, adverse event-free survival, and primary patency-survival were 2.65 (95%CI: 1.56-4.50), 1.57 (95%CI: 1.23-2.01), and 5.67 (95%CI: 3.56-9.03), respectively.

Research conclusions

In our conclusion, DES is a good option for IPADs to maintain efficacy for a long time.

Research perspectives

DES is a highly anticipated therapeutic device. We believe that there will be more and more randomized controlled trials about its application for IPADs in the future.

FOOTNOTES

Author contributions: Li MX designed the study; Li MX and Tu HX performed the article search, data extraction, and evidence quality assessment; Li MX and Yin MC performed the risk of bias assessment of the studies; Li MX performed the data analyses and manuscript writing; Tu HX reviewed the article independently and made minor revisions after consultation with Li MX; All the authors read and gave final approval of the version to be submitted; Tu HX is the guarantor of the review.

Conflict-of-interest statement: The authors declare that they have no conflicts of interest to report.

PRISMA 2009 Checklist statement: The authors have read the PRISMA 2009 Checklist, and the manuscript was prepared and revised according to the PRISMA 2009 Checklist.

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <https://creativecommons.org/licenses/by-nc/4.0/>

Country/Territory of origin: China

ORCID number: Ming-Xuan Li 0000-0002-0473-5667; Hai-Xia Tu 0000-0002-0252-5762; Meng-Chen Yin 0009-0003-7970-5862.

S-Editor: Liu JH

L-Editor: A

P-Editor: Cai YX

REFERENCES

- 1 Rueda CA, Nehler MR, Perry DJ, McLafferty RB, Casserly IP, Hiatt WR, Peyton BD. Patterns of artery disease in 450 patients undergoing revascularization for critical limb ischemia: implications for clinical trial design. *J Vasc Surg* 2008; **47**: 995-9; discussion 999 [PMID: 18372151 DOI: 10.1016/j.jvs.2007.11.055]
- 2 Katsanos K, Spiliopoulos S, Diamantopoulos A, Karnabatidis D, Sabharwal T, Siablis D. Systematic review of infrapopliteal drug-eluting stents: a meta-analysis of randomized controlled trials. *Cardiovasc Intervent Radiol* 2013; **36**: 645-658 [PMID: 23435741 DOI: 10.1007/s00270-013-0578-2]
- 3 Faries PL, Arora S, Pomposelli FB Jr, Pulling MC, Smakowski P, Rohan DI, Gibbons GW, Akbari CM, Campbell DR, LoGerfo FW. The use of arm vein in lower-extremity revascularization: results of 520 procedures performed in eight years. *J Vasc Surg* 2000; **31**: 50-59 [PMID: 10642708 DOI: 10.1016/s0741-5214(00)70067-x]
- 4 Ballard JL, Mills JL Sr. Surgical management of critical limb ischemia. *Tech Vasc Interv Radiol* 2005; **8**: 169-174 [PMID: 16849097 DOI: 10.1053/j.tvir.2006.04.004]
- 5 Norgren L, Hiatt WR, Dormandy JA, Nehler MR, Harris KA, Fowkes FG; TASC II Working Group. Inter-Society Consensus for the Management of Peripheral Arterial Disease (TASC II). *J Vasc Surg* 2007; **45** Suppl S: S5-67 [PMID: 17223489 DOI: 10.1016/j.jvs.2006.12.037]
- 6 Horvath W, Oertl M, Haidinger D. Percutaneous transluminal angioplasty of crural arteries. *Radiology* 1990; **177**: 565-569 [PMID: 2145609 DOI: 10.1148/radiology.177.2.2145609]
- 7 Brown KT, Moore ED, Getrajdman GI, Saddekni S. Infrapopliteal angioplasty: long-term follow-up. *J Vasc Interv Radiol* 1993; **4**: 139-144 [PMID: 8425091 DOI: 10.1016/s1051-0443(93)71836-9]

- 8 **Söder HK**, Manninen HI, Jaakkola P, Matsi PJ, Räsänen HT, Kaukanen E, Lopenon P, Soimakallio S. Prospective trial of infrapopliteal artery balloon angioplasty for critical limb ischemia: angiographic and clinical results. *J Vasc Interv Radiol* 2000; **11**: 1021-1031 [PMID: [10997465](#) DOI: [10.1016/s1051-0443\(07\)61332-3](#)]
- 9 **Rastogi S**, Stavropoulos SW. Infrapopliteal angioplasty. *Tech Vasc Interv Radiol* 2004; **7**: 33-39 [PMID: [15071779](#) DOI: [10.1053/j.tvir.2004.01.001](#)]
- 10 **Tsetis D**, Belli AM. The role of infrapopliteal angioplasty. *Br J Radiol* 2004; **77**: 1007-1015 [PMID: [15569642](#) DOI: [10.1259/bjr/97382129](#)]
- 11 **Albers M**, Romiti M, Brochado-Neto FC, De Luccia N, Pereira CA. Meta-analysis of popliteal-to-distal vein bypass grafts for critical ischemia. *J Vasc Surg* 2006; **43**: 498-503 [PMID: [16520163](#) DOI: [10.1016/j.jvs.2005.11.025](#)]
- 12 **Haider SN**, Kavanagh EG, Forlee M, Colgan MP, Madhavan P, Moore DJ, Shanik GD. Two-year outcome with preferential use of infrainguinal angioplasty for critical ischemia. *J Vasc Surg* 2006; **43**: 504-512 [PMID: [16520164](#) DOI: [10.1016/j.jvs.2005.11.016](#)]
- 13 **Romiti M**, Albers M, Brochado-Neto FC, Durazzo AE, Pereira CA, De Luccia N. Meta-analysis of infrapopliteal angioplasty for chronic critical limb ischemia. *J Vasc Surg* 2008; **47**: 975-981 [PMID: [18372148](#) DOI: [10.1016/j.jvs.2008.01.005](#)]
- 14 **Katsanos K**, Spiliopoulos S, Karunanithy N, Krokidis M, Sabharwal T, Taylor P. Bayesian network meta-analysis of nitinol stents, covered stents, drug-eluting stents, and drug-coated balloons in the femoropopliteal artery. *J Vasc Surg* 2014; **59**: 1123-1133.e8 [PMID: [24661896](#) DOI: [10.1016/j.jvs.2014.01.041](#)]
- 15 **Meng FC**, Chen PL, Lee CY, Shih CC, Chen IM. Real-World Comparison of Drug-Eluting and Bare-Metal Stents in Superficial Femoral Artery Occlusive Disease with Trans-Atlantic Intersociety Consensus B Lesions: A 2-Year, Single-Institute Study. *Acta Cardiol Sin* 2018; **34**: 130-136 [PMID: [29643698](#) DOI: [10.6515/ACS.201803_34\(2\).20171126A](#)]
- 16 **Ding Y**, Zhou M, Wang Y, Cai L, Shi Z. Comparison of Drug-Eluting Stent with Bare-Metal Stent Implantation in Femoropopliteal Artery Disease: A Systematic Review and Meta-Analysis. *Ann Vasc Surg* 2018; **50**: 96-105 [PMID: [29514049](#) DOI: [10.1016/j.avsg.2017.12.003](#)]
- 17 **Siablis D**, Kraniotis P, Karnabatidis D, Kagadis GC, Katsanos K, Tsolakis J. Sirolimus-eluting versus bare stents for bailout after suboptimal infrapopliteal angioplasty for critical limb ischemia: 6-month angiographic results from a nonrandomized prospective single-center study. *J Endovasc Ther* 2005; **12**: 685-695 [PMID: [16363898](#) DOI: [10.1583/05-1620MR.1](#)]
- 18 **Commeau P**, Barragan P, Roquebert PO. Sirolimus for below the knee lesions: mid-term results of SiroBTK study. *Catheter Cardiovasc Interv* 2006; **68**: 793-798 [PMID: [17039538](#) DOI: [10.1002/ccd.20893](#)]
- 19 **Werner M**, Schmidt A, Freyer M, Bausback Y, Bräunlich S, Friedenberger J, Schuster J, Botsios S, Scheinert D, Ulrich M. Sirolimus-eluting stents for the treatment of infrapopliteal arteries in chronic limb ischemia: long-term clinical and angiographic follow-up. *J Endovasc Ther* 2012; **19**: 12-19 [PMID: [22313195](#) DOI: [10.1583/11-3665.1](#)]
- 20 **Fusaro M**, Cassese S, Ndrepepa G, Tepe G, King L, Ott I, Nerad M, Schunkert H, Kastrati A. Drug-eluting stents for revascularization of infrapopliteal arteries: updated meta-analysis of randomized trials. *JACC Cardiovasc Interv* 2013; **6**: 1284-1293 [PMID: [24355118](#) DOI: [10.1016/j.jcin.2013.08.007](#)]
- 21 **Liu X**, Zheng G, Wen S. Drug-eluting stents versus control therapy in the infrapopliteal disease: A meta-analysis of eight randomized controlled trials and two cohort studies. *Int J Surg* 2017; **44**: 166-175 [PMID: [28648791](#) DOI: [10.1016/j.ijsu.2017.06.075](#)]
- 22 **Tudur C**, Williamson PR, Khan S, Best L. The value of the aggregate data approach in meta-analysis with time-to-event outcomes. *Journal of the Royal Statistical Society A* 2001; **164**: 357-370 [DOI: [10.1111/1467-985X.00207](#)]
- 23 **Spreen MI**, Martens JM, Hansen BE, Knippenberg B, Verhey E, van Dijk LC, de Vries JP, Vos JA, de Borst GJ, Vonken EJ, Wever JJ, Statius van Eps RG, Mali WP, van Overhagen H. Percutaneous Transluminal Angioplasty and Drug-Eluting Stents for Infrapopliteal Lesions in Critical Limb Ischemia (PADI) Trial. *Circ Cardiovasc Interv* 2016; **9**: e002376 [PMID: [26861113](#) DOI: [10.1161/CIRCINTERVENTIONS.114.002376](#)]
- 24 **Spreen MI**, Martens JM, Knippenberg B, van Dijk LC, de Vries JPM, Vos JA, de Borst GJ, Vonken EPA, Bijlstra OD, Wever JJ, Statius van Eps RG, Mali WPTM, van Overhagen H. Long-Term Follow-up of the PADI Trial: Percutaneous Transluminal Angioplasty Versus Drug-Eluting Stents for Infrapopliteal Lesions in Critical Limb Ischemia. *J Am Heart Assoc* 2017; **6** [PMID: [28411244](#) DOI: [10.1161/JAHA.116.004877](#)]
- 25 **Konijn LCD**, Wakkie T, Spreen MI, de Jong PA, van Dijk LC, Wever JJ, Veger HTC, Statius van Eps RG, Mali WPTM, van Overhagen H. 10-Year Paclitaxel Dose-Related Outcomes of Drug-Eluting Stents Treated Below the Knee in Patients with Chronic Limb-Threatening Ischemia (The PADI Trial). *Cardiovasc Intervent Radiol* 2020; **43**: 1881-1888 [PMID: [32725411](#) DOI: [10.1007/s00270-020-02602-6](#)]
- 26 **Rastan A**, Brechtel K, Krankenberg H, Zahorsky R, Tepe G, Noory E, Schwarzwälder U, Macharzina R, Schwarz T, Bürgelin K, Sixt S, Tübler T, Neumann FJ, Zeller T. Sirolimus-eluting stents for treatment of infrapopliteal arteries reduce clinical event rate compared to bare-metal stents: long-term results from a randomized trial. *J Am Coll Cardiol* 2012; **60**: 587-591 [PMID: [22878166](#) DOI: [10.1016/j.jacc.2012.04.035](#)]
- 27 **Zuzek Z**, Arora S, Helmy I, Jani C, Jaswaney R, Patel K, Patel HP, Patel M, Osman MN, Li J, Shishehbor MH. Underutilization of Drug-Eluting Stents in Infrapopliteal Intervention for Chronic Limb-Threatening Ischemia. *J Endovasc Ther* 2023; **30**: 45-56 [PMID: [35075941](#) DOI: [10.1177/15266028211068763](#)]
- 28 **Tierney JF**, Stewart LA, Ghersi D, Burdett S, Sydes MR. Practical methods for incorporating summary time-to-event data into meta-analysis. *Trials* 2007; **8**: 16 [PMID: [17555582](#) DOI: [10.1186/1745-6215-8-16](#)]
- 29 **Liberati A**, Altman DG, Tetzlaff J, Mulrow C, Gotzsche PC, Ioannidis JP, Clarke M, Devereaux PJ, Kleijnen J, Moher D. The PRISMA statement for reporting systematic reviews and meta-analyses of studies that evaluate healthcare interventions: explanation and elaboration. *BMJ* 2009; **339**: b2700 [PMID: [19622552](#) DOI: [10.1136/bmj.b2700](#)]
- 30 **Parmar MK**, Torri V, Stewart L. Extracting summary statistics to perform meta-analyses of the published literature for survival endpoints. *Stat Med* 1998; **17**: 2815-2834 [PMID: [9921604](#) DOI: [10.1002/\(sici\)1097-0258\(19981230\)17:24<2815::aid-sim110>3.0.co;2-8](#)]
- 31 **Williamson PR**, Smith CT, Hutton JL, Marson AG. Aggregate data meta-analysis with time-to-event outcomes. *Stat Med* 2002; **21**: 3337-3351 [PMID: [12407676](#) DOI: [10.1002/sim.1303](#)]
- 32 **Sterne JAC**, Savović J, Page MJ, Elbers RG, Blencowe NS, Boutron I, Cates CJ, Cheng HY, Corbett MS, Eldridge SM, Emberson JR, Hernán MA, Hopewell S, Hróbjartsson A, Junqueira DR, Jüni P, Kirkham JJ, Lasserson T, Li T, McAleenan A, Reeves BC, Shepperd S, Shrier I, Stewart LA, Tilling K, White IR, Whiting PF, Higgins JPT. RoB 2: a revised tool for assessing risk of bias in randomised trials. *BMJ* 2019; **366**: 14898 [PMID: [31462531](#) DOI: [10.1136/bmj.14898](#)]
- 33 **Sterne JA**, Hernán MA, Reeves BC, Savović J, Berkman ND, Viswanathan M, Henry D, Altman DG, Ansari MT, Boutron I, Carpenter JR, Chan AW, Churchill R, Deeks JJ, Hróbjartsson A, Kirkham J, Jüni P, Loke YK, Pigott TD, Ramsay CR, Regidor D, Rothstein HR, Sandhu L, Santaguida PL, Schünemann HJ, Shea B, Shrier I, Tugwell P, Turner L, Valentine JC, Waddington H, Waters E, Wells GA, Whiting PF,

- Higgins JP. ROBINS-I: a tool for assessing risk of bias in non-randomised studies of interventions. *BMJ* 2016; **355**: i4919 [PMID: [27733354](#) DOI: [10.1136/bmj.i4919](#)]
- 34 **DerSimonian R**, Kacker R. Random-effects model for meta-analysis of clinical trials: an update. *Contemp Clin Trials* 2007; **28**: 105-114 [PMID: [16807131](#) DOI: [10.1016/j.cct.2006.04.004](#)]
 - 35 **Lindstrom MJ**, Bates DM. Newton-Raphson and EM Algorithms for Linear Mixed-Effects Models for Repeated-Measures Data. *J Am Stat Assoc* 1988; **83**: 1014-1022 [DOI: [10.1080/01621459.1988.10478693](#)]
 - 36 **Cox DR**. Regression models and life tables. *Journal of the Royal Statistical Society* 1972; **34**: 187 [DOI: [10.1007/978-1-4612-4380-9_37](#)]
 - 37 **Thakkinian A**, McElduff P, D'Este C, Duffy D, Attia J. A method for meta-analysis of molecular association studies. *Stat Med* 2005; **24**: 1291-1306 [PMID: [15568190](#) DOI: [10.1002/sim.2010](#)]
 - 38 **Whitehead A**, Whitehead J. A general parametric approach to the meta-analysis of randomized clinical trials. *Stat Med* 1991; **10**: 1665-1677 [PMID: [1792461](#) DOI: [10.1002/sim.4780101105](#)]
 - 39 **Egger M**, Davey Smith G, Schneider M, Minder C. Bias in meta-analysis detected by a simple, graphical test. *BMJ* 1997; **315**: 629-634 [PMID: [9310563](#) DOI: [10.1136/bmj.315.7109.629](#)]
 - 40 **Biljana M**, Jelena M, Branislav J, Milorad R. Bias in meta-analysis and funnel plot asymmetry. *Stud Health Technol Inform* 1999; **68**: 323-328 [PMID: [10724898](#)]
 - 41 **Guyatt GH**, Oxman AD, Vist GE, Kunz R, Falck-Ytter Y, Alonso-Coello P, Schünemann HJ; GRADE Working Group. GRADE: an emerging consensus on rating quality of evidence and strength of recommendations. *BMJ* 2008; **336**: 924-926 [PMID: [18436948](#) DOI: [10.1136/bmj.39489.470347.AD](#)]
 - 42 **Siablis D**, Karnabatidis D, Katsanos K, Diamantopoulos A, Spiliopoulos S, Kagadis GC, Tsolakis J. Infrapopliteal application of sirolimus-eluting versus bare metal stents for critical limb ischemia: analysis of long-term angiographic and clinical outcome. *J Vasc Interv Radiol* 2009; **20**: 1141-1150 [PMID: [19620014](#) DOI: [10.1016/j.jvir.2009.05.031](#)]
 - 43 **Bosiers M**, Scheinert D, Peeters P, Torsello G, Zeller T, Deloose K, Schmidt A, Tessarek J, Vinck E, Schwartz LB. Randomized comparison of everolimus-eluting versus bare-metal stents in patients with critical limb ischemia and infrapopliteal arterial occlusive disease. *J Vasc Surg* 2012; **55**: 390-398 [PMID: [22169682](#) DOI: [10.1016/j.jvs.2011.07.099](#)]
 - 44 **Karnabatidis D**, Spiliopoulos S, Diamantopoulos A, Katsanos K, Kagadis GC, Kakkos S, Siablis D. Primary everolimus-eluting stenting versus balloon angioplasty with bailout bare metal stenting of long infrapopliteal lesions for treatment of critical limb ischemia. *J Endovasc Ther* 2011; **18**: 1-12 [PMID: [21314342](#) DOI: [10.1583/10-3242.1](#)]
 - 45 **Scheinert D**, Katsanos K, Zeller T, Koppensteiner R, Commeau P, Bosiers M, Krankenberg H, Baumgartner I, Siablis D, Lammer J, Van Ransbeeck M, Qureshi AC, Stoll HP; ACHILLES Investigators. A prospective randomized multicenter comparison of balloon angioplasty and infrapopliteal stenting with the sirolimus-eluting stent in patients with ischemic peripheral arterial disease: 1-year results from the ACHILLES trial. *J Am Coll Cardiol* 2012; **60**: 2290-2295 [PMID: [23194941](#) DOI: [10.1016/j.jacc.2012.08.989](#)]
 - 46 **Morice MC**, Serruys PW, Sousa JE, Fajadet J, Ban Hayashi E, Perin M, Colombo A, Schuler G, Barragan P, Guagliumi G, Molnar F, Falotico R; RAVEL Study Group. Randomized Study with the Sirolimus-Coated Bx Velocity Balloon-Expandable Stent in the Treatment of Patients with de Novo Native Coronary Artery Lesions. A randomized comparison of a sirolimus-eluting stent with a standard stent for coronary revascularization. *N Engl J Med* 2002; **346**: 1773-1780 [PMID: [12050336](#) DOI: [10.1056/NEJMoa012843](#)]
 - 47 **Grube E**, Silber S, Hauptmann KE, Mueller R, Buellesfeld L, Gerckens U, Russell ME. TAXUS I: six- and twelve-month results from a randomized, double-blind trial on a slow-release paclitaxel-eluting stent for de novo coronary lesions. *Circulation* 2003; **107**: 38-42 [PMID: [12515740](#) DOI: [10.1161/01.cir.0000047700.58683.a1](#)]
 - 48 **Otsuka F**, Vorpahl M, Nakano M, Foerst J, Newell JB, Sakakura K, Kutys R, Ladich E, Finn AV, Kolodgie FD, Virmani R. Pathology of second-generation everolimus-eluting stents versus first-generation sirolimus- and paclitaxel-eluting stents in humans. *Circulation* 2014; **129**: 211-223 [PMID: [24163064](#) DOI: [10.1161/CIRCULATIONAHA.113.001790](#)]
 - 49 **Neumann FJ**, Sousa-Uva M, Ahlsson A, Alfonso F, Banning AP, Benedetto U, Byrne RA, Collet JP, Falk V, Head SJ, Jüni P, Kastrati A, Koller A, Kristensen SD, Niebauer J, Richter DJ, Seferovic PM, Sibbing D, Stefanini GG, Windecker S, Yadav R, Zembala MO; ESC Scientific Document Group. 2018 ESC/EACTS Guidelines on myocardial revascularization. *Eur Heart J* 2019; **40**: 87-165 [PMID: [30165437](#) DOI: [10.1093/eurheartj/ehy394](#)]
 - 50 **Varcoe RL**, Paravastu SC, Thomas SD, Bennett MH. The use of drug-eluting stents in infrapopliteal arteries: an updated systematic review and meta-analysis of randomized trials. *Int Angiol* 2019; **38**: 121-135 [PMID: [30650949](#) DOI: [10.23736/S0392-9590.19.04049-5](#)]
 - 51 **Antoniou GA**, Chalmers N, Kanesalingham K, Antoniou SA, Schiro A, Serracino-Inglott F, Smyth JV, Murray D. Meta-analysis of outcomes of endovascular treatment of infrapopliteal occlusive disease with drug-eluting stents. *J Endovasc Ther* 2013; **20**: 131-144 [PMID: [23581752](#) DOI: [10.1583/1545-1550-20.2.131](#)]
 - 52 **Katsanos K**, Kitrou P, Spiliopoulos S, Diamantopoulos A, Karnabatidis D. Comparative Effectiveness of Plain Balloon Angioplasty, Bare Metal Stents, Drug-Coated Balloons, and Drug-Eluting Stents for the Treatment of Infrapopliteal Artery Disease: Systematic Review and Bayesian Network Meta-analysis of Randomized Controlled Trials. *J Endovasc Ther* 2016; **23**: 851-863 [PMID: [27708143](#) DOI: [10.1177/1526602816671740](#)]
 - 53 **Zhou Y**, Lin S, Zhang Z, Xiao J, Ai W, Wang J, Li Y, Li Q. A Network Meta-analysis of Randomized Controlled Trials Comparing Treatment Modalities for Infrapopliteal Lesions in Critical Limb Ischemia. *Ann Vasc Surg* 2019; **60**: 424-434 [PMID: [31075473](#) DOI: [10.1016/j.avsg.2019.02.021](#)]
 - 54 **Xiao Y**, Chen Z, Yang Y, Kou L. Network meta-analysis of balloon angioplasty, nondrug metal stent, drug-eluting balloon, and drug-eluting stent for treatment of infrapopliteal artery occlusive disease. *Diagn Interv Radiol* 2016; **22**: 436-443 [PMID: [27559766](#) DOI: [10.5152/dir.2016.15430](#)]
 - 55 **Phair J**, Carnevale M, Lipsitz EC, Shariff S, Scher L, Garg K. Amputation-free Survival in Patients with Critical Limb Ischemia Treated with Paclitaxel-eluting Stents and Paclitaxel-coated Balloons. *Ann Vasc Surg* 2020; **62**: 8-14 [PMID: [31207400](#) DOI: [10.1016/j.avsg.2019.05.013](#)]
 - 56 **Ishihara T**, Iida O, Awata M, Nanto K, Nanto S, Uematsu M. Angioscopic assessment of early phase arterial repair after paclitaxel-coated nitinol drug-eluting stent implantation in the superficial femoral artery. *Circ J* 2013; **77**: 1838-1843 [PMID: [23595033](#) DOI: [10.1253/circj.cj-13-0012](#)]
 - 57 **Vent PA**, Kaladji A, Davaine JM, Guyomarch B, Chaillou P, Costargent A, Quillard T, Gouëffic Y. Bare Metal Versus Paclitaxel-Eluting Stents for Long Femoropopliteal Lesions: Prospective Cohorts Comparison Using a Propensity Score-Matched Analysis. *Ann Vasc Surg* 2017; **43**: 166-175 [PMID: [28300676](#) DOI: [10.1016/j.avsg.2016.10.058](#)]
 - 58 **Soga Y**, Takahara M, Iida O, Mii S, Okazaki J, Nakano M, Yamauchi Y, Ando K. Bypass Surgery vs. Drug-Eluting Stent for Trans-Atlantic

Inter-Society Consensus-II (TASCII) C or D Femoropopliteal Lesions. *Circ J* 2015; **79**: 2688-2695 [PMID: [26466986](#) DOI: [10.1253/circj.CJ-15-0597](#)]

- 59 **Jeon-Slaughter H**, Khalili H, Tsai S, Armstrong EJ, Shammass NW, Jawaaid O, Lu H, Addo T, Gigliotti O, Abu-Fadel M, Banerjee S. Comparative Effectiveness Study of Drug-Eluting and Bare-Metal Peripheral Artery Stents in Endovascular Femoropopliteal Artery Revascularization. *J Invasive Cardiol* 2018; **30**: 373-379 [PMID: [30279293](#)]
- 60 **Haine A**, Schmid MJ, Schindewolf M, Lenz A, Bernhard SM, Drexel H, Baumgartner I, Doppeide JF. Comparison Between Interwoven Nitinol and Drug Eluting Stents for Endovascular Treatment of Femoropopliteal Artery Disease. *Eur J Vasc Endovasc Surg* 2019; **58**: 865-873 [PMID: [31668949](#) DOI: [10.1016/j.ejvs.2019.09.002](#)]
- 61 **Gray WA**, Griffiths RI, Elroy PWM, Amorosi SL, McGovern AM, Jaff MR, Akehurst R, Müller-Hülsbeck S. Cost-effectiveness of a paclitaxel-eluting stent (Eluvia) compared to Zilver PTX for endovascular femoropopliteal intervention. *J Med Econ* 2022; **25**: 880-887 [PMID: [35703041](#) DOI: [10.1080/13696998.2022.2088965](#)]
- 62 **Lee YJ**, Kook H, Ko YG, Yu CW, Joo HJ, Ahn CM, Choi D. Drug Eluting Stent vs. Drug Coated Balloon for Native Femoropopliteal Artery Disease: A Two Centre Experience. *Eur J Vasc Endovasc Surg* 2021; **61**: 287-295 [PMID: [33223281](#) DOI: [10.1016/j.ejvs.2020.10.008](#)]
- 63 **Tomoi Y**, Soga Y, Okazaki J, Iida O, Shiraki T, Hiramori S, Ando K. Drug-coated stent implantation vs. bypass surgery for in-stent occlusion after femoropopliteal stenting. *Heart Vessels* 2021; **36**: 646-653 [PMID: [33392645](#) DOI: [10.1007/s00380-020-01740-8](#)]
- 64 **Duda SH**, Bosiers M, Lammer J, Scheinert D, Zeller T, Oliva V, Tielbeek A, Anderson J, Wiesinger B, Tepe G, Lansky A, Jaff MR, Mudde C, Tieleman H, Beregi JP. Drug-eluting and bare nitinol stents for the treatment of atherosclerotic lesions in the superficial femoral artery: long-term results from the SIROCCO trial. *J Endovasc Ther* 2006; **13**: 701-710 [PMID: [17154704](#) DOI: [10.1583/05-1704.1](#)]
- 65 **Liistro F**, Angioli P, Porto I, Ducci K, Falsini G, Ventoruzzo G, Ricci L, Scatena A, Grotti S, Bolognese L. Drug-Eluting Balloon Versus Drug-Eluting Stent for Complex Femoropopliteal Arterial Lesions: The DRASTICO Study. *J Am Coll Cardiol* 2019; **74**: 205-215 [PMID: [31296293](#) DOI: [10.1016/j.jacc.2019.04.057](#)]
- 66 **Bausback Y**, Wittig T, Schmidt A, Zeller T, Bosiers M, Peeters P, Brucks S, Lottes AE, Scheinert D, Steiner S. Drug-Eluting Stent Versus Drug-Coated Balloon Revascularization in Patients With Femoropopliteal Arterial Disease. *J Am Coll Cardiol* 2019; **73**: 667-679 [PMID: [30765033](#) DOI: [10.1016/j.jacc.2018.11.039](#)]



Published by **Baishideng Publishing Group Inc**
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA

Telephone: +1-925-3991568

E-mail: bpgoffice@wjgnet.com

Help Desk: <https://www.f6publishing.com/helpdesk>

<https://www.wjgnet.com>

