



Kawasaki disease with peritonsillar abscess as the first symptom: A case report

Li-Man Huo, Li-Min Li, Hao-Yang Peng, Li-Jia Wang, Zhang-Ying Feng

Specialty type: Pediatrics

Provenance and peer review:

Unsolicited article; Externally peer reviewed.

Peer-review model: Single blind

Peer-review report's scientific quality classification

Grade A (Excellent): 0

Grade B (Very good): B

Grade C (Good): C

Grade D (Fair): 0

Grade E (Poor): 0

P-Reviewer: Dauey K, Kazakhstan; Kim BS, South Korea

Received: May 21, 2023

Peer-review started: May 21, 2023

First decision: June 13, 2023

Revised: June 25, 2023

Accepted: July 17, 2023

Article in press: July 17, 2023

Published online: August 6, 2023



Li-Man Huo, Department of Pharmacy, The Fourth Hospital of Hebei Medical University, Shijiazhuang 050011, Hebei Province, China

Li-Min Li, Department of Paediatrics, The Fourth Hospital of Hebei Medical University, Shijiazhuang 050011, Hebei Province, China

Hao-Yang Peng, Department of Ultrasound, The Fourth Hospital of Hebei Medical University, Shijiazhuang 050011, Hebei Province, China

Li-Jia Wang, Department of Medical Imaging, The Fourth Hospital of Hebei Medical University, Shijiazhuang 050011, Hebei Province, China

Zhang-Ying Feng, Department of Clinical Pharmacology, The Fourth Hospital of Hebei Medical University, Shijiazhuang 050011, Hebei Province, China

Corresponding author: Zhang-Ying Feng, PhD, Chief Pharmacist, Department of Clinical Pharmacology, The Fourth Hospital of Hebei Medical University, No. 12 Jiankang Road, Chang'an District, Shijiazhuang 050011, Hebei Province, China. fengzhangying@126.com

Abstract

BACKGROUND

Kawasaki disease (KD), also known as mucocutaneous lymph node syndrome, is an acute, self-limiting vasculitis of unknown aetiology that mainly involves the medium and small arteries and can lead to serious cardiovascular complications, with a 25% incidence of coronary artery aneurysms. Periton-Sillar abscesses are a rare symptom of KD and is easily misdiagnosed at its early stages.

CASE SUMMARY

A 5-year-old boy who presented to a community hospital with a 3-d fever, difficulty in opening his mouth, and neck pain and was originally treated for throat infection without improvement. On the basis of laboratory tests, ultrasound of submandibular and superficial lymph nodes and computed tomography of the neck, the clinician diagnosed the periamygda abscess and sepsis that did not resolve after antibiotic therapy. On the fifth day of admission, the child developed conjunctival congestion, prune tongue, perianal congestion and desquamation, and slightly stiff and swollen bunions on both feet. A diagnosis of KD was reached with complete remission after intravenous immunoglobulin treatment.

CONCLUSION

Children with neck pain, lymph node enlargement, or airway obstruction as the main manifestations are poorly treated with intravenous broad-spectrum antibiotics. Clinicians should not rush invasive operations such as neck puncture, incision, and drainage and should be alert for KD when it cannot be explained by deep neck space infection and early treatment with aspirin combined with gammaglobulin.

Key Words: Peritonsillar abscess; Kawasaki disease; Deep neck space involvement; Lymphadenitis; Gammaglobulin; Case report

©The Author(s) 2023. Published by Baishideng Publishing Group Inc. All rights reserved.

Core Tip: Some typical symptoms of a suppurative infection of the deep neck fever, for example elevated white blood cell count, lymph node enlargement, and abnormal changes in the head and neck, may also be atypical manifestations of Kawasaki disease (KD). For cases with a long course of fever and no response to antibiotic treatment, careful observation of symptoms, signs, and cervical imaging findings in the progression of the disease can be used to detect KD with deep neck changes as the first symptom early and avoid the occurrence of serious complications.

Citation: Huo LM, Li LM, Peng HY, Wang LJ, Feng ZY. Kawasaki disease with peritonsillar abscess as the first symptom: A case report. *World J Clin Cases* 2023; 11(22): 5391-5397

URL: <https://www.wjgnet.com/2307-8960/full/v11/i22/5391.htm>

DOI: <https://dx.doi.org/10.12998/wjcc.v11.i22.5391>

INTRODUCTION

Kawasaki disease (KD), also known as mucocutaneous lymph node syndrome, is an acute, self-limiting vasculitis of unknown aetiology that mainly involves the medium and small arteries and can lead to serious cardiovascular complications, with a 25% incidence of coronary artery aneurysms. Periton-Sillar abscesses are a rare symptom of KD and is easily misdiagnosed at its early stages. There are a few children with early meningitis, parotitis, changes in renal function indicators, pancreatitis, cholecystitis, intestinal obstruction[1,2]. Rare symptoms such as inflammatory changes in the parapharyngeal space or postpharyngeal space, rather than typical KD clinical manifestations, often lead to clinical misdiagnosis, delayed diagnosis and unnecessary treatment. In this paper, we report a case of KD with peri-tonsillar abscess as the first presentation and review the literature to summarise the clinical features and treatment course to improve the ability of clinicians to identify the disease early.

CASE PRESENTATION

Chief complaints

A 5-year-old boy presented with a 3-d history of fever, neck pain, and difficulty in opening his mouth. After anti-infective symptomatic treatment at the community hospital, the child's fever and sore throat did not improve. The boy was admitted to our hospital on May 20, 2021, for further treatment.

Physical examination

On admission, the patient had a febrile temperature of 40 °C, a heart rate of 100 beats/min, and a respiratory rate of 40 breaths/min. Congestion and swelling were observed in the neck and behind the ear, with significant tenderness causing the child to resist manipulation of the area. An enlarged lymph node measuring approximately 1.5 cm × 1 cm was palpable under the right jaw. The bilateral Bulbar conjunctiva was mildly congested, the pharynx was congested, tonsils were enlarged to degree II, and no herpes or exudate was seen.

Laboratory examinations

The laboratory tests showed the following: white blood cell count, $10.27 \times 10^9/L$; neutrophil percentage, 80.3%; lymphocyte percentage, 10.0%; C-reactive protein 21.49 mg/L; and erythrocyte sedimentation rate 56.00 mm/h. A blood smear showed no significant abnormalities.

Imaging examinations

An ultrasound showed cortical thickening of multiple lymph nodes adjacent to cervical vessels bilaterally, including both posterior cervical and left submandibular nodes, with the largest lymph node measuring 2.4 cm × 1.0 cm.

FINAL DIAGNOSIS

KD.

TREATMENT

On the first day of hospitalisation, intravenous antibiotic therapy was initiated with ceftriaxone, based on the suspicion of lymphadenitis and acute upper respiratory tract infection. The child had persistent high fever with chills, and computed tomography (CT) of the neck showed a widening of the retropharyngeal space with liquid hypodensity, thickening of the pharyngeal lymphatic ring, and multiple slightly large lymph nodes in the neck space (Figure 1). Based on these findings, the diagnoses were modified to peri-tonsillar abscess and sepsis, which were treated with norvancomycin. On the fifth day of admission (i.e., after more than seven days with fever), the child developed conjunctival congestion, prune tongue, perianal congestion and desquamation, and slightly stiff and swollen bunions on both feet. Based on this clinical presentation and lack of response to antibiotics, the patient was diagnosed with KD. Echocardiography performed on the same day did not reveal any coronary artery abnormalities. One dose of intravenous immunoglobulin (IVIG; 2 g/kg) was administered with rapid defervescence, and acetylsalicylic acid (4 mg/kg/day) was started and continued at home for 8 wk. On day 15, a CT scan of the neck suggested that multiple small lymph nodes in the neck were reduced in size compared to the previous film, and the oropharyngeal and retropharyngeal space effusion had disappeared (Figure 2). The child's temperature was normal, his sore throat and neck pain were relieved, and the neck swelling subsided. No coronary artery dilatation was detected on cardiac ultrasound during hospitalisation; therefore, the child was discharged in good general condition. After discharge, the boy continued to take aspirin 100 mg 1/d orally for 2 mo. The timeline of diagnosis and treatment of KD in this child (Figure 3).

OUTCOME AND FOLLOW-UP


Six months later, the child was followed-up. The family complained that the child's temperature was normal, there was no swelling or pain in the neck, and the blood count was normal. Cardiac ultrasound showed no abnormalities in the morphology, structure, or function of the heart.

DISCUSSION

Deep-neck infections occur primarily in the underlying deep cervical space formed by the deep cervical fascia, including the parapharyngeal, retropharyngeal, submandibular, and parotid spaces. Peritonsillar, parapharyngeal, and retropharyngeal involvement with oedema, cellulitis, and abscess-like lesions are rare but possible manifestations of KD. According to a cross-sectional study in the United States, approximately 0.6% of patients with KD have a combination of deep cervical space involvement[3]. A Japanese study found that 3.6% of patients with KD had CT scans suggestive of hypodense lesions in the retropharyngeal space[4]. Roh *et al*[5] found that 61% (34/56) of the cases were finally diagnosed with KD in 56 cases with fever, cervical lymphadenopathy as the main clinical manifestation, and CT enhancement of the neck suggesting retropharyngeal hypointense shadow). Another Korean study proposed that 25% of patients newly diagnosed with deep cervical space infection according to neck CT examination were diagnosed with KD[6].


The pathogenesis of KD remains unclear, and its epidemiology in recent years is highly suggestive of infection-related pathogenesis. A single-retrospective study found the presence of infection in 33% of patients with KD[7]. The microorganisms currently reported to be associated with KD include bacteria, viruses, *Mycoplasma pneumoniae*, and *Chlamydia*. The microbial toxin-like superantigen pathogenesis theory and pathogenic role of bacteria mimicking the host's own antigens may play an important role in pathogenesis. The main bacterial species associated with upper respiratory tract infections and peri-tonsillar abscesses include streptococci, *Staphylococcus aureus*, and oropharyngeal anaerobes. The microbiological mechanisms of KD and deep cervical space infections share similar pathogenic bacterial profiles. Katano *et al*[8] detected the streptococcal genomes in the cervical lymph nodes of patients with KD. Therefore, scholars have suggested that bacterial infection may be involved in KD pathogenesis with deep cervical space involvement, but no bacteria have been found in the cervical puncture drainage fluid of patients with KD combined with cervical space involvement[9-11]. In view of this, anti-infective treatment may also be necessary. However, antibiotic treatment is ineffective, and aspirin and gamma globulin are still needed to relieve the typical symptoms of KD.

The diagnostic criteria for KD include clinical features that are common to other febrile illnesses in children. KD is sometimes confused with bacterial infections, which leads to antibiotic therapy in patients with KD. In actual clinical practice, a high percentage of patients with KD receive antibiotic therapy because of empirical diagnoses and misdiagnoses. It has been reported that more than half of the patients (54.3%) received antibiotics before receiving standard treatment for KD[11]. A retrospective study reported that 33% of patients had a confirmed infection at the time of KD diagnosis, but (64%) (83/129) of patients received oral antibiotics for suspected infection prior to the diagnosis of KD[7]. When KD is combined with head and neck complications, the first symptoms are fever with limited neck movement, neck swelling and pain, slumped neck, and painful swallowing[12-16], and these clinical manifestations are very similar to those of cervical infections. Patients with KD and combined cervical involvement receive antibiotics more frequently and



DOI: 10.12998/wjcc.v11.i22.5391 Copyright ©The Author(s) 2023.

Figure 1 Neck computed tomography of the patient on the first day of hospitalisation. Computed tomography scan of the neck showed widening of the retropharyngeal space with liquid hypodensity, thickening of the pharyngeal lymphatic ring, and multiple slightly large lymph nodes in the neck space.




DOI: 10.12998/wjcc.v11.i22.5391 Copyright ©The Author(s) 2023.

Figure 2 Neck computed tomography of the patient on the fifteenth day of hospitalisation. Computed tomography scan suggested the presence of multiple small lymph nodes in the neck, which were reduced in size compared to the film scanned on the first day. Additionally, oropharyngeal and retropharyngeal space effusion were disappeared.

have a longer treatment time compared to those without combined cervical involvement[17]. However, KD coinfection does not affect the patient's response to treatment or coronary prognosis.

A peri-tonsillar abscess following acute tonsillitis is often associated with sore throat, dysphagia, restricted mouth opening, slurred pronunciation on the affected side, erythema, and elevation of the soft palate on examination[18]. In addition to clinical manifestations, enhanced CT has a higher sensitivity for the diagnosis of peritonsillar abscesses[19]. The initial presentation in our case was fever, enlarged lymph nodes and tonsils, neck pain, and difficulty in opening the mouth. CT of the cervical region showed a widening of the posterior oropharyngeal space with fluid hypodensity, thickening of the pharyngeal lymphatic ring, and multiple, slightly large lymph nodes in the cervical space. The child was considered to have a peri-tonsillar abscess, but anti-infective treatment was ineffective. It was not until the child presented with typical KD features (fever for more than seven days, visible conjunctival congestion, prune tongue, perianal congestion and desquamation, and slightly stiff and swollen bunions) that the diagnosis of KD was confirmed and treated with aspirin combined with gammaglobulin. The temperature decreased to normal after 1 d of aspirin and gammaglobulin treatment. Therefore, in infants and children with an initial diagnosis of peri-tonsillar abscess, KD needs to be considered as a possibility, especially if antibiotic therapy is not effective.

KD is a febrile multisystem vasculitis of unknown aetiology that primarily affects the coronary arteries and may result in serious cardiovascular disease if not diagnosed and treated promptly. Although antimicrobial therapy does not lead to a poor prognosis, delayed treatment and misdiagnosis may lead to poor prognosis in KD. Early recognition in clinical workup, especially in the differential diagnosis of infectious diseases, is particularly important. CT or magnetic resonance imaging (MRI) can be used for differential diagnosis. From an imaging perspective, septic infections and cellulitis changes have different presentations. In the case of septic changes, CT is mainly characterised by marginal strengthening hypointense images, whereas in patients with KD with deep neck changes, CT is mainly characterised by retropharyngeal



DOI: 10.12998/wjcc.v11.i22.5391 Copyright ©The Author(s) 2023.

Figure 3 The timeline of diagnosis and treatment of Kawasaki disease in this child.

hypointensity without marginal strengthening[20-22]. On MRI, septic neck infection is mainly characterised by central T1 hypo, T2 hyper, and peripheral marginal strengthening[23]. In some cases, CT and MRI of the cervical region can be used to differentiate deep cervical lesions of KD from bacterial septic infections.

It is difficult to distinguish patients with early KD and peri-tonsillar abscesses as the first manifestation of peritonsillar abscesses due to infection, thus leading to treatment with local incision and drainage or even tonsillectomy. Ravi and Brooks reported a 7-year-old child with clinically diagnosed tonsillar abscess who underwent right tonsillectomy on day 6 after admission and was diagnosed with KD on day 2 postoperatively, with resolution of symptoms after treatment with aspirin and intravenous gammaglobulin[24]. Cai *et al*[25] reported a 3-year-old child with parapharyngeal swelling secondary to KD who underwent parapharyngeal abscess incision and drainage on day 3 of admission, and for whom KD was diagnosed on day 8 of admission, with the symptoms resolving after treatment with aspirin and intravenous gammaglobulin. A 14-year-old male was initially diagnosed with a peri-tonsillar abscess, and the patient presented with clinical manifestations typical of KD before surgical treatment was performed[26]. All of these cases which initially lacked the typical clinical manifestations of KD, were typically treated with local surgery. Deep abscess-like lesions in the neck develop from local inflammation and oedema in the setting of systemic vasculitis, and these abscess-like lesions are absorbed and disappear with IVIG and aspirin therapy. Therefore, delayed surgical treatment is essential in these patients.

In our case, even after imaging, we had difficulty ruling out a cervical infection and instead added anti-infective treatment with norvancomycin. The diagnosis of KD was not confirmed until the child presented with typical KD features. The child received aspirin combined with gammaglobulin for one day to reduce his temperature to normal.

CONCLUSION

Children with neck pain, lymph node enlargement, or airway obstruction as the main manifestations are poorly treated with intravenous broad-spectrum antibiotics. Clinicians should not rush invasive operations such as neck puncture, incision, and drainage and should be alert for KD when it cannot be explained by deep neck space infection or early treatment with aspirin combined with gammaglobulin.

FOOTNOTES

Author contributions: Feng ZY conceived of the study and revised manuscript; Huo LM and Li LM collected the data and wrote the manuscript; Peng HY and Wang LJ analyzed the corresponding ultrasound and CT results and wrote the manuscript.

Informed consent statement: Written informed consent was obtained from the patient's parent for publication of this case report and any accompanying images.

Conflict-of-interest statement: The authors declare that they have no conflict of interest.

CARE Checklist (2016) statement: The authors have read the CARE Checklist (2016), and the manuscript was prepared and revised according to the CARE Checklist (2016).

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers.

It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <https://creativecommons.org/licenses/by-nc/4.0/>

Country/Territory of origin: China

ORCID number: Li-Man Huo 0000-0003-0078-4047; Li-Min Li 0009-0004-7963-3640; Zhang-Ying Feng 0000-0002-7881-6937.

S-Editor: Yan JP

L-Editor: A

P-Editor: Yan JP

REFERENCES

- 1 **Zhang B**, Hao Y, Zhang Y, Yang N, Li H, Liang J. Kawasaki disease manifesting as bilateral facial nerve palsy and meningitis: a case report and literature review. *J Int Med Res* 2019; **47**: 4014-4018 [PMID: 31364426 DOI: 10.1177/0300060519854287]
- 2 **Lai CC**, Lin WT, Lin HC. Parotitis: An Initial Manifestation of Kawasaki Disease. *J Pediatr* 2019; **214**: 235-235.e1 [PMID: 31378519 DOI: 10.1016/j.jpeds.2019.06.067]
- 3 **Inagaki K**, Blackshear C, Hobbs CV. Deep Neck Space Involvement of Kawasaki Disease in the US: A Population-Based Study. *J Pediatr* 2019; **215**: 118-122 [PMID: 31477383 DOI: 10.1016/j.jpeds.2019.07.054]
- 4 **Tona R**, Shinohara S, Fujiwara K, Kikuchi M, Kanazawa Y, Kishimoto I, Harada H, Naito Y. Risk factors for retropharyngeal cellulitis in Kawasaki disease. *Auris Nasus Larynx* 2014; **41**: 455-458 [PMID: 24958366 DOI: 10.1016/j.anl.2014.05.017]
- 5 **Roh K**, Lee SW, Yoo J. CT analysis of retropharyngeal abnormality in Kawasaki disease. *Korean J Radiol* 2011; **12**: 700-707 [PMID: 22043152 DOI: 10.3348/kjr.2011.12.6.700]
- 6 **Lim S**, Lee NY, Han SB, Jeong DC, Kang JH. Deep Neck Inflammation: Probable Kawasaki Disease in Korean Children. *Clin Exp Otorhinolaryngol* 2020; **13**: 77-82 [PMID: 31599139 DOI: 10.21053/ceo.2019.00948]
- 7 **Benseler SM**, McCrindle BW, Silverman ED, Tyrrell PN, Wong J, Yeung RS. Infections and Kawasaki disease: implications for coronary artery outcome. *Pediatrics* 2005; **116**: e760-e766 [PMID: 16322132 DOI: 10.1542/peds.2005-0559]
- 8 **Katano H**, Sato S, Sekizuka T, Kinumaki A, Fukumoto H, Sato Y, Hasegawa H, Morikawa S, Saijo M, Mizutani T, Kuroda M. Pathogenic characterization of a cervical lymph node derived from a patient with Kawasaki disease. *Int J Clin Exp Pathol* 2012; **5**: 814-823 [PMID: 23071864]
- 9 **Pontell J**, Rosenfeld RM, Kohn B. Kawasaki disease mimicking retropharyngeal abscess. *Otolaryngol Head Neck Surg* 1994; **110**: 428-430 [PMID: 8170688 DOI: 10.1177/019459989411000413]
- 10 **Principi N**, Esposito S. Antibiotic-related adverse events in paediatrics: unique characteristics. *Expert Opin Drug Saf* 2019; **18**: 795-802 [PMID: 31305171 DOI: 10.1080/14740338.2019.1640678]
- 11 **Langley EW**, Kirse DK, Barnes CE, Covitz W, Shetty AK. Retropharyngeal edema: an unusual manifestation of Kawasaki disease. *J Emerg Med* 2010; **39**: 181-185 [PMID: 19150196 DOI: 10.1016/j.jemermed.2008.08.004]
- 12 **Isidori C**, Sebastiani L, Esposito S. A Case of Incomplete and Atypical Kawasaki Disease Presenting with Retropharyngeal Involvement. *Int J Environ Res Public Health* 2019; **16** [PMID: 31491922 DOI: 10.3390/ijerph16183262]
- 13 **Aldemir-Kocabaş B**, Kcal MM, Ramoğlu MG, Tutar E, Fitöz S, Çiftçi E, İnce E. Recurrent Kawasaki disease in a child with retropharyngeal involvement: a case report and literature review. *Medicine (Baltimore)* 2014; **93**: e139 [PMID: 25546657 DOI: 10.1097/MD.0000000000000139]
- 14 **Hung MC**, Wu KG, Hwang B, Lee PC, Meng CC. Kawasaki disease resembling a retropharyngeal abscess--case report and literature review. *Int J Cardiol* 2007; **115**: e94-e96 [PMID: 17126927 DOI: 10.1016/j.ijcard.2006.08.095]
- 15 **Homicz MR**, Carvalho D, Kearns DB, Edmonds J. An atypical presentation of Kawasaki disease resembling a retropharyngeal abscess. *Int J Pediatr Otorhinolaryngol* 2000; **54**: 45-49 [PMID: 10960696 DOI: 10.1016/s0165-5876(00)00337-2]
- 16 **Kasem Ali Sliman R**, van Montfrans JM, Nassrallah N, Hamad Saied M. Retropharyngeal abscess-like as an atypical presentation of Kawasaki disease: a case report and literature review. *Pediatr Rheumatol Online J* 2023; **21**: 34 [PMID: 37046311 DOI: 10.1186/s12969-023-00812-z]
- 17 **Cheng XY**. Analysis of clinical features of Kawasaki disease combined with deep cervical space involvement. Master Thesis, China Medical University, China, 2021 [DOI: 10.27652/d.cnki.gzyku.2021.001131]
- 18 **AlAwadh I**, Aldrees T, AlQaryan S, Alharethy S, AlShehri H. Bilateral peritonsillar abscess: A case report and pertinent literature review. *Int J Surg Case Rep* 2017; **36**: 34-37 [PMID: 28531866 DOI: 10.1016/j.ijscr.2017.04.028]
- 19 **Baker KA**, Stuart J, Sykes KJ, Sinclair KA, Wei JL. Use of computed tomography in the emergency department for the diagnosis of pediatric peritonsillar abscess. *Pediatr Emerg Care* 2012; **28**: 962-965 [PMID: 23023458 DOI: 10.1097/PEC.0b013e31826c6c36]
- 20 **Nomura O**, Hashimoto N, Ishiguro A, Miyasaka M, Nosaka S, Oana S, Sakai H, Takayama JI. Comparison of patients with Kawasaki disease with retropharyngeal edema and patients with retropharyngeal abscess. *Eur J Pediatr* 2014; **173**: 381-386 [PMID: 24146166 DOI: 10.1007/s00431-013-2179-0]
- 21 **Vural C**, Gungor A, Comerçi S. Accuracy of computerized tomography in deep neck infections in the pediatric population. *Am J Otolaryngol* 2003; **24**: 143-148 [PMID: 12761699 DOI: 10.1016/s0196-0709(03)00008-5]
- 22 **Tsujioka Y**, Handa A, Nishimura G, Miura M, Yokoyama K, Sato K, Handa H, Jinzaki M, Nosaka S, Kono T. Multisystem Imaging Manifestations of Kawasaki Disease. *Radiographics* 2022; **42**: 268-288 [PMID: 34826255 DOI: 10.1148/rg.210070]
- 23 **Chiappini E**, Camaioni A, Benazzo M, Biondi A, Bottero S, De Masi S, Di Mauro G, Doria M, Esposito S, Felisati G, Felisati D, Festini F, Gaini RM, Galli L, Gambini C, Gianelli U, Landi M, Lucioni M, Mansi N, Mazzantini R, Marchisio P, Marseglia GL, Miniello VL, Nicola M, Novelli A, Paulli M, Picca M, Pillon M, Pisani P, Pipolo C, Principi N, Sardi I, Succo G, Tomà P, Tortoli E, Tucci F, Varricchio A, de Martino M; Italian Guideline Panel For Management Of Cervical Lymphadenopathy In Children. Development of an algorithm for the management of cervical lymphadenopathy in children: consensus of the Italian Society of Preventive and Social Pediatrics, jointly with the Italian Society of

Pediatric Infectious Diseases and the Italian Society of Pediatric Otorhinolaryngology. *Expert Rev Anti Infect Ther* 2015; **13**: 1557-1567 [PMID: 26558951 DOI: 10.1586/14787210.2015.1096777]

- 24 **Ravi KV**, Brooks JR. Peritonsillar abscess--an unusual presentation of Kawasaki disease. *J Laryngol Otol* 1997; **111**: 73-74 [PMID: 9292139 DOI: 10.1017/s0022215100136485]
- 25 **Cai Q**, Luo R, Gan J, Zhang L, Qu Y, Mu D. Kawasaki disease mimicking a parapharyngeal abscess: a case report. *Medicine (Baltimore)* 2015; **94**: e761 [PMID: 25929913 DOI: 10.1097/MD.0000000000000761]
- 26 **Connell JT**, Park JH. Acute peritonsillar swelling: a unique presentation for Kawasaki disease in adolescence. *BMJ Case Rep* 2018; **2018** [PMID: 30042103 DOI: 10.1136/bcr-2018-224441]



Published by **Baishideng Publishing Group Inc**
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA

Telephone: +1-925-3991568

E-mail: bpgoffice@wjgnet.com

Help Desk: <https://www.f6publishing.com/helpdesk>

<https://www.wjgnet.com>

