

Fatal hemophagocytic lymphohistiocytosis-induced multiorgan dysfunction secondary to *Burkholderia pseudomallei* sepsis: A case report

Ming-Ze Sui, Ke-Cheng Wan, Yuan-Lu Chen, Huan-Long Li, Shan-Shan Wang, Ze-Fu Chen

Specialty type: Medicine, research and experimental

Provenance and peer review:

Unsolicited article; Externally peer reviewed.

Peer-review model: Single blind

Peer-review report's scientific quality classification

Grade A (Excellent): 0

Grade B (Very good): 0

Grade C (Good): C

Grade D (Fair): 0

Grade E (Poor): 0

P-Reviewer: Moldovan CA, Romania

Received: September 4, 2023

Peer-review started: September 4, 2023

First decision: September 13, 2023

Revised: September 24, 2023

Accepted: October 8, 2023

Article in press: October 8, 2023

Published online: October 26, 2023



Ming-Ze Sui, Department of Pulmonary and Critical Care Medicine, Kunming Children's Hospital, Yunnan Key Laboratory of Children's Major Disease Research, Kunming 650034, Yunnan Province, China

Ke-Cheng Wan, Yuan-Lu Chen, Huan-Long Li, Shan-Shan Wang, Ze-Fu Chen, Department of Pediatrics, Hainan General Hospital, Haikou 570311, Hainan Province, China

Corresponding author: Ze-Fu Chen, BMed, Attending Doctor, Department of Pediatrics, Hainan General Hospital, Xiuhua Road, Haikou 570311, Hainan Province, China. 42648804@hainmc.edu.cn

Abstract

BACKGROUND

Burkholderia pseudomallei (*B. pseudomallei*) is a short, straight, medium-sized Gram-negative bacterium that mostly exists alone, without a capsule or spores, has more than three flagella at one end, and actively moves. *B. pseudomallei* confers high morbidity and mortality, with frequent granulocytopenia in *B. pseudomallei* sepsis-related deaths. However, mortality may be related to hemophagocytic lymphohistiocytosis (HLH) secondary to *B. pseudomallei* infection.

CASE SUMMARY

A 12-year-old female was referred from a local hospital to the pediatric intensive care unit with suspected septic shock and fever, cough, dyspnea, and malaise. After admission, supportive symptomatic treatments including fluid resuscitation, anti-infective therapy, mechanical ventilation, and a vasoactive drug maintenance cycle were carefully initiated. The patient became unconscious, her blood pressure could not be maintained even under the exposure of vasoactive drugs, and she experienced cardiorespiratory arrest. The patient died due to ineffective high-quality in-hospital cardiopulmonary resuscitation. A subsequent bone marrow smear examination revealed extensive phagocytosis, and the blood culture was positive for *B. pseudomallei*. Family history revealed a sibling death from *B. pseudomallei* sepsis 5 years earlier.

CONCLUSION

The higher mortality rate in patients with *B. pseudomallei* sepsis may be related to secondary HLH after infection, wherein multiorgan dysfunction syndrome may be directly related to infection or immune damage caused by secondary HLH.

Patients with *B. pseudomallei* can be asymptomatic and can become an infective source.

Key Words: *Burkholderia pseudomallei*; Sepsis; Septic shock; Hemophagocytic lymphohistiocytosis; Asymptomatic carrier; Case report

©The Author(s) 2023. Published by Baishideng Publishing Group Inc. All rights reserved.

Core Tip: Given the high mortality rate associated with *Burkholderia pseudomallei* (*B. pseudomallei*), it is particularly important to fully understand the pathogenesis. This report presents the clinical characteristics of a case of *B. pseudomallei* infection and some clinical data of the patient's brother, who also died from *B. pseudomallei* infection. The chronic carrier status of *B. pseudomallei* and secondary hemophagocytic lymphohistiocytosis warrants attention in research on the pathogenesis and treatment of *B. pseudomallei* sepsis.

Citation: Sui MZ, Wan KC, Chen YL, Li HL, Wang SS, Chen ZF. Fatal hemophagocytic lymphohistiocytosis-induced multiorgan dysfunction secondary to *Burkholderia pseudomallei* sepsis: A case report. *World J Clin Cases* 2023; 11(30): 7372-7379

URL: <https://www.wjgnet.com/2307-8960/full/v11/i30/7372.htm>

DOI: <https://dx.doi.org/10.12998/wjcc.v11.i30.7372>

INTRODUCTION

Burkholderia pseudomallei (*B. pseudomallei*) is a non-fermentative Gram-negative bacterium that is positive for oxidases and enzymes, does not form spores, and does not contain metachromatic particles. Approximately 165000 cases of *B. pseudomallei* infection and 89000 deaths are reported annually worldwide. The incidence rates in South and East Asia and the Pacific are 44% and 40%, respectively, and the mortality rates are 47% and 35%, respectively[1].

Sequential dysfunction of two or more organs is usually referred to as multiorgan dysfunction syndrome (MODS). MODS caused by sepsis likely contributes to the high mortality rates associated with *B. pseudomallei* infections. However, granulocytopenia, which is common in areas where such cases have been reported, has been largely neglected by the research community. Combined with the clinical data on elevated ferritin levels, there is a need to examine the status of hemophagocytosis in deaths due to *B. pseudomallei*[2].

Herein, we have described the clinical characteristics of a patient infected with *B. pseudomallei* and the clinical data of her brother, who died from the same illness.

CASE PRESENTATION

Chief complaints

A 12-year-old female was transferred from a local hospital to our pediatric intensive care unit after 4 d of fever, cough for 2 d, and dyspnea and malaise for 1 d.

History of present illness

The patient had originally presented to the local clinic with a very high fever ($> 40.0^{\circ}\text{C}$) and had been administered oral antibiotic treatment after a routine blood examination; however, the patient's parents were unaware of the type of antibiotics administered. After treatment, the patient's high fever continued and she developed a cough and other new symptoms. She then presented to a local hospital, where routine blood examination indicated agranulocytosis and blood cell and platelet levels that were significantly lower than those in the last test. After receiving ceftazidime at the hospital, the body temperature remained high and dyspnea and fatigue persisted. Arterial blood gas analysis suggested lactic acidosis, indicating that the patient was experiencing consolidated septic shock.

History of past illness

The patient's parents denied a history of hepatitis, tuberculosis, measles, mumps, or other common infectious diseases. The patient had no history of surgery, trauma, or blood transfusions.

Personal and family history

The patient did not have a history of preterm delivery, birth asphyxia, intrauterine hypoxia, intrauterine conditions, or infection. The patient's brother had died 5 years earlier due to *B. pseudomallei* sepsis and septic shock (results of laboratory tests are shown in Table 1).

Table 1 Laboratory data of the patient's brother

Variable	Reference range	Initial value
Routine blood examination		
White cell count as $\times 10^9/L$	4.30-11.30	0.60
Neutrophil count as $\times 10^9/L$	1.60-7.80	0.40
Hemoglobin in g/L	118-156	104
Platelet count as $\times 10^9/L$	150-407	184
Inflammatory indicators		
C-reactive protein in mg/L	0-8.00	200.00
Procalcitonin in ng/mL	0-0.046	98.670
Erythrocyte sedimentation rate in mm/h	0-15	33
Arterial blood gas analysis		
pH	7.350-7.450	7.076
PaO ₂ in mmHg	83.0-108.0	43.4
PaCO ₂ in mmHg	35-45	37
HCO ₃ ⁻ in mmol/L	22.0-27.0	12.5
BE in mmol/	-3.0 to 3.0	-19.0
Lactic acid in mmol/L	0.5-1.7	13.1
Coagulative function		
Prothrombin time in s	9.8-13.2	21.1
International normalized ratio	0.85-1.20	1.64
D-dimer in ng/mL	0-0.50	4209.00
Biochemical examination		
Glutamic pyruvic transaminase in U/L	7-30	43
Albumin in g/L	39.0-54.0	27.4
Total bilirubin in $\mu\text{mol/L}$	0-21.0	27.2
Direct bilirubin in $\mu\text{mol/L}$	0-8.0	20.1
Urea nitrogen in mmol/L	2.5-6.5	8.8

BE: Base excess.

Physical examination

Body temperature, respiratory rate, and heart rate were 37.5 °C, 34 times/min, and 146 beats/min, respectively. Furthermore, the body weight was 40 kg, and skin oxygen saturation was 87% in room air. The patient's mentality was depressed. Both pupils were equal in size and responsive to light. The nasal wings were flapped, the lips were cyanotic, breathing was rapid, the three concave signs were positive, and auscultation revealed dense moist rales. Heartbeat sounds were low. The liver and spleen were palpable under the costal margin (1.5 cm below the right midclavicular costal margin and 1.5 cm below the left midclavicular costal margin, respectively). The extremities were cold, and capillary filling time was 6 s; however, no obvious abnormality was observed in the neurological examination.

Laboratory examinations

Routine blood tests revealed agranulocytosis, thrombocytopenia, and anemia. Arterial blood gas analysis revealed sustained hypoxia and acidosis. Monitoring of coagulative parameters indicated hypofibrinogenemia. The levels of C-reactive protein, procalcitonin, erythrocyte sedimentation rate, and other inflammatory indicators significantly increased. A complete biochemical examination revealed varying degrees of multiorgan dysfunction. The levels of interleukin and inflammatory factors, such as ferritin, were higher than normal, and no biomarkers related to hemophagocytic syndrome were found in the whole-exon test (described in Table 2).

Table 2 Laboratory data of the patient

Variable	Reference range	Initial value	Value retested after 4 h	Last value before death
Routine blood examination				
White cell count as $\times 10^9/L$	4.30-11.30	0.69	0.83	-
Neutrophil count as $\times 10^9/L$	1.60-7.80	0.08	0.08	-
Hemoglobin in g/L	118-156	112	83	-
Platelet count as $\times 10^9/L$	150-407	50	23	-
Inflammatory indicators				
C-reactive protein in mg/L	0-8.00	239.04	135.83	-
Procalcitonin in ng/mL	0-0.046	73.540	-	-
Erythrocyte sedimentation rate in mm/h	0-20	8	-	-
Ferritin in ng/mL	4.63-204.00	14586.00	-	-
Arterial blood gas analysis				
pH	7.350-7.450	7.150	7.230	6.840
PaO ₂ in mmHg	83.0-108.0	69.0	73.0	22.0
PaCO ₂ in mmHg	35-45	28	28	88
BE in mmol/L	-3.0 to 3.0	-17.6	-14.5	-19.2
Oxygenation index	-	191	202	36
Lactic acid in mmol/L	0.5-1.7	15.8	14.5	16.7
Coagulation function				
Prothrombin time in s	9.8-13.2	22.9	53.6	-
International normalized ratio	0.85-1.20	1.97	5.99	-
Thrombin time in s	14.0-21.0	14.7	240.0	-
Activated partial thromboplastin time in s	28.0-43.0	58.1	180.0	-
Fibrinogen in g/L	2.00-4	-	1.56	-
D-dimer in ng/mL	0-0.50	24.93	11.43	-
Biochemical examination				
Glutamic pyruvic transaminase in U/L	7.0-30.0	101.4	192.6	-
Albumin in g/L	39.0-54.0	27.3	19.7	-
Total bilirubin in $\mu\text{mol/L}$	0-21.00	89.07	54.83	-
Direct bilirubin in $\mu\text{mol/L}$	0-8.00	51.46	31.99	-
Indirect bilirubin in $\mu\text{mol/L}$	0-21.00	37.61	22.84	-
Urea nitrogen in $\mu\text{mol/L}$	2.50-6.50	10.75	11.02	-
Creatinine in $\mu\text{mol/L}$	27-66	157	82	-
K ⁺ in mmol/L	3.50-5.50	3.41	2.65	-
Na ⁺ in mmol/L	130.0-150.0	135.9	142.6	-
Ca ²⁺ in mmol/L	2.10-2.80	1.76	1.45	-
Triglyceride in mmol/L	0-1.70	1.57	-	-
Cytokines in pg/mL				
Interleukin-1B	0-12.40	623.47	-	-
Interleukin-6	0-5.30	1452.90	-	-
Interleukin-8	0-53.09	1876.33	-	-
Interleukin-10	0-4.91	5754.38	-	-

Interferon- γ	0-7.42	3143.99	-	-
Tumor necrosis factor	0-4.60	42.28	-	-

BE: Base excess.

Imaging examinations

Chest radiography revealed exudative lesions in both lungs (Figure 1).

Further diagnostic analysis

A bone marrow smear showed hemophagocytosis (Figure 2).

FINAL DIAGNOSIS

The patient was finally diagnosed with *B. pseudomallei* sepsis, septic shock, MODS, acute respiratory distress syndrome, respiratory failure, severe pneumonia, metabolic acidosis, disseminated intravascular coagulation, electrolyte metabolism disorder, agranulocytosis, and thrombocytopenia, with a high suspicion of secondary hemophagocytic lymphohistiocytosis (HLH).

TREATMENT

Meropenem was initiated. The patient immediately received a 2:1 isotonic solution to expand the volume, twice along with nasal catheter oxygen inhalation (oxygen flow ≤ 5 L/min). We established a femoral vein infusion path to simultaneously implement subsequent fluid support. As the blood pressure of the patient could not be maintained after volume expansion and the analysis of arterial blood gas indicated continuous hypoxia due to irregular spontaneous respiration, we administered norepinephrine 0.5 $\mu\text{g/kg/min}$ continuous pumping to maintain blood pressure and invasive mechanical ventilation with endotracheal intubation (PC-SIMV mode: FiO_2 60%, VT 300 mL, PEEP 5 cmH_2O , RR25 times/min). Supportive treatments, such as granulocyte-stimulating factor, plasma, coagulation factor cryoprecipitate, and suspended red blood cells, were also administered.

Nevertheless, the patient experienced respiratory and cardiac arrest 4 h after admission, and a pink foam-like liquid gushed from the endotracheal tube. After cardiopulmonary resuscitation and intravenous morphine administration, the patient's heart rate recovered, but the results of arterial blood gas analysis worsened (Table 2). In addition to hypoxia, the patient had serious carbon dioxide retention; we changed the invasive mechanical ventilation mode to high-frequency mode (FiO_2 60%, average airway pressure 35 cmH_2O , amplitude 75 cmH_2O , sighing time 0.3s, frequency 7 Hz).

After the above rescue, the patient's condition continued to deteriorate, and norepinephrine (1 $\mu\text{g/kg/min}$) combined with dopamine (6 $\mu\text{g/kg/min}$) still failed to maintain normal blood pressure. The patient's consciousness gradually turned into coma, and diffuse bleeding spots appeared over the patient's entire body. We urgently punctured the bone marrow, injected vitamin K1, ethylphenesulfonate, and snake venom hemocoagulase to stop bleeding and added m-hydroxylamine to raise the blood pressure. At the same time, we actively administered bedside blood purification treatment and adjusted norepinephrine to 1.4 $\mu\text{g/kg/min}$, dopamine to 12 $\mu\text{g/kg/min}$, and m-hydroxylamine to 2 $\mu\text{g/kg/min}$ during continuous renal replacement therapy. The patient's blood pressure remained unstable, and the maintenance of transcutaneous oxygen saturation was unsatisfactory. After 12 h of hospitalization, the patient died.

OUTCOME AND FOLLOW-UP

The patient eventually died, and blood culture was positive for *B. pseudomallei*. The patient's living brother also had a fever at the same time. The parents requested that the patient's brother be hospitalized. Based on the patient's brother's blood culture and drug sensitivity results, imipenem was administered to him. Finally, the patient's brother was discharged with a normal body temperature, and no pathogenic bacterial growth was observed in subsequent blood cultures.


DISCUSSION

HLH is a macrophage proliferative disease mostly caused by Epstein-Barr virus infection, whereas HLH caused by bacterial infections is relatively rare. According to the HLH-2004 guidelines, the diagnosis of HLH needs to meet five of the eight diagnostic criteria: (1) Fever; (2) spleen enlargement; (3) decrease of peripheral blood cells, involving 2-3 lines, that is hemoglobin < 90 g/L, platelet count $< 100 \times 10^9/\text{L}$, and neutrophils $< 1.0 \times 10^9/\text{L}$; (4) hypertriglyceridemia and/or



DOI: 10.12998/wjcc.v11.i30.7372 Copyright ©The Author(s) 2023.

Figure 1 Chest X-ray showed that the density of both lung fields was increased, suggesting the possibility of exudative lesions and suspected bilateral pleural effusion. The endotracheal intubation head was at the lower edge of the thoracic cone 2.



DOI: 10.12998/wjcc.v11.i30.7372 Copyright ©The Author(s) 2023.

Figure 2 Bone marrow smear showing phagocytic cells, phagocytic erythrocytes, and platelets. A: Macrophages engulfing erythrocytes; B: Macrophages engulfing platelets.

low fibrinogen, with fasting triglyceride ≥ 3.0 mmol/L (≥ 2.65 g/L) and fibrinogen ≤ 1.5 g/L; (5) blood phagocytic cells in the bone marrow, spleen, or lymph nodes, and no evidence of malignant tumor; (6) activity of natural killer cells being reduced or completely absent; (7) serum ferritin ≥ 500 $\mu\text{g/L}$; and (8) soluble CD25 (interleukin-2 receptor) ≥ 2400 U/mL [3]. According to the guidelines for fever, a temperature $\geq 38.5^\circ\text{C}$ for more than 7 d is required. The patient reported in this article was declared clinically dead on the 5th d of the fever. Therefore, the patient's clinical data only met criteria 2, 3, 5, and 7. Unfortunately, the whole-exon test failed to identify the molecular biological markers supporting HLH. Many indicators cannot be reviewed or improved over time because of the rapid worsening of a patient's condition. Although the clinical diagnosis of HLH was not confirmed, the results of the patient's bone marrow examination and MODS caused by HLH cannot be ignored among the many factors that lead to the patient's death.

In addition, case reports of focal infections have shown that most patients have a good prognosis, and the examination indicators for these patients are quite different from the clinical manifestations of HLH[4-6]. Systemic infections are not complicated by MODS[2]. Therefore, the final progression of secondary HLH to *B. pseudomallei* sepsis contributes to patient mortality, and *B. pseudomallei* infection is mainly observed during the rainy season in tropical and subtropical regions. The seasonal incidence may be related to the survival of bacteria in the soil. Knowledge of the history of contact between pestilence-related soil and water sources is particularly important for early treatment by doctors.

Several preclinical studies have shown that the lungs, liver, and spleen are the most common target organs for chronic *B. pseudomallei* infection[7]. Studies have shown that *B. pseudomallei* regulates phagocytic death and aids in the progression of acute or chronic infections[8]. After *B. pseudomallei* invade the body, they recruit host complement regulatory proteins for immune evasion[9]. Furthermore, bacteria can survive in small abscesses formed in target organs. Over time, the chronic infection site forms granulomas with neutrophils, macrophages, and lymphocytes as the main components[10]. During this process, the patient's symptoms gradually improve, which is often considered a clinical cure. However, this patient was a chronic disease carrier. The patient reported in this article did not have any history of living in an endemic area. Based on the patient's brother's history, we suspected that the source of infection was an asymptomatic carrier who came into contact with the patient.

Most *B. pseudomallei* strains are sensitive to imipenem. The most important step is to identify the causative pathogen as soon as possible and provide effective interventions before severe clinical events occur. Early diagnosis based on bioinformatic analysis may help solve these problems[11]. Future vaccine development and bacteriophage therapy will help to reduce the incidence and mortality of *B. pseudomallei*[12,13].

CONCLUSION

A person infected with *B. pseudomallei* may be an asymptomatic carrier owing to the unique mechanism of chronic *B. pseudomallei* infection. The high fatality rate of *B. pseudomallei* may be related to MODS caused by secondary HLH, as observed in this case.

FOOTNOTES

Author contributions: Sui MZ and Wan KC contributed equally to this work; Sui MZ wrote the original draft of the manuscript; Wan KC performed the analysis and interpretation of imaging findings; Chen YL and Li HL were responsible for the treatment of the patient and provided clinical data; Wang SS was responsible for literature retrieval; Chen ZF was responsible for the revision and editing of the manuscript; All authors approved the final version to be submitted.

Informed consent statement: The patient's parents signed a written informed consent form.

Conflict-of-interest statement: The authors declare having no conflicts of interest.

CARE Checklist (2016) statement: The authors read the CARE Checklist (2016) and the manuscript was prepared and revised according to it Checklist (2016).

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <https://creativecommons.org/licenses/by-nc/4.0/>

Country/Territory of origin: China

ORCID number: Ming-Ze Sui 0000-0002-1424-7384; Ke-Cheng Wan 0000-0001-9856-5509; Yuan-Lu Chen 0000-0002-4515-7824; Huan-Long Li 0000-0002-3849-7038; Shan-Shan Wang 0000-0002-0789-9958; Ze-Fu Chen 0000-0003-4785-4882.

S-Editor: Lin C

L-Editor: Filipodia

P-Editor: Yu HG

REFERENCES

- 1 Limmathurotsakul D, Golding N, Dance DA, Messina JP, Pigott DM, Moyes CL, Rolim DB, Bertherat E, Day NP, Peacock SJ, Hay SI. Predicted global distribution of *Burkholderia pseudomallei* and burden of melioidosis. *Nat Microbiol* 2016; **1** [PMID: 26877885 DOI: 10.1038/nmicrobiol.2015.8]
- 2 Bhaskaran P, Prasad V, Gopinathan A, Shaw T, Sivadas S, Jayakumar C, Chowdhury S, Dravid A, Mukhopadhyay C, Kumar A. *Burkholderia pseudomallei* in Environment of Adolescent Siblings with Melioidosis, Kerala, India, 2019. *Emerg Infect Dis* 2022; **28**: 1246-1249 [PMID: 35608830 DOI: 10.3201/eid2806.211526]

- 3 **Henter JI**, Horne A, Aricó M, Egeler RM, Filipovich AH, Imashuku S, Ladisch S, McClain K, Webb D, Winiarski J, Janka G. HLH-2004: Diagnostic and therapeutic guidelines for hemophagocytic lymphohistiocytosis. *Pediatr Blood Cancer* 2007; **48**: 124-131 [PMID: [16937360](#) DOI: [10.1002/pbc.21039](#)]
- 4 **Dimitriou NG**, Flüh G, Zange S, Aytulun A, Turowski B, Hartung HP, Meuth SG, Gliem M. Case report: First case of neuromelioidosis in Europe: CNS infection caused by *Burkholderia pseudomallei*. *Front Neurol* 2022; **13**: 899396 [PMID: [35968288](#) DOI: [10.3389/fneur.2022.899396](#)]
- 5 **Santos Júnior ELD**, Moura JCR, Protásio BKPF, Parente VAS, Veiga MHND. Clinical repercussions of Glanders (*Burkholderia mallei* infection) in a Brazilian child: a case report. *Rev Soc Bras Med Trop* 2020; **53**: e20200054 [PMID: [32578712](#) DOI: [10.1590/0037-8682-0054-2020](#)]
- 6 **Verdecia JL**, Jankowski CA, Isache CL, Neilsen CD, McCarter YS, Sands ML, Ravi M. A Case Report of Melioidotic Prostatic Abscess in a Traveler. *Open Forum Infect Dis* 2022; **9**: ofac284 [PMID: [35891686](#) DOI: [10.1093/ofid/ofac284](#)]
- 7 **Amemiya K**, Dankmeyer JL, Fetterer DP, Worsham PL, Welkos SL, Cote CK. Comparison of the early host immune response to two widely diverse virulent strains of *Burkholderia pseudomallei* that cause acute or chronic infections in BALB/c mice. *Microb Pathog* 2015; **86**: 53-63 [PMID: [26162294](#) DOI: [10.1016/j.micpath.2015.07.004](#)]
- 8 **Krakauer T**. Living dangerously: *Burkholderia pseudomallei* modulates phagocyte cell death to survive. *Med Hypotheses* 2018; **121**: 64-69 [PMID: [30396496](#) DOI: [10.1016/j.mehy.2018.09.028](#)]
- 9 **Syed I**, Wooten RM. Interactions Between Pathogenic *Burkholderia* and the Complement System: A Review of Potential Immune Evasion Mechanisms. *Front Cell Infect Microbiol* 2021; **11**: 701362 [PMID: [34660335](#) DOI: [10.3389/fcimb.2021.701362](#)]
- 10 **Lewis ER**, Torres AG. The art of persistence-the secrets to *Burkholderia* chronic infections. *Pathog Dis* 2016; **74** [PMID: [27440810](#) DOI: [10.1093/femspd/ftw070](#)]
- 11 **Yin L**, Chen Y, Fu T, Liu L, Xia Q. Identification of candidate blood biomarkers for the diagnosis of septicaemic melioidosis based on WGCNA. *Artif Cells Nanomed Biotechnol* 2022; **50**: 252-259 [PMID: [36191125](#) DOI: [10.1080/21691401.2022.2126490](#)]
- 12 **Lauman P**, Dennis JJ. Advances in Phage Therapy: Targeting the *Burkholderia cepacia* Complex. *Viruses* 2021; **13** [PMID: [34372537](#) DOI: [10.3390/v13071331](#)]
- 13 **Wang G**, Zarodkiewicz P, Valvano MA. Current Advances in *Burkholderia* Vaccines Development. *Cells* 2020; **9** [PMID: [33322641](#) DOI: [10.3390/cells9122671](#)]



Published by **Baishideng Publishing Group Inc**
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA

Telephone: +1-925-3991568

E-mail: bpgoffice@wjgnet.com

Help Desk: <https://www.f6publishing.com/helpdesk>

<https://www.wjgnet.com>

