



Fournier gangrene in an infant, complicated with severe sepsis and liver dysfunction: A case report

Ilijana Bakalli, Saimir Heta, Ermira Kola, Ermela Celaj

Specialty type: Medicine, research and experimental

Provenance and peer review: Unsolicited article; Externally peer reviewed.

Peer-review model: Single blind

Peer-review report's scientific quality classification

Grade A (Excellent): 0
Grade B (Very good): 0
Grade C (Good): C, C, C
Grade D (Fair): 0
Grade E (Poor): 0

P-Reviewer: Leowattana W, Thailand; Manrai M, India

Received: July 2, 2023

Peer-review started: July 2, 2023

First decision: July 28, 2023

Revised: August 9, 2023

Accepted: September 6, 2023

Article in press: September 6, 2023

Published online: October 26, 2023



Ilijana Bakalli, Ermira Kola, Ermela Celaj, Department of Pediatrics, UHC “Mother Theresa”, Tirana 1024, Albania

Saimir Heta, Department of Pediatric Surgery, UHC “Mother Theresa”, Tirana 1024, Albania

Corresponding author: Ilijana Bakalli, PhD, Associate Professor, Department of Pediatrics, UHC “Mother Theresa”, Rruga e Dibres, 376, Tirana 1024, Albania.

ilijanabakalli@yahoo.com

Abstract

BACKGROUND

Fournier gangrene is a rare, life-threatening infection characterized by necrotizing fasciitis in the perineal, genital and/or lower abdominal regions. Despite its rarity, the unfavorable prognosis associated with this disease is dependent on the timing of medical care.

CASE SUMMARY

A 3-month-old boy was admitted to our pediatric intensive care unit in critical condition after a 5-day history of fever and scrotal erythema with breaching skin lesions and swelling. Despite ambulatory antibiotic treatment, the child's clinical condition deteriorated. At the time of admission, the child had necrotizing scrotal fasciitis that had spread to the abdomen. Following reanimation, the surgeon decided on an immediate intervention to rule out testicular torsion and to debride the affected area. Despite optimal antibiotic and supportive therapy, the patient developed severe sepsis with liver dysfunction, making treatment more challenging.

CONCLUSION

Recognizing Fournier gangrene, prompt referral to pediatric surgery, and appropriate antibiotic coverage are critical for avoiding sepsis and multiorgan dysfunction.

Key Words: Fournier gangrene; Infant; Early diagnosis; Sepsis; Liver dysfunction; Case report

©The Author(s) 2023. Published by Baishideng Publishing Group Inc. All rights reserved.

Core Tip: This is a case report of an infant with Fournier gangrene. Despite its rarity, Fournier gangrene remains a disease with severe complications and high mortality. The prognosis is influenced by the timing of medical treatment. Despite receiving optimal antibiotic and supportive care after hospital admission, the patient developed sepsis with severe liver damage, making treatment more challenging. The prognosis to sepsis and liver dysfunction was influenced by delayed diagnosis. Through this case report we highlight the importance of early recognition and high clinical suspicion by all doctors (pediatricians, dermatologist, surgeons, *etc.*), to make an accurate diagnosis.

Citation: Bakalli I, Heta S, Kola E, Celaj E. Fournier gangrene in an infant, complicated with severe sepsis and liver dysfunction: A case report. *World J Clin Cases* 2023; 11(30): 7398-7402

URL: <https://www.wjgnet.com/2307-8960/full/v11/i30/7398.htm>

DOI: <https://dx.doi.org/10.12998/wjcc.v11.i30.7398>

INTRODUCTION

Fournier gangrene is a severe and aggressive form of infective necrotizing fasciitis that affects the perineum and genitalia as a result of polymicrobial infection[1-4]. Despite being named Fournier (Jean-Alfred Fournier) after he published a series of five cases, the condition was first described as an idiopathic, rapidly progressive necrotic process of the soft tissues leading to genital organ gangrene in 1764 by a physician named Baurienne[5]. Fournier gangrene in children is rare, with an annual incidence of 0.8 per million patients, and little is known about the disease in this age group[2,4-10]. A review of the literature revealed only 56 pediatric cases, with 66% of those involving infants under the age of 3 mo[5,7]. Despite its rarity, Fournier gangrene was and continues to be a disease with severe complications and high mortality. The prognosis for this disease is dependent on the timing of medical care. Because early manifestations are frequently subtle, through this case report we would like to emphasize the importance of early recognition and high clinical suspicion by all doctors in order to make an accurate diagnosis.

CASE PRESENTATION

Chief complaints

A 3-month-old boy was admitted to our pediatric intensive care unit in critical condition, intoxicated and drowsy, after a 5-day history of fever and progressively increasing scrotal swelling with scrotal zone discoloration.

History of present illness

He initially displayed skin breaches in the scrotum and right inguinal area (Figure 1A). The child was examined 3 days before hospitalization and began antibiotic treatment (oral cefaclor and topical fusidic acid). Despite outpatient treatment, the child's clinical condition worsened (Figure 1B).

History of past illness

His medical history revealed that he was born at term, appropriate for gestational age, and weighed 3500 g. The child was in good general condition 5 d before admission. There was no history of surgical intervention, perineum or lower abdomen injury, catheterization, insect bite, or other predisposing conditions.

Personal and family history

His parents are in good health.

Physical examination

The scrotal area appeared edematous at admission, with a dark to black color that extended towards the dexter inguinal area and the abdomen, with some ulcerative areas (Figure 1C).

Laboratory examinations

The initial laboratory results revealed leukocytosis (white blood cell count of 23200/mm³) with 78% neutrophils, high C-reactive protein (9.95 mg/dL), blood urea nitrogen 80 mg/dL; creatinine 0.72 mg/dL, high aspartate transaminase (AST) levels of 563 IU/L, high alanine transaminase (ALT) levels of 1871 U/L, total bilirubin of 1.39 mg/dL, low total protein of 3.5 g/dL, prothrombin time of 21 s, activated partial thromboplastin time of 47.7 s, and fibrinogen level of 302 mg/dL.

Imaging examinations

Ultrasound was performed to rule out a deep soft tissue infection and a torsion testis. The presence of liquid with corpuscular content was confirmed by ultrasound, as was thickening of the scrotal walls, but without a clear assessment of blood flow in the appropriate testis.



DOI: 10.12998/wjcc.v11.i30.7398 Copyright ©The Author(s) 2023.

Figure 1 Evolution of scrotal necrosis from the day 2 of history to day 1 of hospital admission. A: Skin breaches in the scrotum and right inguinal area; B: The child's clinical condition worsened; C: The scrotal area appeared edematous at admission, with a dark to black color that extended towards the dexter inguinal area and the abdomen, with some ulcerative areas.

MULTIDISCIPLINARY EXPERT CONSULTATION

After vigorous resuscitation, the surgeon decided on an immediate intervention to rule out testicular torsion and to debride the affected region.

FINAL DIAGNOSIS

The intervention clarified the child's diagnosis of Fournier's gangrene.

TREATMENT

In addition to other supportive measures, the baby was given intravenous fluids and broad-spectrum antibiotics that covered both aerobic and anaerobic bacteria.

OUTCOME AND FOLLOW-UP

Despite receiving optimal antibiotic and supportive therapy, the patient developed sepsis with severe liver injury that led to liver failure, making treatment more challenging. AST, ALT and bilirubin levels peaked on the second day of admission (ALT: 651 IU/L, AST: 2993 IU/L, total bilirubin: 3.9 mg/dL, prothrombin time: 6 s). The liver parameters improved 4 d after admission and returned to normal levels after 10 d. The results of blood and urine cultures were both negative. The immunological parameters were within the normal range. In the postoperative period, antibiotics and other supportive treatments, as well as regular dressing, were continued. The child was discharged from the hospital in a good clinical condition.

DISCUSSION

Fournier gangrene is a type of necrotizing fasciitis that affects the external genital organs as well as the perianal region. It is characterized by thrombosis of the feeding arteries, which leads to gangrene of the skin and subcutaneous tissues, as well as severe intoxication and multiple organ failure. The necrotizing process is frequently caused by an infection in the anorectum, urogenital tract, or genital skin[1,17,18]. Omphalitis, strangulated hernia, prematurity, diaper rash, varicella infection, circumcision, and perineal abscesses have all been reported as etiological factors in the pediatric age group[1,8]. Trauma, minor skin breaches (as in our case), insect bites, surgeries or invasive procedures in the perineal region, urethral instrumentation, burns, and systemic infections are some of the other causes in children[1,5].

Despite its rarity, the unfavorable prognosis is dependent on the timing of medical care. The treatment delay is accompanied by a high lethality rate, reaching 90%, due to the development of septic shock and its associated complications[13]. Fournier gangrene progresses more quickly in children than in older people as their immune systems are immature and their skin barrier is more fragile. As a result, the situation is more likely to lead to septic shock, dysfunction of many organs, and a poor prognosis[12].

The liver plays an important role in defensive responses to scavenge bacteria and produce inflammatory mediators during sepsis. However, the liver can also be a target of a dysregulated inflammatory response. Although liver dysfunction is not the most common type of organ injury seen in septic patients, when it progresses to liver failure, it

becomes a serious complication. Based on the new consensus for sepsis definition, the clinical situation, and the additional investigation, sepsis appears to have been complicated by significant liver injury in our case[14-16].

Lu *et al*[12] also reported the case of an 8-month-old girl with Fournier gangrene caused by *Escherichia coli*, which was complicated by septic shock and septic associated encephalopathy[12]. The majority of pediatric cases of Fournier gangrene have been reported in previously healthy children, who may appear well in the early stages of the disease and thus escape diagnosis, as in our case[3].

Radiography, ultrasound, computed tomography, and magnetic resonance imaging (MRI) are all used to diagnose Fournier gangrene. An X-ray of the region may show gas in the soft tissues, which is an absolute indication of surgical intervention. However, the diagnosis is usually clinical[5].

In soft tissues, ultrasound can detect fluid or gas. Gas in the scrotal wall is the sonographic hallmark of Fournier gangrene. Scrotal wall edema is possible. Blood flow and testicular status can also be determined using ultrasound. Contrary to most urological diseases, Fournier gangrene does not affect the testes. This is due to their supply's independence from the blood supply of the scrotum and penis[13]. Even in our case, we were able to keep the tests. The use of MRI in the treatment of Fournier gangrene is not well documented in the literature. MRI should not be used to postpone surgery if the diagnosis is highly suspect.

The treatment for Fournier gangrene is based on an emergency surgical intervention combined with antibacterial therapy. Surgery is required for a definitive diagnosis and the removal of necrotic tissue. A delay to the surgical intervention is the most significant modifiable risk factor associated with mortality. Once Fournier gangrene diagnosis has been made, all necrotic tissues must be removed. A systemic pediatric review determined that early aggressive surgical treatment was the best option[17]. A study confirmed the opinion that early surgical intervention reduces mortality in a large retrospective review of 379 patients. Those who received earlier intervention had a lower fatality rate (odds ratio = 0.38) than those who received intervention 3 d or later[5].

In approximately 95% of cases, the cause of Fournier gangrene can be identified. Necrotizing fasciitis is classified into two types based on the organisms that cause it. Type 1 bacteria are typically polymicrobial, containing both aerobic and anaerobic bacteria. Type 2 is monomicrobial and is frequently caused by *Streptococcus* group A alone or in combination with *Staphylococcus aureus*[18]. Zundel *et al*[17] with their systematic review, aimed to better characterize the pediatric Fournier gangrene. In their study, 53 pediatric patients were included. Monomicrobial necrotizing fasciitis (type 2) was far more common than polymicrobial fasciitis (type 1). *Pseudomonas aeruginosa* was frequently isolated alongside streptococci and staphylococci[17]. In our case, we were unable to determine the etiology.

Considering the polymicrobial (aerobic and anaerobic) causative microflora, the drugs of choice for antibacterial therapy of Fournier gangrene are second or third generation cephalosporins combined with antibiotics from the nitroimidazole, fluoroquinolone and aminoglycoside groups. Antibiotics from the carbapenem class are used in the complex processes of antibacterial therapy in severe cases of the disease[13]. Streptococci and staphylococci are more common in children than Gram-negative and anaerobic bacteria, which predominate in adults[1,17,19]. As a result, multiple antibiotics are recommended, including one specifically targeted toward *S. aureus*[12,20]. This aggressive treatment modality was effective in our patient.

According to various authors, mortality ranged from 3% to 45% due to severe sepsis, coagulopathy and renal failure[1,17,20,21]. Necrotizing fasciitis in neonates is associated with a nearly 50% mortality rate[10]. The prognosis of this disease is influenced by age, the presence of infectious agents, extent of the disease, and underlying medical conditions. A multicenter retrospective study found that older age, diabetes, anemia, sepsis, a delay in initial treatment, and a Fournier Gangrene Severity Index score of 9 are important predictors of disease severity[12,22]. The mainstays of management in children with Fournier gangrene are a high index of suspicion and prompt diagnosis, as well as surgical debridement and a multidisciplinary approach[12,20,21].

CONCLUSION

Recognizing Fournier gangrene, referring the children for a pediatric surgery as soon as possible, and providing the appropriate antibiotic coverage are crucial for avoiding sepsis and multiorgan dysfunction and improving prognosis.

ACKNOWLEDGMENTS

We thank all the doctors and nurses who have contributed to this child's treatment.

FOOTNOTES

Author contributions: Bakalli I conceptualized, supervised data collection, drafted the initial manuscript, and corrected the final English version; Heta S and Celaj E contributed to the acquisition of data; Kola E revised the manuscript; and all authors have read and approved the manuscript critically.

Informed consent statement: The present manuscript used anonymous images to produce its analyzes and results, in a method that obeys the norms of medical bioethics. Thus, there was no direct or even indirect contact between researchers and patients, with no

necessity for “Signed Informed Consent Form” to carry out our study.

Conflict-of-interest statement: All the authors report no relevant conflicts of interest for this article.

CARE Checklist (2016) statement: The authors have read the CARE Checklist (2016), and the manuscript was prepared and revised according to the CARE Checklist (2016).

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <https://creativecommons.org/licenses/by-nc/4.0/>

Country/Territory of origin: Albania

ORCID number: Ilirjana Bakalli 0000-0002-7063-3765.

S-Editor: Wang JJ

L-Editor: Kerr C

P-Editor: Wang JJ

REFERENCES

- Bains SP, Singh V, Gill MK, Jain A, Arry V. Fournier's Gangrene in a Two Year Old Child: A Case Report. *J Clin Diagn Res* 2014; **8**: ND01-ND02 [PMID: 25302233 DOI: 10.7860/JCDR/2014/9257.4662]
- Neogi S, Kariholu PL, Chatterjee D, Singh BK, Kumar R. Fournier's gangrene in a pediatric patient after prolonged neglected diarrhea: A case report. *IJCR* 2013; **4**: 135-137 [DOI: 10.5348/ijeri-2013-02-279-CR-13]
- Tsinti M, Tsekoura T, Blevrakis E, Vlachakis I, Tsilimigaki-Christaki A. Fournier's Gangrene Associated with Local Cutaneous HPV Lesions in a Previously Healthy Girl. *Case Rep Pediatr* 2013; **2013**: 704532 [PMID: 23710402 DOI: 10.1155/2013/704532]
- Jefferies M, Saw NK, Jones P. Fournier's gangrene in a five year old boy - beware of the child post varicella infection. *Ann R Coll Surg Engl* 2010; **92**: W62-W63 [PMID: 20626965 DOI: 10.1308/147870810X12699662980790]
- Pais MV. Fournier gangrene. [cited 15 May 2023]. Available from: <https://emedicine.medscape.com/article/2028899-overview>
- Ameh EA, Dauda MM, Sabiu L, Mshelbwala PM, Mbibu HN, Nmadu PT. Fournier's gangrene in neonates and infants. *Eur J Pediatr Surg* 2004; **14**: 418-421 [PMID: 15630645 DOI: 10.1055/s-2004-821138]
- Irom KS. Fournier's gangrene in an infant: Is primary surgical repair a better alternative for management? A case report. *Ped Urol Case Rep* 2015; **2**: 1 [DOI: 10.14534/PUCR.2015614083]
- Ekingen G, Isken T, Agir H, Oncel S, Günlemez A. Fournier's gangrene in childhood: a report of 3 infant patients. *J Pediatr Surg* 2008; **43**: e39-e42 [PMID: 19040919 DOI: 10.1016/j.jpedsurg.2008.09.014]
- Dey S, Bhutia KL, Baruah AK, Kharga B, Mohanta PK, Singh VK. Neonatal Fournier's gangrene. *Arch Iran Med* 2010; **13**: 360-362 [PMID: 20597572]
- Zgraj O, Paran S, O'Sullivan M, Quinn F. Neonatal scrotal wall necrotizing fasciitis (Fournier gangrene): a case report. *J Med Case Rep* 2011; **5**: 576 [PMID: 22151925 DOI: 10.1186/1752-1947-5-576]
- Hota PK. Fournier's Gangrene: Report of 2 Cases. *Case Rep Emerg Med* 2012; **2012**: 984195 [PMID: 23326734 DOI: 10.1155/2012/984195]
- Lu G, Liu H, Li D, Qiao L, Liu Z, Ma Z. Fournier gangrene caused by Escherichia coli complicated with septic shock and sepsis-associated encephalopathy in an 8-month-old girl. *Minerva Med* 2023; **114**: 133-135 [PMID: 33047940 DOI: 10.23736/S0026-4806.20.07034-2]
- Chernyadyev SA, Ufimtseva MA, Vishnevskaya IF, Bochkarev YM, Ushakov AA, Beresneva TA, Galimzyanov FV, Khodakov VV. Fournier's Gangrene: Literature Review and Clinical Cases. *Urol Int* 2018; **101**: 91-97 [PMID: 29949811 DOI: 10.1159/000490108]
- Wang D, Yin Y, Yao Y. Advances in sepsis-associated liver dysfunction. *Burns Trauma* 2014; **2**: 97-105 [PMID: 27602369 DOI: 10.4103/2321-3868.132689]
- Singer M, Deutschman CS, Seymour CW, Shankar-Hari M, Annane D, Bauer M, Bellomo R, Bernard GR, Chiche JD, Coopersmith CM, Hotchkiss RS, Levy MM, Marshall JC, Martin GS, Opal SM, Rubenfeld GD, van der Poll T, Vincent JL, Angus DC. The Third International Consensus Definitions for Sepsis and Septic Shock (Sepsis-3). *JAMA* 2016; **315**: 801-810 [PMID: 26903338 DOI: 10.1001/jama.2016.0287]
- Bakalli I. Liver Dysfunction in Severe Sepsis from Respiratory Syncytial Virus. *J Pediatr Intensive Care* 2018; **7**: 110-114 [PMID: 31073482 DOI: 10.1055/s-0037-1612609]
- Zundel S, Lemaréchal A, Kaiser P, Szavay P. Diagnosis and Treatment of Pediatric Necrotizing Fasciitis: A Systematic Review of the Literature. *Eur J Pediatr Surg* 2017; **27**: 127-137 [PMID: 27380058 DOI: 10.1055/s-0036-1584531]
- Mosayebi Z, Omidian A, Movahedian AH, Kompani F, Hosseiniodeh SS. Fournier's Gangrene in a Neonate With Acute Myeloid Leukemia: A Case Report. *Iran J Pediatr* 2016; **26**: e4537 [PMID: 27617071 DOI: 10.5812/ijp.4537]
- Idweini Y. Necrotizing Scrotum Fasciitis (Fournier's Gangrene) in an Infant: A Case Report and Review of the Literature. *Clin Res Urol* 2018; **1** [DOI: 10.33309/2638-7670.010101]
- Singh AP, Gupta A, Pardeshi R, Garg D. Neonatal Fournier's gangrene. *J Clin Neonatol* 2018 [DOI: 10.4103/jcn.jcn_3_18]
- Rouzkrokh M, Tavassoli A, Mirshemirani A. Fournier's Gangrene in Children: Report on 7 Cases and Review of Literature. *Iran J Pediatr* 2014; **24**: 660-661 [PMID: 25793080]
- Verma S, Sayana A, Kala S, Rai S. Evaluation of the Utility of the Fournier's Gangrene Severity Index in the Management of Fournier's Gangrene in North India: A Multicentre Retrospective Study. *J Cutan Aesthet Surg* 2012; **5**: 273-276 [PMID: 23378710 DOI: 10.4103/0974-2077.104916]



Published by **Baishideng Publishing Group Inc**
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA

Telephone: +1-925-3991568

E-mail: bpgoffice@wjgnet.com

Help Desk: <https://www.f6publishing.com/helpdesk>

<https://www.wjgnet.com>

