



Circulating copeptin level and the clinical prognosis of patients with chronic liver disease

Hao-Qian Tan, Ming Zhao, Zan Huang, Yang Liu, Han Li, Long-Hui Ma, Jun-Ying Liu

Specialty type: Gastroenterology and hepatology

Provenance and peer review: Unsolicited article; Externally peer reviewed.

Peer-review model: Single blind

Peer-review report's scientific quality classification

Grade A (Excellent): A
Grade B (Very good): B
Grade C (Good): 0
Grade D (Fair): 0
Grade E (Poor): 0

P-Reviewer: El-Shabrawi MH, Egypt; Rodrigues AT, Brazil

Received: June 15, 2023

Peer-review started: June 15, 2023

First decision: July 14, 2023

Revised: July 21, 2023

Accepted: July 28, 2023

Article in press: July 28, 2023

Published online: August 21, 2023



Hao-Qian Tan, Department of Gastroenterology, Zhoukou Central Hospital Affiliated to Xinxiang Medical University, Zhoukou 466000, Henan Province, China

Ming Zhao, Yang Liu, Han Li, Jun-Ying Liu, Department of Gastroenterology, Zhoukou Central Hospital, Zhoukou 466000, Henan Province, China

Zan Huang, Long-Hui Ma, Department of Teaching and Research, Zhoukou Central Hospital, Zhoukou 466000, Henan Province, China

Corresponding author: Jun-Ying Liu, MD, Director, Department of Gastroenterology, Zhoukou Central Hospital, No. 26 Renmin Road East Section, Chuanhui District, Zhoukou 466000, Henan Province, China. liujunying781025@163.com

Abstract

BACKGROUND

The relationship between copeptin and the severity of circulatory dysfunction and systemic stress response in patients with chronic liver disease (CLD) has been established. Nevertheless, the potential of serum copeptin levels to predict the prognosis of CLD patients remains unclear.

AIM

To conduct a systematic review and meta-analysis to investigate the correlation between serum copeptin and transplant-free survival (TFS) in this population.

METHODS

To achieve the objective of the meta-analysis, PubMed, Embase, the Cochrane Library, and the Web of Science were searched to identify observational studies with longitudinal follow-up. The Cochrane Q test was utilized to assess between-study heterogeneity, and the I^2 statistic was estimated. Random-effects models were employed to combine the outcomes, taking into account the potential influence of heterogeneity.

RESULTS

Ten datasets including 3133 patients were involved. The follow-up durations were 1 to 48 mo (mean: 12.5 mo). Overall, it was shown that a high level of serum copeptin was associated with a poor TFS [risk ratio (RR): 1.82, 95% confidence interval: 1.52-2.19, $P < 0.001$; $P = 0\%$]. In addition, sensitivity analysis by omitting one dataset at a time showed consistent results (RR: 1.73-2.00, $P < 0.05$). Finally, subgroup analyses according to study country, study design, patient diagnosis,

cutoff of copeptin, follow-up duration, and study quality score also showed similar results (P for subgroup difference all > 0.05).

CONCLUSION

Patients with CLD who have high serum copeptin concentrations may be associated with a poor clinical prognosis.

Key Words: Chronic liver disease; Liver cirrhosis; Copeptin; Prognosis; Mortality; Meta-analysis

©The Author(s) 2023. Published by Baishideng Publishing Group Inc. All rights reserved.

Core Tip: Serum copeptin has been related to the severity of circulatory dysfunction and systemic stress response in patients with chronic liver disease (CLD). However, little is known about the relationship between serum copeptin and the prognosis of patients with CLD. In this systematic review and meta-analysis, evidence from ten datasets including 3133 patients were integrated. The results showed that a high level of serum copeptin was associated with a poor transplant-free survival in these patients. These findings support the use of serum copeptin as a prognostic biomarker for patients with CLD.

Citation: Tan HQ, Zhao M, Huang Z, Liu Y, Li H, Ma LH, Liu JY. Circulating copeptin level and the clinical prognosis of patients with chronic liver disease. *World J Gastroenterol* 2023; 29(31): 4797-4808

URL: <https://www.wjgnet.com/1007-9327/full/v29/i31/4797.htm>

DOI: <https://dx.doi.org/10.3748/wjg.v29.i31.4797>

INTRODUCTION

Chronic liver disease (CLD) patients are more likely to develop complications associated with the progression of the disease, such as portal hypertension, ascites, spontaneous bacterial peritonitis, gastroesophageal varices, and hepatic encephalopathy[1,2]. For those with advanced CLD (ACLD), the clinical prognosis is generally poor, which has become a substantial cause of morbidity and mortality worldwide[3,4]. Accumulating evidence suggests that patients with ACLD may have multiple features of circulatory dysfunction, such as incremental intrahepatic vascular resistance, decrease of portal blood flow, and a reduced systemic vascular resistance[5,6], all of which could subsequently activate the neuro-hormonal system for counter regulation, including the upregulated systemic arginine vasopressin (AVP)[7,8]. As a biomarker of disease progression, the AVP increment in patients with CLD has been hypothesized as a predictor of poor prognosis[9]. However, as a small peptide of nine amino acids, the blood level of AVP is difficult to quantitatively analyze due to its short half-life[10]. Copeptin is a 39-amino-acid glycopeptide which contains the C-terminus of AVP precursor, making it an effective surrogate marker for AVP release[11,12]. As a result of osmotic stress and osmotic stimulation, a high level of copeptin has been demonstrated in patients with ACLD[13,14]. However, it remains unknown whether the serum level of copeptin may predict the prognosis of patients with CLD. Therefore, in this study, we conducted a systematic review and meta-analysis to investigate the potential association between serum copeptin and clinical outcomes in this patient population.

MATERIALS AND METHODS

Throughout the process of planning, conducting, and reporting the study, the Preferred Reporting Items for Systematic Reviews and Meta-Analyses statement[15,16] and Cochrane Handbook[17] were followed.

Search of databases

We searched electronic databases including PubMed, Embase, the Cochrane Library, and Web of Science, starting from inception and ending April 21st, 2023, for studies that had been published up to that date. The search was performed with terms related to our study including: (1) "Copeptin" OR "C-terminal provasopressin"; and (2) "cirrhosis" OR "cirrhotic" OR "liver" OR "hepatic" OR "hepatitis". There was no limitation on the language of the publication in the search for human studies. As part of our manual screening process, references from relevant original and review articles were screened for possible relevant studies.

Inclusion and exclusion criteria of studies

Inclusion criteria were developed in accordance with PICOS recommendations and according to the aim of the meta-analysis.

P (patients): Adult patients (18 years or older) with a confirmed diagnosis of CLD; I (exposure): Patients with a high serum concentration of copeptin at baseline. Methods for measuring serum copeptin and cutoffs for defining high serum copeptin were consistent with those of the original studies; C (control): Patients with a low serum concentration of

copeptin at baseline; O (outcomes): Incidence of transplant-free survival (TFS) compared between CLD patients with high vs low serum levels of copeptin at baseline. S (study design): Studies with longitudinal follow-up, including cohort and case-control studies, as well as post-hoc analyses of clinical trials.

Excluded from the meta-analysis were reviews, editorials, preclinical studies, and studies that did not involve patients with CLD, failed to measure serum copeptin, or did not report the relevant outcome. In instances where there was a patient population overlap, the study with the greatest sample size was incorporated into the meta-analysis.

Data extraction and quality evaluation

Two of the authors conducted literature searches, data collection, and assessments of study quality independently. In instances where discrepancies arose, a third author was consulted for discussion, and a consensus was reached. The analysis of studies included the collection of information pertaining to study information, design characteristics, patient diagnosis, demographic factors, measuring methods, serum copeptin cutoffs, follow-up durations, and adjusted variables for the evaluation of the association between serum copeptin levels and TFS in patients with CLD. In terms of quality, the study was scored using the Newcastle-Ottawa Scale[18] based on the criteria for participant selection, the comparability of the groups, and the validity of the outcomes. There were nine stars on the scale, with a larger number of stars representing a higher quality study.

Statistics

Risk ratios (RRs) and corresponding 95% confidence interval (CI) were used as the variables to indicate the association between serum concentration of copeptin and the survival of patients with CLD. A logarithmical transformation was performed on the RR and its corresponding standard error from each study to stabilize and normalize its variance[19]. In order to estimate between-study heterogeneity, the Cochrane Q test and the I^2 statistic[20] were used. An $I^2 > 50\%$ indicates that there is significant heterogeneity between studies. The utilization of a random-effects model was employed to amalgamate the findings, as it has been acknowledged to encompass the impact of potential heterogeneity[17]. In order to assess the impact of individual studies on the meta-analysis outcomes, sensitivity analyses were conducted by eliminating one dataset at a time[21]. To ascertain the effect of study characteristics on the outcome, subgroup analyses were executed based on the study country, design, patient diagnosis, cutoffs of copeptin, follow-up duration, and study quality scores. The subgroups were defined based on the medians of continuous variables. A funnel plot was used to estimate publication bias based on visual judgments of symmetry, along with Egger's regression asymmetry test[22]. The statistical analyses were carried out with RevMan (Version 5.1; Cochrane Collaboration, Oxford, United Kingdom) and Stata software (version 12.0; Stata Corporation, College Station, TX, United States).

RESULTS

Database search and study retrieval

Figure 1 shows the process of the literature search and study retrieval. Initially, 253 records were obtained from the database, and 77 duplicate entries were removed. After screening the titles and abstracts, a further 159 studies were removed as they did not fit the meta-analysis' objectives. Following full-text reviews of 17 studies, eight were excluded because of the reasons listed in Figure 1. Accordingly, 9 studies were obtained for subsequent meta-analysis[23-31].

Study characteristics

One of the studies reported two datasets from two cohort studies[26], and these datasets were independently incorporated into the meta-analysis. Overall, ten datasets from nine cohort studies[23-31], which included 3133 patients with CLD, were used for the meta-analysis. The characteristics of the included studies are summarized in Table 1. These studies were published between 2013 and 2021, and performed in European and Asian countries. As for the study design, four cohorts were prospective[23,25,26,29], and six were retrospective[24,26-28,30,31]. Patients with cirrhosis were included in six cohorts of the included studies[23-27], while patients with various CLD were included in the other four cohorts[28-31]. The mean ages of the patients were 49.0 years to 67.3 years, and the proportions of men were 58.2% to 83.4%. Serum copeptin was measured with the Kryptor immunoassay in eight cohorts[23-28,30], while other methods such as enzyme-linked immunosorbent assay and the automated copeptin immunofluorescent assay were used in the other two studies[29,31]. A high serum level of copeptin was defined according to receiver operating characteristic analysis in four cohorts[24,25,27,31], medians in three cohorts[26,29], tertiles in two cohorts[23,28], and the upper limit of normal level in one cohort[30]. The follow-up durations were 1 mo to 48 mo (mean: 12.5 mo). Multivariate regression analyses were applied in all of the included studies when the association between serum copeptin and TFS of patients with CLD were estimated, and factors including age, sex, and scores for hepatic dysfunction were adjusted, such as the Child-Pugh score, the Model for End-Stage Liver Disease score, and the albumin-bilirubin score. Among the included studies, all had quality scores between seven and nine stars, indicating that they were of good quality (Table 2).

Meta-analysis results

Overall, pooled results of ten datasets from nine cohort studies showed that a high level of serum copeptin at baseline was associated with a poor TFS (RR: 1.82, 95%CI: 1.52-2.19, $P < 0.001$; Figure 2) with no evidence of significant heterogeneity (P for Cochrane Q test = 0.54, $I^2 = 0\%$). In addition, sensitivity analysis by omitting one dataset at a time showed consistent results (RR: 1.73-2.00, $P < 0.05$). Finally, subgroup analyses showed that the association between serum




Figure 1 Flowchart of database search and study inclusion. ACLD: Advanced chronic liver diseases.





Figure 2 Forest plots for the overall meta-analyses regarding the association between serum copeptin and transplant-free survival of patients with chronic liver diseases. 95%CI: 95% confidence interval.

copeptin and poor TFS was consistent between European and Asian studies (P for subgroup difference = 0.16, Figure 3A), between prospective and retrospective cohorts (P for subgroup difference = 0.48, Figure 3B), between studies of patients with cirrhosis and all CLD (P for subgroup difference = 0.42, Figure 4A), between studies with cutoffs for copeptin < and ≥ 15 pmol/L (P for subgroup difference = 0.17, Figure 4B), between studies with mean follow-up durations < and ≥ 1 year (P for subgroup difference = 0.29, Figure 5A), and between studies with different quality scores (P for subgroup difference = 0.14, Figure 5B).


Publication bias

The funnel plots for the meta-analysis of the association between copeptin and TFS in patients with CLD are shown in Figure 6. Based on visual examination, the plots are symmetrical, suggesting that publication bias is low. Additionally, Egger's regression tests indicated a low likelihood of publication bias ($P = 0.47$).

A



B



DOI: 10.3748/wjg.v29.i31.4797 Copyright ©The Author(s) 2023.

Figure 3 Forest plots for the subgroup analyses regarding the association between serum copeptin and transplant-free survival of patients with chronic liver diseases. A: subgroup analysis according to study country; B: Subgroup analysis according to study design. 95%CI: 95% confidence interval.

DISCUSSION

This systematic review and meta-analysis synthesized data from ten cohort studies to investigate the association between serum copeptin concentration and poor TFS in patients with CLD. Our findings indicate that patients with a high serum copeptin level at baseline are at a greater risk of experiencing poor TFS during follow-up, compared to those with a low serum copeptin concentration. The robustness of our results was confirmed through sensitivity analyses that excluded

Table 1 Characteristics of the included studies

Ref.	Country	Design	Diagnosis	Patient number	Mean age (yr)	Male (%)	CP class C (%)	Methods for copeptin measuring	Copeptin analysis	Median follow-up duration (mo)	Variables adjusted
Moreno <i>et al</i> [23], 2013	France	PC	Cirrhosis (alcohol 84.2%)	125	58	69	33.7	Kryptor immunoassay	13 pmol/L, T3:T1-2	11	Age, sex, CP Class, and CRP
Kerbert <i>et al</i> [24], 2015	The Netherlands	RC	Cirrhosis (alcohol 34.4%, viral 18%)	61	54	75.4	31.1	Kryptor immunoassay	21.9 pmol/L, ROC analysis	12	Age, sex, and MELD score
Sola 2016-original	Spain	PC	Cirrhosis (alcohol 43%, HCV 34%)	265	60	66	NR	Kryptor immunoassay	14 pmol/L, median	3	Age, sex, MELD score, and leukocyte count
Sola 2016-validation	Multiple European countries	RC	Cirrhosis (alcohol 51%, HCV 19%)	120	57	72	NR	Kryptor immunoassay	19 pmol/L, median	3	Age, sex, and MELD score
Kerbert <i>et al</i> [25], 2016	The Netherlands and France	PC	Cirrhosis (alcohol 70.1%, viral 11.4%)	184	55.7	70.7	21.2	Kryptor immunoassay	12.3 pmol/L, ROC analysis	12	Age, sex, CP class, and CRP
Kerbert <i>et al</i> [27], 2017	Multiple European countries	RC	Acute decompensated cirrhosis (alcohol 61%, HCV 31.9%, and HBV 5.7%)	779	58	65.7	NR	Kryptor immunoassay	13.6 pmol/L, ROC analysis	3	Age, sex, WBC, sodium, and MELD score
Schneider <i>et al</i> [28], 2019	Germany	RC	End-stage liver disease (alcohol 63.4%, viral 8.5%)	615	57.2	61.9	NR	Kryptor immunoassay	16.3 pmol/L, T3:T1-2	9.3	Age, sex, and MELD-Na score
Zhao <i>et al</i> [29], 2019	China	PC	HBV related ACLF	151	49	83.4	NR	ELISA	18.7 pmol/L, median	1	Age, sex, and MELD score
Hartl <i>et al</i> [30], 2021	Austria	RC	ACLD (alcohol 36.2%, viral 35.9%)	663	56.6	68.2	16.5	Kryptor immunoassay	11.4 pmol/L, ULN	26.2	Age, sex, MELD score, HVPG, albumin, sodium, and presence of HCC
Shigefuku <i>et al</i> [36], 2021	Japan	RC	CLD (alcohol 45.9%, viral 57.0%), cirrhosis 66.5%	170	67.3	58.2	NR	Automated copeptin immuno-fluorescent assay	4.8 pmol/L, ROC analysis	48	Age, sex, eGFR, ALBI score, and presence of HCC

CP: Child-Pugh; PC: Prospective cohort; RC: Retrospective cohort; HBV: Hepatitis B virus; HCV: Hepatitis C virus; ACLF: Acute-on-chronic liver failure; ACLD: Advanced chronic liver diseases; CLD: Chronic liver diseases; NR: Not Reported; ELISA: Enzyme-linked immunosorbent assay; T: Tertile; ROC: Receiver operating characteristic; ULN: Upper limit of normal; CRP: C-reactive protein; MELD: Model for end-stage liver disease; WBC: White blood cell; HVPG: Hepatic venous pressure gradient; HCC: Hepatocellular carcinoma; ALBI: Albumin-bilirubin; eGFR: Estimated glomerular infiltrating rate.

individual datasets and subgroup analyses based on various study characteristics, including country, design, patient diagnosis, copeptin cutoffs, follow-up duration, and study quality scores. These findings demonstrate that a high serum copeptin level may be a useful index which is associated with poor prognosis for patients with CLD.

To the best of our knowledge, this is the first meta-analysis investigating the potential role of serum copeptin concentration as a prognostic factor of patients with CLD. Several advantages in meta-analysis methodologies deserve to be noticed. For example, we performed a comprehensive literature search in four commonly used databases, which could provide current evidence regarding the relationship of serum copeptin and TFS of patients with CLD. Furthermore, it is noteworthy that all of the studies incorporated in this analysis were cohort studies, implying a potential longitudinal correlation between elevated serum copeptin levels and diminished transplant-free survival among individuals with

Table 2 Quality evaluation of the included studies via the Newcastle-Ottawa scale


Ref.	Representativeness of the exposed cohort	Selection of the non-exposed cohort	Ascertainment of exposure	Outcome not present at baseline	Control for age and sex	Control for other confounding factors	Assessment of outcome	Suitable follow-up duration	Adequacy of follow-up of cohorts	Total
Moreno <i>et al</i> [23], 2013	1	1	1	1	1	1	1	0	1	8
Kerbert <i>et al</i> [24], 2015	0	1	1	1	1	1	1	1	1	8
Sola 2016-original	1	1	1	1	1	1	1	0	1	8
Sola 2016-validation	0	1	1	1	1	1	1	0	1	7
Kerbert <i>et al</i> [25], 2016	1	1	1	1	1	1	1	1	1	9
Kerbert <i>et al</i> [27], 2017	0	1	1	1	1	1	1	0	1	7
Schneider <i>et al</i> [28], 2019	0	1	1	1	1	1	1	0	1	7
Zhao <i>et al</i> [29], 2019	1	1	1	1	1	1	1	0	1	8
Hartl <i>et al</i> [30], 2021	0	1	1	1	1	1	1	1	1	8
Shigefuku <i>et al</i> [36], 2021	1	1	1	1	1	1	1	1	1	9

CLD. Additionally, the utilization of multivariate regression analysis in all of the studies included in this review indicates that the relationship between heightened serum copeptin levels and reduced TFS in this population may be autonomous of potential confounding factors, such as age, gender, and hepatic dysfunction scores. Ultimately, multiple sensitivity and subgroup analyses were executed, and the uniform outcomes reinforced the durability and stability of the conclusions. Collectively, these conclusions substantiate that elevated serum levels of copeptin in individuals with CLD may serve as an indicator of unfavorable prognosis.


There may be multiple mechanisms underlying the relationship between a high serum copeptin level and a poor survival outcome in patients with CLD. Previous studies have shown that a high copeptin level in patients with cirrhosis was correlated to the risk of various complications that may lead to a poor prognosis of these patients, such as gastrointestinal hemorrhage due to portal hypertension, hepatorenal syndrome, hepatic encephalopathy, and larger amounts of ascites[32,33]. Pathophysiologically, as a surrogate marker of AVP, increased copeptin may reflect the enhanced systemic release of AVP in these patients[34]. A high AVP in patients with cirrhosis may deteriorate the status of vasoconstriction, water retention, and hyponatremia, which have all been recognized as key risk factors for poor survival of these patients[35]. The role of copeptin in the pathogenesis and progression of CLD should be further investigated, either as a simple biomarker or an active participant in the disease.

This study is subject to certain limitations. Firstly, the meta-analysis results were predominantly influenced by studies involving patients with ACLD, including those with decompensated cirrhosis and other end-stage liver diseases. The efficacy of copeptin as a prognostic factor in patients with early CLD requires further validation in future research. Additionally, despite the utilization of multivariate regression analyses across all the studies included, the potential for residual factors to confound the association between copeptin and TFS cannot be entirely ruled out. For example, serum copeptin may predict the response to tolvaptan in patients with decompensated cirrhosis[36]. Accordingly, use of treatments such as tolvaptan may confound the association between copeptin and TFS. Furthermore, the determination of an optimal serum copeptin cutoff for predicting the survival of patients with CLD remains elusive, necessitating further investigation. Moreover, the absence of a causal association between elevated copeptin levels and unfavorable TFS in CLD patients is attributable to the reliance on observational studies in the meta-analysis.

A



B




DOI: 10.3748/wjg.v29.i31.4797 Copyright ©The Author(s) 2023.

Figure 4 Forest plots for the subgroup analyses regarding the association between serum copeptin and transplant-free survival of patients with chronic liver diseases. A: Subgroup analysis according to the diagnosis of the patients; B: Subgroup analysis according to the cutoffs of serum copeptin. 95%CI: 95% confidence interval.

CONCLUSION

The findings of the meta-analysis indicate that an elevated serum copeptin concentration in individuals with CLD is linked to unfavorable TFS. The assessment of serum copeptin levels may hold significance in the stratification of risk among CLD patients. Furthermore, it is imperative to investigate whether the reduction of copeptin levels in these patients is correlated with a better clinical outcome, particularly in those with ACLD.



DOI: 10.3748/wjg.v29.i31.4797 Copyright ©The Author(s) 2023.

Figure 5 Forest plots for the subgroup analyses regarding the association between serum copeptin and transplant-free survival of patients with chronic liver diseases. A: Subgroup analysis according to the follow-up durations; B: Subgroup analysis according to the study quality scores. 95%CI: 95% confidence interval.




Figure 6 Funnel plots for the publication bias underlying the meta-analysis regarding the association between serum copeptin and transplant-free survival of patients with chronic liver diseases. RR: Risk ratio.

ARTICLE HIGHLIGHTS

Research background

Patients with chronic liver disease (CLD) will develop various complications with the progression of the disease. Upregulated systemic arginine vasopressin (AVP) has been observed in patients with advanced CLD. However, measuring AVP is clinically challenging due to the short half-life. Copeptin is a C-terminus of AVP precursor, which may be of importance for prognostic prediction in patients with CLD.

Research motivation

Identifying biomarkers that predict the prognosis of patients with CLD is clinically important. Although there are pilot studies aiming to correlate copeptin with survival of patients with CLD, the results are not always consistent. In this regard, a systematic review with meta-analysis is particularly useful.

Research objectives

To investigate the correlation between serum copeptin and transplant-free survival (TFS) in patients with CLD with a systematic review and meta-analysis.

Research methods

Studies were obtained by search of PubMed, Embase, the Cochrane Library, and Web of Science. Two authors independently screened the studies, assessed the study quality with Newcastle-Ottawa Scale, and extracted the data. Risk ratios and corresponding 95% confidence intervals were used as the variables to indicate the association between serum concentration of copeptin and the survival of patients with CLD. The RevMan and Stata software were used for the statistical analyses.

Research results

This meta-analysis enrolled ten datasets involving 3133 patients, who were followed for 1 to 48 mo (mean: 12.5 mo). We found that a high level of serum copeptin was associated with a poor TFS, with a risk ratio of 1.82. Additionally, sensitivity analysis retrieved similar results by omitting one dataset at a time. The robustness of the finding was further evidenced by consistent results of subgroup analyses according to study country, study design, patient diagnosis, cutoff of copeptin, follow-up duration, and study quality score.

Research conclusions

High serum concentration of copeptin may be associated with a poor clinical prognosis in patients with CLD. These findings were not significantly affected by either of the included studies and were not influenced by multiple study characteristics within the subgroup analysis.

Research perspectives

In view of the standard methods for the measuring copeptin in clinical practice, as well as the finding of the meta-analysis, evaluating serum copeptin may be considered at the initial management of patients with CLD, which may provide prognostic significance.

FOOTNOTES

Author contributions: Tan HQ and Liu JY conceived and designed the study; Tan HQ, Zhao M, Huang Z, and Liu Y performed database search, data collection, and study quality evaluation; Tan HQ, Li H, and Ma LH performed statistical analysis; Tan HQ, Zhao M, Huang Z, and Liu JY interpreted the results; Tan HQ wrote the initial draft; all authors revised the manuscript, read and approved the final version of the manuscript.

Conflict-of-interest statement: The authors have no conflicts of interest to declare.

PRISMA 2009 Checklist statement: The authors have read the PRISMA 2009 Checklist, and the manuscript was prepared and revised according to the PRISMA 2009 Checklist.

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <https://creativecommons.org/licenses/by-nc/4.0/>

Country/Territory of origin: China

ORCID number: Hao-Qian Tan 0000-0001-5349-0507; Ming Zhao 0009-0005-8792-0612; Zan Huang 0009-0000-6942-212X; Yang Liu 0009-0003-8587-2800; Han Li 0009-0001-6171-8783; Long-Hui Ma 0009-0003-4569-2346; Jun-Ying Liu 0009-0002-8058-6197.

S-Editor: Chen YL

L-Editor: A

P-Editor: Cai YX

REFERENCES

- 1 Br VK, Sarin SK. Acute-on-chronic liver failure: Terminology, mechanisms and management. *Clin Mol Hepatol* 2023; **29**: 670-689 [PMID: 36938601 DOI: 10.3350/cmh.2022.0103]
- 2 Huang DQ, Terrault NA, Tacke F, Gluud LL, Arrese M, Bugianesi E, Loomba R. Global epidemiology of cirrhosis - aetiology, trends and predictions. *Nat Rev Gastroenterol Hepatol* 2023; **20**: 388-398 [PMID: 36977794 DOI: 10.1038/s41575-023-00759-2]
- 3 Moon AM, Singal AG, Tapper EB. Contemporary Epidemiology of Chronic Liver Disease and Cirrhosis. *Clin Gastroenterol Hepatol* 2020; **18**: 2650-2666 [PMID: 31401364 DOI: 10.1016/j.cgh.2019.07.060]
- 4 Younossi ZM, Wong G, Anstee QM, Henry L. The Global Burden of Liver Disease. *Clin Gastroenterol Hepatol* 2023; **21**: 1978-1991 [PMID: 37121527 DOI: 10.1016/j.cgh.2023.04.015]
- 5 Engelmann C, Clària J, Szabo G, Bosch J, Bernardi M. Pathophysiology of decompensated cirrhosis: Portal hypertension, circulatory dysfunction, inflammation, metabolism and mitochondrial dysfunction. *J Hepatol* 2021; **75** Suppl 1: S49-S66 [PMID: 34039492 DOI: 10.1016/j.jhep.2021.01.002]
- 6 Møller S, Bendtsen F. The pathophysiology of arterial vasodilatation and hyperdynamic circulation in cirrhosis. *Liver Int* 2018; **38**: 570-580 [PMID: 28921803 DOI: 10.1111/liv.13589]
- 7 Tapper EB, Parikh ND. Diagnosis and Management of Cirrhosis and Its Complications: A Review. *JAMA* 2023; **329**: 1589-1602 [PMID: 37159031 DOI: 10.1001/jama.2023.5997]
- 8 Wagener G, Bakker J. Vasopressin in cirrhosis and sepsis: physiology and clinical implications. *Minerva Anestesiol* 2015; **81**: 1377-1383 [PMID: 25384691]
- 9 Rondon-Berrios H, Velez JCQ. Hyponatremia in Cirrhosis. *Clin Liver Dis* 2022; **26**: 149-164 [PMID: 35487602 DOI: 10.1016/j.cld.2022.01.001]
- 10 Leng G, Sabatier N. Measuring Oxytocin and Vasopressin: Bioassays, Immunoassays and Random Numbers. *J Neuroendocrinol* 2016; **28** [PMID: 27467712 DOI: 10.1111/jne.12413]
- 11 Bankir L, Bichet DG, Morgenthaler NG. Vasopressin: physiology, assessment and osmosensation. *J Intern Med* 2017; **282**: 284-297 [PMID: 28649750 DOI: 10.1111/joim.12645]
- 12 Szymgyn H, Szydelko J, Matyjaszek-Matuszek B. Copeptin as a novel biomarker of cardiometabolic syndrome. *Endokrynol Pol* 2021; **72**: 566-571 [PMID: 34378786 DOI: 10.5603/EP.a2021.0072]
- 13 Di Martino V, Weil D, Cervoni JP, Thevenot T. New prognostic markers in liver cirrhosis. *World J Hepatol* 2015; **7**: 1244-1250 [PMID: 26019739 DOI: 10.4254/wjh.v7.i9.1244]
- 14 Mookerjee RP. Prognosis and Biomarkers in Acute-on-Chronic Liver Failure. *Semin Liver Dis* 2016; **36**: 127-132 [PMID: 27172354 DOI: 10.1055/s-0036-1583200]
- 15 Page MJ, Moher D, Bossuyt PM, Boutron I, Hoffmann TC, Mulrow CD, Shamseer L, Tetzlaff JM, Akl EA, Brennan SE, Chou R, Glanville J, Grimshaw JM, Hróbjartsson A, Lalu MM, Li T, Loder EW, Mayo-Wilson E, McDonald S, McGuinness LA, Stewart LA, Thomas J, Tricco AC, Welch VA, Whiting P, McKenzie JE. PRISMA 2020 explanation and elaboration: updated guidance and exemplars for reporting systematic reviews. *BMJ* 2021; **372**: n160 [PMID: 33781993 DOI: 10.1136/bmj.n160]
- 16 Page MJ, McKenzie JE, Bossuyt PM, Boutron I, Hoffmann TC, Mulrow CD, Shamseer L, Tetzlaff JM, Akl EA, Brennan SE, Chou R, Glanville J, Grimshaw JM, Hróbjartsson A, Lalu MM, Li T, Loder EW, Mayo-Wilson E, McDonald S, McGuinness LA, Stewart LA, Thomas J, Tricco AC, Welch VA, Whiting P, Moher D. The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. *BMJ* 2021; **372**: n71 [PMID: 33782057 DOI: 10.1136/bmj.n71]
- 17 Higgins J, Thomas J, Chandler J, Cumpston M, Li T, Page M, Welch V. Cochrane Handbook for Systematic Reviews of Interventions version 6.2. 2021. [cited 5 July 2023]. Available from: <https://training.cochrane.org/handbook>

- 18 Wells GA, Shea B, O'Connell D, Peterson J, Welch V, Losos M, Tugwell P. The Newcastle-Ottawa Scale (NOS) for assessing the quality of nonrandomised studies in meta-analyses. 2010. [cited 5 July 2023]. Available from: [https://www.semanticscholar.org/paper/The-Newcastle-Ottawa-Scale-\(NOS\)-for-Assessing-the-Wells-Wells/c293fb316b6176154c3fdbb8340a107d9c8c82bf](https://www.semanticscholar.org/paper/The-Newcastle-Ottawa-Scale-(NOS)-for-Assessing-the-Wells-Wells/c293fb316b6176154c3fdbb8340a107d9c8c82bf)
- 19 Higgins J, Green S. Cochrane Handbook for Systematic Reviews of Interventions Version 5.1.0. 2011. [cited 5 July 2023]. Available from: <http://handbook-5-1.cochrane.org/>
- 20 Higgins JP, Thompson SG. Quantifying heterogeneity in a meta-analysis. *Stat Med* 2002; **21**: 1539-1558 [PMID: 12111919 DOI: 10.1002/sim.1186]
- 21 Patsopoulos NA, Evangelou E, Ioannidis JP. Sensitivity of between-study heterogeneity in meta-analysis: proposed metrics and empirical evaluation. *Int J Epidemiol* 2008; **37**: 1148-1157 [PMID: 18424475 DOI: 10.1093/ije/dyn065]
- 22 Egger M, Davey Smith G, Schneider M, Minder C. Bias in meta-analysis detected by a simple, graphical test. *BMJ* 1997; **315**: 629-634 [PMID: 9310563]
- 23 Moreno JP, Grandclement E, Monnet E, Clerc B, Agin A, Cervoni JP, Richou C, Vanlemmens C, Dritsas S, Dumoulin G, Di Martino V, Thevenot T. Plasma copeptin, a possible prognostic marker in cirrhosis. *Liver Int* 2013; **33**: 843-851 [PMID: 23560938 DOI: 10.1111/liv.12175]
- 24 Kerbert AJ, Verbeke L, Chiang FW, Laleman W, van der Reijden JJ, van Duijn W, Nevens F, Wolterbeek R, van Hoek B, Verspaget HW, Coenraad MJ. Copeptin as an Indicator of Hemodynamic Derangement and Prognosis in Liver Cirrhosis. *PLoS One* 2015; **10**: e0138264 [PMID: 26378453 DOI: 10.1371/journal.pone.0138264]
- 25 Kerbert AJ, Weil D, Verspaget HW, Moréno JP, van Hoek B, Cervoni JP, Di Martino V, Coenraad MJ, Thevenot T. Copeptin is an independent prognostic factor for transplant-free survival in cirrhosis. *Liver Int* 2016; **36**: 530-537 [PMID: 26502363 DOI: 10.1111/liv.12992]
- 26 Solà E, Kerbert AJ, Verspaget HW, Moreira R, Pose E, Ruiz P, Cela R, Morales-Ruiz M, López E, Graupera I, Solé C, Huelin P, Navarro AA, Ariza X, Jalan R, Fabrellas N, Bente D, de Prada G, Durand F, Jimenez W, van der Reijden JJ, Fernandez J, van Hoek B, Coenraad MJ, Ginès P. Plasma copeptin as biomarker of disease progression and prognosis in cirrhosis. *J Hepatol* 2016; **65**: 914-920 [PMID: 27422752 DOI: 10.1016/j.jhep.2016.07.003]
- 27 Kerbert AJC, Verspaget HW, Navarro AA, Jalan R, Solà E, Bente D, Durand F, Ginès P, van der Reijden JJ, van Hoek B, Coenraad MJ; CANONIC Study Investigators of the EASL-CLIF Consortium. Copeptin in acute decompensation of liver cirrhosis: relationship with acute-on-chronic liver failure and short-term survival. *Crit Care* 2017; **21**: 321 [PMID: 29268760 DOI: 10.1186/s13054-017-1894-8]
- 28 Schneider C, Remmler J, Netto J, Seehofer D, Engelmann C, Berg T, Thiery J, Kaiser T. Copeptin - a biomarker of short-term mortality risk (7 days) in patients with end-stage liver disease. *Clin Chem Lab Med* 2019; **57**: 1897-1905 [PMID: 31199758 DOI: 10.1515/ccm-2019-0023]
- 29 Zhao R, Wu W, Zhou Z, Zheng X, Sun W, Shi Y, Yu H, Wang F, Zhao H, Sun S, Jin L, Sheng J. Prognostic utility of novel biomarkers in acute-on-chronic liver failure (ACLF) associated with hepatitis B: A multicenter prospective study. *Hepatol Res* 2019; **49**: 42-50 [PMID: 30246902 DOI: 10.1111/hepr.13251]
- 30 Hardt L, Jachs M, Desbalmes C, Schaufler D, Simbrunner B, Paternostro R, Schwabl P, Bauer DJM, Semmler G, Scheiner B, Bucsics T, Eigenbauer E, Marculescu R, Szekeres T, Peck-Radosavljevic M, Kastl S, Trauner M, Mandorfer M, Reiberger T. The differential activation of cardiovascular hormones across distinct stages of portal hypertension predicts clinical outcomes. *Hepatol Int* 2021; **15**: 1160-1173 [PMID: 34021479 DOI: 10.1007/s12072-021-10203-9]
- 31 Shigefuku R, Iwasa M, Eguchi A, Tamai Y, Yoshikawa K, Sugimoto R, Takei Y. Serum copeptin level is a biomarker associated with ascites retention and the formation of a portosystemic shunt in chronic liver disease. *J Gastroenterol Hepatol* 2021; **36**: 1006-1014 [PMID: 32790956 DOI: 10.1111/jgh.15215]
- 32 Tawfik AK, Helmy A, Yousef M, Abou-Saif S, Kobtan A, Asaad E, Abd-Elsalam S. Copeptin as a novel marker predicting prognosis of liver cirrhosis and its major complications. *Hepat Med* 2018; **10**: 87-93 [PMID: 30214326 DOI: 10.2147/HMER.S174267]
- 33 Abudeif A, Hashim MS, Ahmed NM, Ahmed AO. Serum copeptin is associated with major complications of liver cirrhosis and spontaneous bacterial peritonitis. *Clin Exp Hepatol* 2023; **9**: 71-78 [PMID: 37064833 DOI: 10.5114/ceh.2023.125970]
- 34 Glavaš M, Gitlin-Domagalska A, Dębowski D, Ptaszyńska N, Łęgowska A, Rolka K. Vasopressin and Its Analogues: From Natural Hormones to Multitasking Peptides. *Int J Mol Sci* 2022; **23** [PMID: 35328489 DOI: 10.3390/ijms23063068]
- 35 Alukal JJ, John S, Thuluvath PJ. Hyponatremia in Cirrhosis: An Update. *Am J Gastroenterol* 2020; **115**: 1775-1785 [PMID: 33156095 DOI: 10.14309/ajg.0000000000000786]
- 36 Shigefuku R, Iwasa M, Eguchi A, Tempaku M, Tamai Y, Suzuki T, Takei Y. Serum Copeptin and Zinc-α2-glycoprotein Levels Are Novel Biomarkers of Tolvaptan Treatment in Decompensated Cirrhotic Patients with Ascites. *Intern Med* 2021; **60**: 3359-3368 [PMID: 34719623 DOI: 10.2169/internalmedicine.7291-21]



Published by **Baishideng Publishing Group Inc**
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA

Telephone: +1-925-3991568

E-mail: bpgoffice@wjgnet.com

Help Desk: <https://www.f6publishing.com/helpdesk>

<https://www.wjgnet.com>

