WJD

# World Journal of Diabetes

Submit a Manuscript: https://www.f6publishing.com

World J Diabetes 2024 March 15; 15(3): 418-428

DOI: 10.4239/wjd.v15.i3.418

**Case Control Study** 

ISSN 1948-9358 (online)

ORIGINAL ARTICLE

# Predictive value of angiopoietin-like protein 8 in metabolic dysfunction-associated fatty liver disease and its progression: A case-control study

Lu-Lu Gan, Can Xia, Xuan Zhu, Yue Gao, Wen-Chang Wu, Qi Li, Ling Li, Zhe Dai, Yi-Min Yan

Lu-Lu Gan, Can Xia, Xuan Zhu, Yue Gao, Wen-Chang Wu, Yi-Min Yan, Medical College, Wuhan Specialty type: Endocrinology and University of Science and Technology, Wuhan 430071, Hubei Province, China metabolism Lu-Lu Gan, Qi Li, Ling Li, Yi-Min Yan, Department of Endocrinology, Xiaogan Hospital Affiliated Provenance and peer review: with Wuhan University of Science and Technology, The Central Hospital of Xiaogan, Xiaogan Unsolicited article; Externally peer 432000, Hubei Province, China reviewed Zhe Dai, Department of Endocrinology, Zhongnan Hospital of Wuhan University, Wuhan Peer-review model: Single blind 430071, Hubei Province, China Peer-review report's scientific Corresponding author: Yi-Min Yan, MD, Deputy Director, Department of Endocrinology, quality classification Xiaogan Hospital Affiliated with Wuhan University of Science and Technology, The Central Grade A (Excellent): 0 Hospital of Xiaogan, No. 6 Plaza Street, Xiaogan 432000, Hubei Province, China. Grade B (Very good): 0 yanyimin180@163.com Grade C (Good): C, C Grade D (Fair): 0 Grade E (Poor): 0 Abstract P-Reviewer: Giacomelli L, Italy; BACKGROUND Horowitz M, Australia The prevalence of metabolic dysfunction-associated fatty liver disease (MAFLD) is rapidly increasing, currently affecting approximately 25% of the global popu-Received: October 28, 2023

lation. Liver fibrosis represents a crucial stage in the development of MAFLD, with advanced liver fibrosis elevating the risks of cirrhosis and hepatocellular carcinoma. Simple serum markers are less effective in diagnosing liver fibrosis First decision: December 23, 2023 compared to more complex markers. However, imaging techniques like transient elastography face limitations in clinical application due to equipment and technical constraints. Consequently, it is imperative to identify a straightforward Article in press: February 18, 2024 yet effective method for assessing MAFLD-associated liver fibrosis.

#### AIM

To investigate the predictive value of angiopoietin-like protein 8 (ANGPTL8) in MAFLD and its progression.

# **METHODS**

We analyzed 160 patients who underwent abdominal ultrasonography in the Endocrinology Department, Xiaogan Central Hospital affiliated to Wuhan University of Science and Technology, during September 2021-July 2022. Using abdominal ultrasonography and MAFLD diagnostic criteria, among the 160

2023

WJD | https://www.wjgnet.com

Peer-review started: October 28,

Revised: January 5, 2024

Accepted: February 18, 2024

Published online: March 15, 2024

patients, 80 patients (50%) were diagnosed with MAFLD. The MAFLD group was divided into the liver fibrosis group (n = 23) and non-liver fibrosis group (n = 57) by using a cut-off fibrosis-4 index  $\ge 1.45$ . Logistical regression was used to analyze the risk of MAFLD and the risk factors for its progression. Receiver operating characteristic curves were used to evaluate the predictive value of serum ANGPTL8 in MAFLD and its progression.

#### RESULTS

Compared with non-MAFLD patients, MAFLD patients had higher serum ANGPTL8 and triglyceride-glucose (TyG) index (both P < 0.05). Serum ANGPTL8 (r = 0.576, P < 0.001) and TyG index (r = 0.473, P < 0.001) were positively correlated with MAFLD. Serum ANGPTL8 was a risk factor for MAFLD [odds ratio (OR): 1.123, 95% confidence interval (CI): 1.066-1.184, P < 0.001). Serum ANGPTL8 and ANGPTL8 + TyG index predicted MAFLD [area under the curve (AUC): 0.832 and 0.886, respectively; both P < 0.05]. Compared with MAFLD patients without fibrosis, those with fibrosis had higher serum ANGPTL8 and TyG index (both P < 0.05), and both parameters were positively correlated with MAFLD-associated fibrosis. Elevated serum ANGPTL8 (OR: 1.093, 95% CI: 1.044-1.144, P < 0.001) and TyG index (OR: 2.383, 95% CI: 1.199-4.736, P < 0.013) were risk factors for MAFLD-associated fibrosis. Serum ANGPTL8 and ANGPTL8 + TyG index predicted MAFLD-associated fibrosis (AUC: 0.812 and 0.835, respectively; both P < 0.05).

#### CONCLUSION

The serum levels of ANGPTL8 are elevated and positively correlated with MAFLD. They can serve as predictors for the risk of MAFLD and liver fibrosis, with the ANGPTL8 + TyG index potentially exhibiting even higher predictive value.

Key Words: Angiopoietin-like protein 8; Metabolic dysfunction-associated fatty liver disease; Fibrosis-4 index; Liver fibrosis

©The Author(s) 2024. Published by Baishideng Publishing Group Inc. All rights reserved.

**Core Tip:** This study unveils elevated serum levels of angiopoietin-like protein 8 (ANGPTL8) in individuals with metabolic dysfunction-associated fatty liver disease (MAFLD). Serum ANGPTL8 emerges not only as a predictive factor for MAFLD risk but also as a powerful indicator for the presence of liver fibrosis in MAFLD. The amalgamation of serum ANGPTL8 with the triglyceride-glucose index demonstrates potential for heightened predictive accuracy in both conditions. Further exploration of serum ANGPTL8 holds promise for enhancing clinical strategies in the prevention and treatment of MAFLD.

**Citation:** Gan LL, Xia C, Zhu X, Gao Y, Wu WC, Li Q, Li L, Dai Z, Yan YM. Predictive value of angiopoietin-like protein 8 in metabolic dysfunction-associated fatty liver disease and its progression: A case-control study. *World J Diabetes* 2024; 15(3): 418-428 **URL:** https://www.wjgnet.com/1948-9358/full/v15/i3/418.htm **DOI:** https://dx.doi.org/10.4239/wjd.v15.i3.418

#### INTRODUCTION

Metabolic dysfunction-associated fatty liver disease (MAFLD), formerly known as non-alcoholic fatty liver disease (NAFLD), is a collective term that encompasses a range of liver diseases from simple steatosis, non-alcoholic steatohepatitis, and liver fibrosis to MAFLD-related cirrhosis and hepatocellular carcinoma[1]. All histological stages of MAFLD are significantly associated with increased overall mortality, and as the disease progresses, the overall mortality increases [2]. The National Health and Nutrition Examination Survey also reported that MAFLD and its associated metabolic disorders are significantly associated with overall and cardiovascular mortality; in addition, the survey found a significant upward trend in the prevalence of MAFLD, which has increased by 14% in the past 3 decades[3]. Nevertheless, there is still a lack of simple and effective methods for the diagnosis of this disease, so it is crucial to find a convenient laboratory index for the early identification of MAFLD and for monitoring its progression.

Angiopoietin-like protein 8 (ANGPTL8) is a 22-kDa protein composed of 198 amino acids; the ANGPTL8 gene is located on chromosome 19p13.2. ANGPTL8 is mainly expressed in the liver, white adipose tissue, and brown adipose tissue, and plays an important role in lipid flux, glucose regulation, and chronic inflammation[4,5]. Many studies have shown that the ANGPTL8 concentration is significantly higher in patients with metabolic diseases such as diabetes[6], obesity[7], and metabolic syndrome[8] than in healthy individuals. Since the pathogenesis of MAFLD is driven by metabolic disorders and changes in glucose-insulin homeostasis, we speculated that ANGPTL8 may be associated with MAFLD. To validate this hypothesis and explore the relationship between MAFLD and ANGPTL8, we evaluated the value of ANGPTL8 in predicting MAFLD and its progression. We hope to provide a simple reference index for the early prediction and prompt management of MAFLD in clinical practice.

Zaishidene® WJD | https://www.wjgnet.com

# MATERIALS AND METHODS

#### Study subjects

We collected the clinical data of 160 patients who underwent abdominal ultrasound examination and were admitted to the Endocrinology Department of Xiaogan Central Hospital, Wuhan University of Science and Technology, between September 2021 and July 2022. The following exclusion criteria were applied: (1) Age < 18 years or > 85 years; (2) patients with mental illness or malignant tumor; (3) patients with autoimmune diseases, active viral hepatitis, drug-induced liver damage, genetic diseases, or chronic schistosomiasis; and (4) patients with acute infection or severe cardiopulmonary, kidney, or cerebrovascular diseases or malnutrition. A total of 160 patients were enrolled according to the above criteria. MAFLD was diagnosed using a combination of abdominal ultrasound findings and relevant diagnostic criteria from the International Expert Consensus on MAFLD[1]: Evidence of fat accumulation in the liver combined with one of the following three criteria, namely, overweight/obesity, presence of type 2 diabetes mellitus, or evidence of metabolic dysregulation (type 2 diabetes was diagnosed using the 1999 World Health Organization diabetes diagnostic criteria). Patients with MAFLD were further divided into those with hepatic fibrosis [defined as fibrosis-4 (FIB-4) index  $\geq$  1.45] and those without hepatic fibrosis (FIB-4 index < 1.45). The study protocol was approved by the ethics committee of Xiaogan Central Hospital, and all study subjects signed informed consent forms.

# Data collection

The age, gender, medical history, height, weight, waist circumference, hip circumference, and blood pressure of all patients were obtained from their medical records. The levels of the following indicators were collected through the Hospital Information Management System: Platelet count (PLT), alanine transaminase (ALT), aspartate transaminase (AST), triglyceride (TG), total cholesterol (TC), high-density lipoprotein-cholesterol (HDL-C), low-density lipoprotein cholesterol (LDL-C), glycated hemoglobin (HbA1c), fasting plasma glucose (FPG), and C-reactive protein (CRP). The homeostasis model assessment of insulin resistance (HOMA-IR) was calculated using the following formula: HOMA-IR =  $[FPG (mmol/L) \times fasting insulin (FINS) (\mu U/mL)]/22.5$ , where FINS stands for fasting insulin level. The homeostasis model assessment of  $\beta$ -cell function (HOMA- $\beta$ ) was calculated using the following formula: HOMA- $\beta$  = [20 × FINS ( $\mu$ U/ mL)]/[FPG (mmol/L) - 3.5]. The TG-glucose (TyG) index was calculated using the formula: TyG = Ln [TG (mg/dL) × FPG (mg/mL)/2]. Liver fibrosis was evaluated using the non-invasive FIB-4 index, which was calculated as follows: FIB-4 = [age (years) × AST (U/L)]/[PLT (× 10<sup>9</sup>/L) × ALT<sup>1/2</sup> (U/L)]. FIB-4 index  $\ge$  1.45 indicated the presence of liver fibrosis[9, 10].

# ANGPTL8 detection

Serum ANGPTL8 concentration was measured using an enzyme-linked immunosorbent assay kit (Wuhan Huamei Biological Engineering Co. Ltd., product number: CSB-EL028107HU, detection range: 6.25-400 pg/mL, sensitivity: 1.56 pg/mL).

#### Statistical analysis

Data were analyzed and processed using SPSS (version 27.0) statistical software. The normality of the data was tested using the Shapiro-Wilk test. Normally distributed data were expressed as mean ± SD and compared using one-way analysis of variance. Non-normally distributed data were expressed as median and interquartile range, and compared using the Wilcoxon rank-sum test. Count data were expressed as percentages and compared using the chi-square test. Spearman correlation analysis was used to analyze the correlation of MAFLD with various indicators as well as the correlation of MAFLD-associated liver fibrosis with ANGPTL8, TyG index, and CRP. Binary logistic regression analysis was used to analyze the risk factors for MAFLD and MAFLD-associated liver fibrosis. Receiver operating characteristic (ROC) curves were used to evaluate the predictive value of ANGPTL8 for the risk of MAFLD and liver fibrosis (FIB-4  $\geq$ 1.45). Differences with P < 0.05 were considered statistically significant.

# RESULTS

# Clinical characteristics of the study subjects

A flow chart of subject selection is presented in Figure 1, and the clinical characteristics of the study subjects are listed in Table 1. A total of 160 patients were enrolled, including 90 men (56.3%) and 70 women (43.7%). The age of the included population ranged from 18 to 85 years. Among the 160 patients, 80 patients (50%) were diagnosed with MAFLD. Among the patients with MAFLD, 23 patients had liver fibrosis, and 57 patients did not have liver fibrosis, based on the FIB-4 index. The following factors did not significantly differ between the MAFLD and non-MAFLD groups: age, gender, systolic blood pressure, diastolic blood pressure, mean arterial pressure, FPG, and HOMA- $\beta$  (P > 0.05). In contrast, body mass index (BMI), waist-to-hip ratio (WHR), HbA1c, FINS, HOMA-IR, TC, TG, TG/HDL-C ratio, LDL-C/HDL-C ratio, CRP, FIB-4 index, TyG index, and ANGPTL8 level were significantly higher in the MAFLD group than in the non-MAFLD group (P < 0.05 for all), while HDL-C was significantly lower in the MAFLD group than in the non-MAFLD group (P = 0.015). The LDL-C was also higher in the MAFLD group than in the non-MAFLD group (2.28 ± 0.99 vs 2.51 ± 1.05 mmol/L), but the difference was not statistically significant (P = 0.144). Finally, the ALT level [14.00 (10.00, 22.75) vs 18.00 (12.00, 25.75) U/L] was significantly higher in the MAFLD group (P = 0.014), while the AST level did not significantly differ between the 2 groups (P = 0.181).



WJD | https://www.wjgnet.com

Table 1 Clinical characteristics of all study subjects							
	Non-MAFLD group ( <i>n</i> = 80)	MAFLD group ( <i>n</i> = 80)	<i>P</i> value <sup>a</sup>				
Sex (female/male)	37/43	33/47	0.524 <sup>1</sup>				
Age (yr)	55.35 ± 12.77	54.70 ± 12.15	0.745 <sup>2</sup>				
WHR	$0.89\pm0.09$	$0.92\pm0.07$	0.039 <sup>2</sup>				
BMI (kg/m <sup>2</sup> )	22.06 (20.16, 24.64)	24.41 (22.89, 27.61)	< 0.001 <sup>3</sup>				
SBP (mmHg)	131.00 (121.75, 147.25)	136.00 (125.00, 150.00)	0.193 <sup>3</sup>				
DBP (mmHg)	83.50 (75.75, 94.00)	87.00 (79.00, 92.00)	0.237 <sup>3</sup>				
MAP (mmHg)	100.00 (91.00, 112.25)	104.00 (95.00, 111.00)	0.151 <sup>3</sup>				
ALT (U/L)	14.00 (10.00, 22.75)	18.00 (12.00, 25.75)	0.014 <sup>3</sup>				
AST (U/L)	18.00 (15.75, 22.00)	19.50 (15.00, 24.00)	0.181 <sup>3</sup>				
TC (mmol/L)	4.29 (3.55, 4.96)	4.39 (3.61, 5.54)	0.222 <sup>3</sup>				
TG (mmol/L)	1.04 (0.62, 1.54)	2.69 (1.39, 4.35)	< 0.001 <sup>3</sup>				
HDL-C (mmol/L)	1.31 (1.04, 1.60)	1.17 (0.85, 1.41)	0.015 <sup>3</sup>				
LDL-C (mmol/L)	$2.28 \pm 0.99$	$2.51 \pm 1.05$	0.144 <sup>2</sup>				
TG/HDL-C	0.84 (0.46, 1.33)	2.73 (1.14, 4.24)	< 0.001 <sup>3</sup>				
LDL-C/HDL-C	1.71 (1.26, 2.29)	2.06 (1.59, 2.92)	0.005 <sup>3</sup>				
HbA1c (%)	6.00 (5.44, 8.70)	7.00 (6.13, 8.58)	0.018 <sup>3</sup>				
FPG (mmol/L)	6.58 (5.63, 8.68)	6.67 (5.93, 9.28)	0.370 <sup>3</sup>				
FINS (µU/mL)	7.01 (4.98, 9.72)	10.00 (5.97, 13.66)	0.004 <sup>3</sup>				
HOMA-β (%)	50.10 (23.25, 87.63)	55.29 (33.96, 94.46)	0.156 <sup>3</sup>				
HOMA-IR	2.08 (1.40, 3.34)	2.99 (1.73, 5.50)	0.008 <sup>3</sup>				
CRP (mg/L)	1.57 (0.98, 2.21)	1.74 (1.19, 2.90)	0.020 <sup>3</sup>				
ANGPTL8 (pg/mL)	23.95 (16.52, 34.20)	44.93 (33.41, 58.01)	< 0.001 <sup>3</sup>				
FIB-4 index	1.18 (0.96, 1.46)	1.39 (1.23, 1.76)	< 0.001 <sup>3</sup>				
TyG index	$8.66 \pm 0.71$	$9.56 \pm 0.92$	< 0.001 <sup>2</sup>				

 $^{a}P < 0.05$  is considered significant.

 $^{2}t$ -test.

<sup>3</sup>Mann Whitney *U* test.

MAFLD: Metabolic dysfunction-associated fatty liver disease; WHR: Waist-to-hip ratio; BMI: Body mass index; SBP: Systolic blood pressure; DBP: Diastolic blood pressure; MAP: Mean arterial pressure; ALT: Alanine transaminase; AST: Aspartate transaminase; TG: Triglyceride; TC: Total cholesterol; HDL-C: High-density lipoprotein-cholesterol; LDL-C: Low-density lipoprotein-cholesterol; HbA1c: Glycated hemoglobin; FPG: Fasting plasma glucose; FINS: Fasting insulin; HOMA-β: Homeostasis model assessment of β-cell function; HOMA-IR: Homeostasis model assessment of insulin resistance; CRP: Creactive protein; ANGPTL8: Angiopoietin-like protein 8; FIB-4 index: Fibrosis-4 index; TyG index: Triglyceride-glucose index.

# Correlation of MAFLD with clinical indicators

Spearman correlation analysis was performed to assess the correlation of the indicators with significant between-group differences with MAFLD. The results are shown in Table 2. Serum ANGPTL8, TyG index, FIB-4 index, BMI, WHR, TG, ALT, FINS, HbA1c, HOMA-IR, CRP, TG/HDL-C, and LDL-C/HDL-C were positively correlated with MAFLD (P < 0.05), while HDL-C was negatively correlated with MAFLD (r = -0.195, P = 0.014). Further, binary logistic regression analysis of the potential risk factors for MAFLD was performed (Table 3). The results showed that ANGPTL8, BMI, and TG had a statistically significant impact on the risk of MAFLD (P < 0.05). Serum ANGPTL8 was a risk factor for the occurrence of MAFLD.

#### Diagnostic performance of ANGPTL8 and TyG index in MAFLD

ROC curves were used to evaluate the diagnostic performance of ANGPTL8, the TyG index, and their combination for the risk of MAFLD (Figure 2). The ROC curve of ANGPTL8 had an area under the curve (AUC) of 0.832 [95% confidence interval (CI): 0.771-0.893, P < 0.001], with an optimal cutoff point of 31.18 pg/mL, sensitivity of 80%, and specificity of 73.08%. The ROC curve of the TyG index had an AUC of 0.773 (95% CI: 0.701-0.845, P < 0.001), with an optimal cutoff



Zaishidena® WJD | https://www.wjgnet.com

 $<sup>^{1}\</sup>chi^{2}$  test.

#### Table 2 Spearman correlation analysis of metabolic dysfunction-associated fatty liver disease and various indicators

	r	<i>P</i> value <sup>a</sup>
ANGPTL8	0.576	< 0.001
TyG index	0.473	< 0.001
FIB-4 index	0.318	< 0.001
BMI	0.394	< 0.001
WHR	0.172	0.031
TG	0.581	< 0.001
HDL-C	-0.195	0.014
ALT	0.195	0.014
FINS	0.231	0.003
HbA1c	0.189	0.017
HOMA-IR	0.211	0.008
CRP	0.189	0.018
TG/HDL-C	0.551	< 0.001
LDL-C/HDL-C	0.226	0.004

#### $^{a}P < 0.05$ is considered significant.

ANGPTL8: Angiopoietin-like protein 8; TyG index: Triglyceride-glucose index; FIB-4 index: Fibrosis-4 index; BMI: Body mass index; WHR: Waist-to-hip ratio; TG: Triglyceride; HDL-C: High-density lipoprotein-cholesterol; ALT: Alanine transaminase; FINS: Fasting insulin; HbA1c: Glycated hemoglobin; HOMA-IR: Homeostasis model assessment of insulin resistance; CRP: C-reactive protein; LDL-C: Low-density lipoprotein-cholesterol.

Table 3 Multifactor logistic regression analysis of factors influencing metabolic dysfunction-associated fatty liver disease						
	В	S.E.	Wals	P value <sup>a</sup>	EXP(B)	95%CI
ANGPTL8	0.116	0.027	19.118	< 0.001	1.123	1.066-1.184
BMI	0.275	0.111	6.142	0.013	1.317	1.059-1.637
ALT	0.032	0.021	2.394	0.122	1.032	0.992-1.075
HDL-C	-0.386	0.804	0.231	0.631	0.680	0.141-3.287
FINS	0.085	0.091	0.875	0.350	1.088	0.911-1.300
HbA1c	-0.248	0.176	1.982	0.159	0.780	0.552-1.102
HOMA-IR	-0.426	0.251	2.869	0.090	0.653	0.399-1.069
CRP	0.160	0.088	3.285	0.070	1.173	0.987-1.395
WHR	-3.793	4.066	0.870	0.351	0.023	0.000-65.098
TG	2.255	0.527	18.277	< 0.001	9.532	3.391-26.800

<sup>a</sup>P < 0.05 is considered significant.

CI: Confidence interval; ANGPTL8: Angiopoietin-like protein 8; BMI: Body mass index; ALT: Alanine transaminase; HDL-C: High-density lipoproteincholesterol; FINS: Fasting insulin; HbA1c: Glycated hemoglobin; HOMA-IR: Homeostasis model assessment of insulin resistance; CRP: C-reactive protein; WHR: Waist-to-hip ratio; TG: Triglyceride.

point of 9.155, sensitivity of 67.50%, and specificity of 80.77%. The combined ROC curve of ANGPTL8 and the TyG index had an AUC of 0.886 (95% CI: 0.846-0.941, P < 0.001), with an optimal cutoff point of 0.546, sensitivity of 75.5%, and specificity of 85.9%.

#### Clinical characteristics of MAFLD patients with liver fibrosis

According to the FIB-4 index, patients with MAFLD were divided into the non-fibrosis group (FIB-4 < 1.45) and the liver fibrosis group (FIB-4  $\ge$  1.45). The serum ANGPTL8 Level (40.75  $\pm$  15.11 vs 58.94  $\pm$  12.81 pg/mL, P < 0.001) and TyG index  $(9.38 \pm 0.85 vs 9.99 \pm 0.96, P = 0.006)$  were significantly higher in the liver fibrosis group than in the non-fibrosis group, while serum CRP level [2.13 (1.31, 4.23) *vs* 1.63 (0.98, 1.96) mg/L, *P* = 0.034] was opposite. However, age, gender, systolic



Raishideng® WJD | https://www.wjgnet.com



Figure 1 Flow chart of subject selection. MAFLD: Metabolic dysfunction-associated fatty liver disease; FIB-4 index: Fibrosis-4 index.



Figure 2 Predictive value of angiopoietin-like protein 8 and triglyceride-glucose index. A: Predictive value in metabolic dysfunction-associated fatty liver disease; B: Predictive value in liver fibrosis. MAFLD: Metabolic dysfunction-associated fatty liver disease; ANGPTL8: Angiopoietin-like protein 8; TyG index: Triglyceride-glucose index.

blood pressure, diastolic blood pressure, mean arterial pressure, FPG, HOMA- $\beta$ , BMI, WHR, HbA1c, FINS, HOMA-IR, TC, TG, HDL-C, LDL-C, TG/HDL-C, LDL-C/HDL-C, ALT, and AST did no significantly differ between the fibrosis and non-fibrosis groups (all *P* > 0.05, Figure 3).

# Correlation of MAFLD-associated liver fibrosis with clinical indicators

Considering the above results, we speculated that there exists a correlation between ANGPTL8 and MAFLD complicated with liver fibrosis. Spearman correlation analysis revealed that liver fibrosis in MAFLD was positively correlated with the serum ANGPTL8 Level (r = 0.489, P < 0.001) and TyG index (r = 0.294, P = 0.008), and negatively correlated with the CRP level (r = -0.238, P = 0.033, Table 4). Furthermore, logistic regression analysis revealed that elevated serum ANGPTL8 level and TyG index were risk factors for liver fibrosis in MAFLD (both P < 0.05, Table 5).

# Diagnostic performance of ANGPTL8 and TyG index in MAFLD with liver fibrosis

ROC curve analysis was performed to evaluate the predictive value of ANGPTL8, TyG index, and their combination for liver fibrosis in MAFLD (Table 6, Figure 2). The results showed that the AUC of ANGPTL8 was 0.812 (95%CI: 0.713-0.910, P < 0.001), with an optimal cutoff point of 44.36 pg/mL, sensitivity of 91.3%, and specificity of 61.36%. The AUC of TyG index was 0.688 (95%CI: 0.553-0.822, P < 0.001), with an optimal cutoff point of 9.925, sensitivity of 60.87%, and specificity

Table 4 Spearman correlation of metabolic dysfunction-associated fatty liver disease complicated with liver fibrosis with serum angiopoietin-like protein 8, C-reactive protein, and triglyceride-glucose index					
	r	<i>P</i> value <sup>a</sup>			
ANGPTL8	0.489	< 0.001			
TyG index	0.294	0.008			
CRP	-0.238	0.033			

 $^{a}P < 0.05$  is considered significant.

ANGPTL8: Angiopoietin-like protein 8; CRP: C-reactive protein; TyG index: Triglyceride-glucose index.

Table 5 Logistical analysis of metabolic dysfunction-associated fatty liver disease with liver fibrosis							
	В	S.E.	Wals	P value <sup>a</sup>	EXP(B)	95%CI	
ANGPTL8	0.089	0.023	14.513	< 0.001	1.093	1.044-1.144	
TyG index	0868	0.350	6.146	0.013	2.383	1.199-4.736	
CRP	-0.560	0.302	3.449	0.063	0.571	0.316-1.032	

<sup>a</sup>P < 0.05 is considered significant.

CI: Confidence interval; ANGPTL8: Angiopoietin-like protein 8; TyG index: Triglyceride-glucose index; CRP: C-reactive protein.

Table 6 Receiver operating characteristic curve analysis of the predictive value of angiopoietin-like protein 8, triglyceride-glucose index, and its combination with triglyceride-glucose index for metabolic dysfunction-associated fatty liver disease and liver fibrosis

	AUC	95%CI	Cutoff	Sensitivity	Specificity	P value <sup>a</sup>	
Prediction of MAFLD							
ANGPTL8	0.832	0.771-0.893	31.18	80.00	73.08	< 0.001	
TyG index	0.773	0.701-0.845	9.155	67.5	80.77	< 0.001	
ANGPTL8 + TyG index	0.886	0.846-0.941	0.546	75.5	85.9	< 0.001	
Prediction of MAFLD with liver fibrosis							
ANGPTL8	0.812	0.713-0.910	44.36	91.30	61.36	< 0.001	
TyG index	0.688	0.553-0.822	9.925	60.87	75.44	< 0.001	
ANGPTL8 + TyG index	0.835	0.735-0.935	0.3897	73.91	78.95	< 0.001	

 $^{a}P < 0.05$  is considered significant.

AUC: Area under the curve; MAFLD: Metabolic dysfunction-associated fatty liver disease; CI: Confidence interval; ANGPTL8: Angiopoietin-like protein 8; TyG index: Triglyceride-glucose index.

of 75.44%. The AUC of their combination was 0.835 (95%CI: 0.735-0.935, P < 0.001), with an optimal cutoff point of 0.3897, sensitivity of 73.91%, and specificity of 78.95%.

# DISCUSSION

The data from this study indicate that serum ANGPTL8 is significantly elevated in MAFLD patients and is closely associated with the risk of MAFLD and MAFLD-associated liver fibrosis. As serum ANGPTL8 levels increase, the risks of MAFLD and liver fibrosis also increase. Hence, serum ANGPTL8 may be a powerful indicator for predicting the risk of MAFLD and MAFLD-associated liver fibrosis. The combination of serum ANGPTL8 and the TyG index may have an even higher predictive value for the risk of MAFLD and the presence of liver fibrosis.

MAFLD is a progressive liver disease that can lead to steatohepatitis, liver fibrosis, cirrhosis, and even hepatocellular carcinoma. Its pathophysiology is complex and involves many interconnected processes, including metabolic dysregulation, lipotoxicity, insulin resistance, chronic inflammation, oxidative stress, mitochondrial autophagy, and gut microbiota[11-13]. Currently, MAFLD has become an increasingly common factor in end-stage liver disease and

Baishidena® WJD | https://www.wjgnet.com



Figure 3 Comparison of 3 indexs between the liver fibrosis and non-fibrosis groups. A: Angiopoietin-like protein 8; B: Triglyceride-glucose index levels; C: C-reactive protein. MAFLD: Metabolic dysfunction-associated fatty liver disease; ANGPTL8: Angiopoietin-like protein 8; CRP: C-reactive protein.

cardiovascular disease, posing a significant economic burden on public health[14,15].

Studies have suggested that ANGPTL8, as a novel protein secreted by the liver or adipose tissue, plays a dual role in lipid and glucose metabolism. The latest research has proposed a dual mode of ANGPTL8 regulation of lipoprotein lipase (LPL), known as the "ANGPTL3-4-8" model[16]. ANGPTL8 can inhibit LPL activity; control postprandial TG flux, and lipid uptake and storage in brown adipose tissue through synergistic interactions with ANGPTL3 and ANGPTL4; reduce TG clearance; and ultimately lead to a significant increase in serum TG levels[17]. Animal experiments also support a positive correlation between ANGPTL8 and TG levels; knockout of ANGPTL8 leads to decreased lipid content in adipocytes[18], which can significantly improve lipid uptake and reduce lipid deposition[4]. It is worth noting that ANGPTL8 is also associated with TC, HDL-C, and LDL-C[19]. Serum ANGPTL8 can also improve insulin resistance by directly activating insulin-mediated AKT phosphorylation[20] or regulating the PI3K/Akt signaling pathway to enhance insulin sensitivity[21], promote glycogen synthesis, and inhibit gluconeogenesis. All of the above findings indicate that ANGPTL8 is an important link in glucose and lipid metabolism, and may be involved in the development of MAFLD.

Research has shown that the ANGPTL8 level is significantly increased in patients with MAFLD[22], and it is positively correlated with the lipid content in liver cells[23]. In a mouse experiment, it was discovered that inhibiting serum ANGPTL8 levels with antisense oligonucleotides can suppress hepatic steatosis in high-fat-fed mice[4]. This further suggests that ANGPTL8 may play an important role in the occurrence and development of MAFLD. ANGPTL8 expression is increased in the livers of MAFLD patients, which is possibly related to endoplasmic reticulum stress in liver cells[24]. ANGPTL8 can also induce the expression of inflammatory factors by regulating the NF-kB signaling pathway [25], thereby participating in the inflammatory response and promoting the progression of MAFLD to liver fibrosis and cirrhosis. The present study also found similar results by comparing the clinical characteristics of the non-MAFLD and MAFLD populations. Serum ANGPTL8 was positively correlated with the risk of MAFLD, and as serum ANGPTL8 levels increased, the risk of MAFLD also increased. Logistic regression analysis showed that ANGPTL8 is a risk factor for MAFLD. The TyG index, which reflects insulin resistance, has been proven to be of significant value in predicting the risk of MAFLD and cardiovascular risk factors such as hypertension and atherosclerosis. In our study, we also found a close correlation between the TyG index and the risk of MAFLD. We further constructed ROC curves for predicting MAFLD by using ANGPTL8 alone and in combination with the TyG index, and obtained AUCs of 0.832 and 0.886, respectively. This indicates that ANGPTL8 can serve as a predictive factor for the risk of MAFLD, and its combination with the TyG index may have an even higher predictive value for the risk of MAFLD.

FIB-4 is a widely used non-invasive index for assessing liver fibrosis, with diagnostic efficacy comparable to the gold standard liver biopsy[26]. Studies have proposed that a FIB-4 value of < 1.45 can exclude progressive fibrosis, and this has been widely validated[9,10]. ANGPTL8 can promote the occurrence and development of MAFLD by participating in glucose and lipid metabolism, insulin resistance, and energy homeostasis[22,27], but the relationship between ANGPTL8 and liver fibrosis remains controversial. In 2016, Cengiz *et al*[28] found that serum ANGPTL8 levels were lower in NAFLD patients than in healthy individuals, and higher in NAFLD patients with mild fibrosis than in patients with significant fibrosis. However, some authors have reported contradictory conclusions. Ke *et al*[22] reported that the circulating level of ANGPTL8 is significantly higher in MAFLD patients than in healthy individuals, and is not affected by race, region, BMI, *etc.*; they concluded that elevated serum ANGPTL8 is positively correlated with MAFLD. In addition, Zhang *et al*[29] suggested that ANGPTL8 can act as a pro-inflammatory factor, and promote the development of liver fibrosis. In our study, we found similar results: ANGPTL8 levels were significantly increased in MAFLD patients with liver fibrosis. Hence, we concluded that elevated serum ANGPTL8 levels are associated with an increased risk of liver fibrosis in MAFLD patients.

Studies have also shown a significant correlation between the TyG index and the severity of hepatic steatosis and liver fibrosis in NAFLD patients[30]. Elevated TyG index is an independent predictor of MAFLD and is positively correlated with the MAFLD fibrosis score and FIB-4 index, suggesting that a significant increase in the TyG index may indicate the development of liver fibrosis[31]. In 113 patients with histologically confirmed NAFLD, the TyG index levels were significantly higher in the F3 fibrosis stage than in the F0–F1 stage (P < 0.0001)[32]. However, Guo *et al*[30] found that the AUC value of the TyG index for predicting NAFLD was 0.761, while its AUC value for predicting liver fibrosis. In our study, indicating that the TyG index has a relatively low accuracy for predicting NAFLD-associated liver fibrosis. In our study,

we constructed ROC curves to analyze whether serum ANGPTL8 as well as the combination of serum ANGPTL8 and the TyG index could predict the coexistence of liver fibrosis with MAFLD, and obtained AUC values of 0.812 and 0.835, respectively. Hence, serum ANGPTL8 is a powerful predictor of the coexistence of liver fibrosis with MAFLD, and the combined use of serum ANGPTL8 and the TyG index may improve the predictive efficacy for liver fibrosis in MAFLD.

This study has the following limitations: First, the gold standard for the diagnosis of MAFLD and liver fibrosis is liver biopsy, but this study only used imaging examinations and the FIB-4 index due to limitations in conditions. Second, this study had a small sample size, and the results may be biased and require further validation with a larger sample size. Third, this study is a cross-sectional study and cannot reflect the specific mechanisms of ANGPTL8 in MAFLD. Further prospective studies are needed to explore and clarify these mechanisms.

# CONCLUSION

In conclusion, the study reveals that serum levels of ANGPTL8 are elevated in patients diagnosed with MAFLD, and notably higher in those with concurrent liver fibrosis. The close correlation of serum ANGPTL8 with the risks associated with MAFLD and the progression of liver fibrosis underscores its significance. Serving as a predictive marker for both MAFLD risks and its associated liver fibrosis, serum ANGPTL8 exhibits enhanced predictive value when combined with the TyG index. Consequently, we posit that serum ANGPTL8 stands as a valuable and straightforward reference marker for the early prediction and timely management of MAFLD, enriching the repertoire of non-invasive assessment methods in clinical settings. Monitoring changes in ANGPTL8 levels facilitates early detection of MAFLD-associated liver fibrosis, enabling prompt intervention and management strategies to mitigate disease progression and reduce the risk of mortality.

# ARTICLE HIGHLIGHTS

#### Research background

Metabolic dysfunction-associated fatty liver disease (MAFLD) is now recognized as a prevalent global chronic liver disorder, standing as the primary contributor to end-stage liver disease and cardiovascular complications. This condition imposes a significant economic burden on public health. Previous research has elucidated the pivotal role of angiopoietinlike protein 8 (ANGPTL8) in the pathogenesis and progression of MAFLD, showcasing variations across different disease stages. Understanding ANGPTL8's involvement in MAFLD development informs early intervention strategies and aids in reducing mortality.

#### Research motivation

This study was motivated by the need for a convenient laboratory indicator facilitating the early identification of MAFLD and dynamic monitoring of its progression. Providing early intervention strategies for MAFLD patients is crucial to delay disease progression and reduce mortality.

#### Research objectives

The primary objective of this study was to conduct a comparative analysis of serum ANGPTL8 levels between individuals with and without MAFLD, investigating the predictive value of ANGPTL8 in relation to MAFLD development and progression.

#### Research methods

In this cross-sectional study, 160 patients were enrolled, with 80 (50%) diagnosed with MAFLD. MAFLD patients were further categorized into hepatic fibrosis and non-hepatic fibrosis groups based on the fibrosis-4 index. Logistic regression analysis and receiver operating characteristic curves were employed to explore the impact and predictive ability of serum ANGPTL8.

#### Research results

Compared with the non-MAFLD group, serum ANGPTL8 levels and the triglyceride-glucose (TyG) index were significantly elevated in the MAFLD group, positively correlated with MAFLD. The combined ANGPTL8 and TyG index showed high predictive accuracy for MAFLD. Similarly, in the liver fibrosis group, both ANGPTL8 and the TyG index were significantly increased, positively correlated with liver fibrosis, with robust predictive accuracy for MAFLDassociated liver fibrosis.

#### Research conclusions

Serum ANGPTL8 appears pivotal in the pathogenesis and progression of MAFLD, emerging as a potential biomarker for predicting both MAFLD and its associated hepatic fibrosis. The combined assessment of serum ANGPTL8 levels and the TyG index enhances predictive accuracy. Understanding the underlying mechanisms linking ANGPTL8 with MAFLD provides valuable insights for early diagnosis, risk stratification, and timely intervention, ultimately alleviating the burden of this disease.



WJD https://www.wjgnet.com

#### Research perspectives

Future studies should consider expanding sample sizes and incorporating liver biopsy and other methods to distinguish MAFLD-related liver fibrosis. Longitudinal studies are necessary to analyze serum ANGPTL8 changes at each MAFLD stage, confirming the relationship between serum ANGPTL8 and the occurrence and progression of MAFLD.

# ACKNOWLEDGEMENTS

We express our gratitude to all the medical and technical personnel who willingly agreed to participate in this study.

#### FOOTNOTES

Author contributions: Gan LL participated in the study design and wrote the manuscript; Yan YM conducted the design of the study and reviewed/edited the drafts, and is guarantor; Gan LL, Zhu X, Xia C, Gao Y and Wu WC collected and analyzed the data; Li Q, Li L and Dai Z revised the manuscript. All authors contributed to the article and approved the submitted article.

Supported by Youth Talents Project of Joint Fund of Hubei Health Commission, No. WJ2019H170; and Xiaogan Natural Science Project, No. XGKJ2020010033.

Institutional review board statement: The study protocol was approved by the ethics committee of Xiaogan Central Hospital (Approval No. XGLY2021-08-01).

Informed consent statement: All study participants or their legal guardian provided informed written consent about personal and medical data collection prior to study enrolment.

Conflict-of-interest statement: The authors have no conflicts of interest to declare.

Data sharing statement: The datasets generated during and/or analyzed during the current study are not publicly available, but are available from the corresponding author on reasonable request.

STROBE statement: The authors have read the STROBE Statement-checklist of items, and the manuscript was prepared and revised according to the STROBE Statement-checklist of items.

**Open-Access:** This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: https://creativecommons.org/Licenses/by-nc/4.0/

#### Country/Territory of origin: China

**ORCID** number: Lu-Lu Gan 0000-0001-8847-0354; Can Xia 0009-0006-5133-5128; Xuan Zhu 0000-0001-6671-897X; Yue Gao 0000-0002-0156-8752; Wen-Chang Wu 0009-0009-5323-367X; Qi Li 0009-0006-0979-2253; Ling Li 0009-0008-7539-2885; Zhe Dai 0000-0003-0321-0225; Yi-Min Yan 0000-0002-3351-4732.

S-Editor: Ou XL L-Editor: A P-Editor: Zhao YQ

#### REFERENCES

- Eslam M, Sanyal AJ, George J; International Consensus Panel. MAFLD: A Consensus-Driven Proposed Nomenclature for Metabolic 1 Associated Fatty Liver Disease. Gastroenterology 2020; 158: 1999-2014.e1 [PMID: 32044314 DOI: 10.1053/j.gastro.2019.11.312]
- Simon TG, Roelstraete B, Khalili H, Hagström H, Ludvigsson JF. Mortality in biopsy-confirmed nonalcoholic fatty liver disease: results from 2 a nationwide cohort. Gut 2021; 70: 1375-1382 [PMID: 33037056 DOI: 10.1136/gutjnl-2020-322786]
- Xie ZQ, Li HX, Wang BK, Yang ZM, Zhang ZY, Tan WL, Li WX, Wang QB, Yang L, Zhuang HK, Tang CW, Shang CZ, Chen YJ. Trends in 3 prevalence and all-cause mortality of metabolic dysfunction-associated fatty liver disease among adults in the past three decades: Results from the NHANES study. Eur J Intern Med 2023; 110: 62-70 [PMID: 36754655 DOI: 10.1016/j.ejim.2023.01.029]
- 4 Vatner DF, Goedeke L, Camporez JG, Lyu K, Nasiri AR, Zhang D, Bhanot S, Murray SF, Still CD, Gerhard GS, Shulman GI, Samuel VT. Angptl8 antisense oligonucleotide improves adipose lipid metabolism and prevents diet-induced NAFLD and hepatic insulin resistance in rodents. Diabetologia 2018; 61: 1435-1446 [PMID: 29497783 DOI: 10.1007/s00125-018-4579-1]
- Zhang Y, Zheng L, Huang K. A new way to regulate inflammation: selective autophagic degradation of IKKy mediated by ANGPTL8. Cell 5 Stress 2018; 2: 66-68 [PMID: 31225468 DOI: 10.15698/cst2018.03.128]
- Yin Y, Ding X, Peng L, Hou Y, Ling Y, Gu M, Wang Y, Peng Y, Sun H. Increased Serum ANGPTL8 Concentrations in Patients with 6

WJD | https://www.wjgnet.com

Prediabetes and Type 2 Diabetes. J Diabetes Res 2017; 2017: 8293207 [PMID: 29082263 DOI: 10.1155/2017/8293207]

- 7 Ye J, Qin Y, Wang D, Yang L, Yuan G. The Relationship between Circulating ANGPTL8/Betatrophin Concentrations and Adult Obesity: A Meta-Analysis. *Dis Markers* 2019; 2019: 5096860 [PMID: 31772689 DOI: 10.1155/2019/5096860]
- 8 Wang H, Lai Y, Han C, Liu A, Fan C, Wang H, Zhang H, Ding S, Teng W, Shan Z. The Effects of Serum ANGPTL8/betatrophin on the Risk of Developing the Metabolic Syndrome A Prospective Study. *Sci Rep* 2016; **6**: 28431 [PMID: 27345212 DOI: 10.1038/srep28431]
- 9 Vallet-Pichard A, Mallet V, Nalpas B, Verkarre V, Nalpas A, Dhalluin-Venier V, Fontaine H, Pol S. FIB-4: an inexpensive and accurate marker of fibrosis in HCV infection. comparison with liver biopsy and fibrotest. *Hepatology* 2007; 46: 32-36 [PMID: 17567829 DOI: 10.1002/hep.21669]
- 10 Chinese Society of Hepatology; Chinese Medical Association; Chinese Society of Gastroenterology, Chinese Medical Association; Chinese Society of Infectious Diseases, Chinese Medical Association. Consensus on the diagnosis and treatment of hepatic fibrosis (2019). J Dig Dis 2020; 21: 127-138 [PMID: 32108989 DOI: 10.1111/1751-2980.12854]
- 11 Nassir F. NAFLD: Mechanisms, Treatments, and Biomarkers. *Biomolecules* 2022; 12 [PMID: 35740949 DOI: 10.3390/biom12060824]
- 12 Ji J, Wu L, Wei J, Wu J, Guo C. The Gut Microbiome and Ferroptosis in MAFLD. J Clin Transl Hepatol 2023; 11: 174-187 [PMID: 36406312 DOI: 10.14218/JCTH.2022.00136]
- Clare K, Dillon JF, Brennan PN. Reactive Oxygen Species and Oxidative Stress in the Pathogenesis of MAFLD. J Clin Transl Hepatol 2022;
  10: 939-946 [PMID: 36304513 DOI: 10.14218/JCTH.2022.00067]
- 14 Zhang P, Dong X, Zhang W, Wang S, Chen C, Tang J, You Y, Hu S, Zhang S, Wang C, Wen W, Zhou M, Tan T, Qi G, Li L, Wang M. Metabolic-associated fatty liver disease and the risk of cardiovascular disease. *Clin Res Hepatol Gastroenterol* 2023; 47: 102063 [PMID: 36494073 DOI: 10.1016/j.clinre.2022.102063]
- 15 Powell EE, Wong VW, Rinella M. Non-alcoholic fatty liver disease. Lancet 2021; 397: 2212-2224 [PMID: 33894145 DOI: 10.1016/S0140-6736(20)32511-3]
- 16 Abu-Farha M, Ghosh A, Al-Khairi I, Madiraju SRM, Abubaker J, Prentki M. The multi-faces of Angptl8 in health and disease: Novel functions beyond lipoprotein lipase modulation. *Prog Lipid Res* 2020; 80: 101067 [PMID: 33011191 DOI: 10.1016/j.plipres.2020.101067]
- 17 **Zhang R**, Zhang K. An updated ANGPTL3-4-8 model as a mechanism of triglyceride partitioning between fat and oxidative tissues. *Prog Lipid Res* 2022; **85**: 101140 [PMID: 34793860 DOI: 10.1016/j.plipres.2021.101140]
- 18 Luo D, Chen X, Yang W, Ran W, Wen Z. Angiopoietin-like 8 Improves Insulin Resistance and Attenuates Adipose Tissue Inflammation in Diet-Induced Obese Mice. *Exp Clin Endocrinol Diabetes* 2020; **128**: 290-296 [PMID: 30257264 DOI: 10.1055/a-0725-7897]
- 19 Yang S, Jiao X, Huo X, Zhu M, Wang Y, Fang X, Yang Y, Yue W, Qin Y. Association between circulating full-length angiopoietin-like protein 8 and non-high-density lipoprotein cholesterol levels in Chinese non-diabetic individuals: a cross-sectional study. *Lipids Health Dis* 2018; 17: 161 [PMID: 30021605 DOI: 10.1186/s12944-018-0802-9]
- Yuan J, Zhang D, Wang Y, Zhu Z, Lin Q, Li M, Zhong W, Han J, Xu F, Dong J. Angiopoietin-Like 8 in Gestational Diabetes Mellitus: Reduced Levels in Third Trimester Maternal Serum and Placenta, Increased Levels in Cord Blood Serum. *Int J Endocrinol* 2022; 2022: 1113811 [PMID: 35529083 DOI: 10.1155/2022/1113811]
- 21 Zhao Z, Deng X, Jia J, Zhao L, Wang C, Cai Z, Guo C, Yang L, Wang D, Ma S, Deng J, Li H, Zhou L, Tu Z, Yuan G. Angiopoietin-like protein 8 (betatrophin) inhibits hepatic gluconeogenesis through PI3K/Akt signaling pathway in diabetic mice. *Metabolism* 2022; 126: 154921 [PMID: 34715116 DOI: 10.1016/j.metabol.2021.154921]
- 22 Ke Y, Liu S, Zhang Z, Hu J. Circulating angiopoietin-like proteins in metabolic-associated fatty liver disease: a systematic review and metaanalysis. *Lipids Health Dis* 2021; 20: 55 [PMID: 34034750 DOI: 10.1186/s12944-021-01481-1]
- Hong BS, Liu J, Zheng J, Ke W, Huang Z, Wan X, He X, Xiao H, Li Y. Angiopoietin-like protein 8/betatrophin correlates with hepatocellular lipid content independent of insulin resistance in non-alcoholic fatty liver disease patients. *J Diabetes Investig* 2018; 9: 952-958 [PMID: 29266821 DOI: 10.1111/jdi.12792]
- 24 Lee YH, Lee SG, Lee CJ, Kim SH, Song YM, Yoon MR, Jeon BH, Lee JH, Lee BW, Kang ES, Lee HC, Cha BS. Association between betatrophin/ANGPTL8 and non-alcoholic fatty liver disease: animal and human studies. *Sci Rep* 2016; 6: 24013 [PMID: 27045862 DOI: 10.1038/srep24013]
- Liao Z, Wu X, Song Y, Luo R, Yin H, Zhan S, Li S, Wang K, Zhang Y, Yang C. Angiopoietin-like protein 8 expression and association with extracellular matrix metabolism and inflammation during intervertebral disc degeneration. J Cell Mol Med 2019; 23: 5737-5750 [PMID: 31211513 DOI: 10.1111/jcmm.14488]
- 26 Lee J, Vali Y, Boursier J, Spijker R, Anstee QM, Bossuyt PM, Zafarmand MH. Prognostic accuracy of FIB-4, NAFLD fibrosis score and APRI for NAFLD-related events: A systematic review. *Liver Int* 2021; 41: 261-270 [PMID: 32946642 DOI: 10.1111/liv.14669]
- 27 Luo M, Peng D. ANGPTL8: An Important Regulator in Metabolic Disorders. Front Endocrinol (Lausanne) 2018; 9: 169 [PMID: 29719529 DOI: 10.3389/fendo.2018.00169]
- 28 Cengiz M, Ozenirler S, Kocabiyik M. Serum β-trophin level as a new marker for noninvasive assessment of nonalcoholic fatty liver disease and liver fibrosis. *Eur J Gastroenterol Hepatol* 2016; 28: 57-63 [PMID: 26513612 DOI: 10.1097/MEG.000000000000502]
- 29 Zhang Z, Yuan Y, Hu L, Tang J, Meng Z, Dai L, Gao Y, Ma S, Wang X, Zhang Q, Cai W, Ruan X, Guo X. ANGPTL8 accelerates liver fibrosis mediated by HFD-induced inflammatory activity *via* LILRB2/ERK signaling pathways. *J Adv Res* 2023; 47: 41-56 [PMID: 36031141 DOI: 10.1016/j.jare.2022.08.006]
- 30 Guo W, Lu J, Qin P, Li X, Zhu W, Wu J, Xu N, Zhang Q. The triglyceride-glucose index is associated with the severity of hepatic steatosis and the presence of liver fibrosis in non-alcoholic fatty liver disease: a cross-sectional study in Chinese adults. *Lipids Health Dis* 2020; 19: 218 [PMID: 33028338 DOI: 10.1186/s12944-020-01393-6]
- 31 Tutunchi H, Naeini F, Mobasseri M, Ostadrahimi A. Triglyceride glucose (TyG) index and the progression of liver fibrosis: A cross-sectional study. *Clin Nutr ESPEN* 2021; 44: 483-487 [PMID: 34330512 DOI: 10.1016/j.clnesp.2021.04.025]
- 32 Amzolini AM, Forţofoiu MC, Barău Abu-Alhija A, Vladu IM, Clenciu D, Mitrea A, Forţofoiu M, Matei D, Enăchescu V, Predescu OI, Micu ES. Triglyceride and glucose index: a useful tool for non-alcoholic liver disease assessed by liver biopsy in patients with metabolic syndrome? Rom J Morphol Embryol 2021; 62: 475-480 [PMID: 35024735 DOI: 10.47162/RJME.62.2.13]

Saisbideng® WJD | https://www.wjgnet.com



# Published by Baishideng Publishing Group Inc 7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA Telephone: +1-925-3991568 E-mail: office@baishideng.com Help Desk: https://www.f6publishing.com/helpdesk https://www.wjgnet.com

