



Poor oral health was associated with higher risk of gastric cancer: Evidence from 1431677 participants

Fei Liu, Shi-Jun Tang, Zi-Wei Li, Xu-Rui Liu, Quan Lv, Wei Zhang, Dong Peng

Specialty type: Gastroenterology and hepatology

Provenance and peer review: Unsolicited article; Externally peer reviewed.

Peer-review model: Single blind

Peer-review report's scientific quality classification

Grade A (Excellent): 0
Grade B (Very good): 0
Grade C (Good): C
Grade D (Fair): D
Grade E (Poor): 0

P-Reviewer: Arigami T, Japan;
Corte-Real A, Portugal

Received: November 2, 2023

Peer-review started: November 2, 2023

First decision: November 30, 2023

Revised: December 26, 2023

Accepted: January 29, 2024

Article in press: January 29, 2024

Published online: February 27, 2024



Fei Liu, Zi-Wei Li, Xu-Rui Liu, Quan Lv, Wei Zhang, Dong Peng, Department of Gastrointestinal Surgery, The First Affiliated Hospital of Chongqing Medical University, Chongqing 400016, China

Shi-Jun Tang, Department of Pediatric Dentistry, Stomatological Hospital of Guizhou Medical University, Guizhou 550000, Guizhou Province, China

Corresponding author: Dong Peng, FAASLD, Academic Editor, Department of Gastrointestinal Surgery, The First Affiliated Hospital of Chongqing Medical University, No. 1 Youyi Road, Yuanjiagang, Yuzhong District, Chongqing 400016, China. carry_dong@126.com

Abstract

BACKGROUND

In recent years, the association between oral health and the risk of gastric cancer (GC) has gradually attracted increased interest. However, in terms of GC incidence, the association between oral health and GC incidence remains controversial. Periodontitis is reported to increase the risk of GC. However, some studies have shown that periodontitis has no effect on the risk of GC. Therefore, the present study aimed to assess whether there is a relationship between oral health and the risk of GC.

AIM

To assess whether there was a relationship between oral health and the risk of GC.

METHODS

Five databases were searched to find eligible studies from inception to April 10, 2023. Newcastle-Ottawa Scale score was used to assess the quality of included studies. The quality of cohort studies and case-control studies were evaluated separately in this study. Incidence of GC were described by odds ratio (OR) and 95% confidence interval (CI). Funnel plot was used to represent the publication bias of included studies. We performed the data analysis by StataSE 16.

RESULTS

A total of 1431677 patients from twelve included studies were enrolled for data analysis in this study. According to our analysis, we found that the poor oral health was associated with higher risk of GC (OR = 1.15, 95%CI: 1.02-1.29; $I^2 = 59.47\%$, $P = 0.00 < 0.01$). Moreover, after subgroup analysis, the outcomes showed that whether tooth loss (OR = 1.12, 95%CI: 0.94-1.29; $I^2 = 6.01\%$, $P > 0.01$), gingivitis (OR = 1.19, 95%CI: 0.71-1.67; $I^2 = 0.00\%$, $P > 0.01$), dentures (OR = 1.27,

95%CI: 0.63-1.19; $I^2 = 68.79\%$, $P > 0.01$), or tooth brushing (OR = 1.25, 95%CI: 0.78-1.71; $I^2 = 88.87\%$, $P > 0.01$) had no influence on the risk of GC. However, patients with periodontitis (OR = 1.13, 95%CI: 1.04-1.23; $I^2 = 0.00\%$, $P < 0.01$) had a higher risk of GC.

CONCLUSION

Patients with poor oral health, especially periodontitis, had a higher risk of GC. Patients should be concerned about their oral health. Improving oral health might reduce the risk of GC.

Key Words: Oral health; Tooth loss; Periodontitis; Gastric cancer; Risk factor

©The Author(s) 2024. Published by Baishideng Publishing Group Inc. All rights reserved.

Core Tip: The aim of this current study was to assess whether there was a relationship between oral health and the risk of gastric cancer (GC). A total of 1431677 patients from twelve included studies were enrolled for data analysis in this study. This article summarised all the papers over the years on the relationship between oral health and the incidence of GC. After analysing them, the existing controversies were resolved to some extent. It was useful to guide clinical work.

Citation: Liu F, Tang SJ, Li ZW, Liu XR, Lv Q, Zhang W, Peng D. Poor oral health was associated with higher risk of gastric cancer: Evidence from 1431677 participants. *World J Gastrointest Surg* 2024; 16(2): 585-595

URL: <https://www.wjgnet.com/1948-9366/full/v16/i2/585.htm>

DOI: <https://dx.doi.org/10.4240/wjgs.v16.i2.585>

INTRODUCTION

Gastric cancer (GC) is one of the most common tumours worldwide and the forth leading cause of cancer death[1-3]. The incidence and mortality of GC continue to increase, and there are approximately 1 million new cases worldwide each year [3-5]. In China, more than 400000 new cases are diagnosed each year, accounting for 50% of new cases worldwide[6-8]. Prevention of GC has become a focal point because of these worrisome numbers. Prevention of GC can be divided into primary prevention (reducing the incidence of GC) and secondary prevention (early detection and treatment). Primary prevention includes smoking cessation, reducing salt intake, increasing fruit and vegetable intake, and other health behaviours, such as oral health behaviours[9].

The oral cavity is the conduit between the external environment and the gastrointestinal tract and is involved in the intake and digestion of food. Oral hygiene plays an important role in human health. Measures of oral health included tooth loss, periodontitis, gingivitis, dentures, and tooth brushing. Poor oral health has been shown to be a risk factor for many diseases, including cardiovascular disease, atherosclerosis, oral cancer, kidney cancer, lung cancer, oesophageal cancer, and pancreatic cancer[10-18].

In recent years, the association between oral health and the risk of GC has gradually attracted increased interest. However, in terms of GC incidence, the association between oral health and GC incidence remains controversial. Periodontitis is reported to increase the risk of GC[19]. However, some studies have shown that periodontitis has no effect on the risk of GC[20,21]. Therefore, the present study aimed to assess whether there is a relationship between oral health and the risk of GC.

MATERIALS AND METHODS

Methods

Our study was produced in accordance with the Preferred Reporting Items for Systematic Reviews and Meta-Analyses statement[22].

Search strategy

The PubMed, EMBASE, Cochrane Library, MEDLINE, and Ovid databases were searched from inception to April 10, 2023. The two keywords used were oral health and GC. For oral health, the search strategy was as follows: "dental" OR "oral" OR "oral health" OR "oral hygiene behaviour" OR "oral hygiene" OR "oral behaviour" OR "tooth loss" OR "tooth missing" OR "dental caries" OR "full teeth" OR "salivary flow" OR "probing depth" OR "periodontal disease" OR "periodontitis" OR "gingivitis" OR "dentures" OR "tooth brushing". In terms of GC, "gastric cancer" OR "gastric carcinoma" OR "gastric neoplasms" OR "stomach cancer" OR "stomach carcinoma" OR "stomach neoplasms" were searched. Then, we used "AND" to combine the two keywords. The language was limited to English.

Inclusion and exclusion criteria

The inclusion and exclusion criteria were established to find eligible studies. The inclusion criteria for patients were as follows: (1) Patients were reported to have oral health problems; and (2) The incidence of GC was reported. The exclusion criteria for patients were as follows: (1) Had case reports, comments, letters to the editor, or conference abstracts; (2) Had data repeated or overlapped; and (3) Incomplete information.

Study selection

The database search was independently conducted by two authors. The steps for screening eligible studies were as follows: (1) Excluded duplicate studies; (2) Scanned the titles and abstracts; and (3) Read the full text, including the reference. All disagreements were settled by group discussion.

Data collection

The baseline characteristics of the individuals included in the studies and the incidence of GC were collected for data analysis in the present study. The baseline information of the enrolled studies included author, publication year, publication country, study period, sample size, study type, follow-up period, diagnosis of GC, definition of oral health, and Newcastle-Ottawa scale (NOS) score.

Quality assessment

The quality of the included studies was evaluated by the NOS score[23]. According to the NOS score, we divided studies into high quality (9 points), median quality (7-8 points), and low quality (< 7 points) groups. We evaluated the quality of the cohort studies and case-control studies separately in this study.

Statistical analysis

We defaulted the risk ratio (RR) and hazard ratio (HR) of GC reported in the included studies to be equivalent to the odds ratio (OR)[24]. The incidence of GC was described by the OR and 95% confidence interval (CI). I^2 values and χ^2 tests were used to assess the statistical heterogeneity[25,26]. The random effects DerSimonian-Laird model was used only. When a random effects model was used, $P < 0.01$ was considered to indicate statistical significance. StataSE 16 was used for the data analysis.

RESULTS

Study selection

A total of 13279 studies were identified from five databases (2652 in PubMed, 3982 in EMBASE, 1262 in the Cochrane Library, 2416 in MEDLINE, 2967 in Ovid, and two from the citation searching) from inception to April 10, 2023. After duplicate removal, 5509 records remained. Three thousand forty-one unqualified studies were excluded according to the exclusion criteria. After excluding unqualified studies, 2168 studies were needed for eligibility, and six records were not retrieved. Finally, twelve eligible studies were included in this study[8,19-21,27-34] (Figure 1).

Baseline characteristics of the included studies

Twelve studies involving 1431677 patients were included in this study. The publication years ranged from 1998 to 2022. The published countries were mainly in China, Japan, United States, and Iran. The study period ranged from 1973 to 2011. There were nine cohort studies and three case-control studies. There were nine studies reporting tooth loss, three studies reporting periodontitis, two studies reporting gingivitis, three studies reporting dentures, and five studies reporting tooth brushing. More details of the included studies' baseline characteristics, including the author, the number of patients, the follow-up period, the diagnosis of GC, and the NOS score, are shown in Table 1.

Tables 2 and 3 show the results of the quality assessment by the NOS score for cohort studies and case-control studies, respectively. For cohort studies, three studies were graded as high quality (nine points), five studies were graded as median quality (seven to eight points), and one study was graded as low quality (six points). For case-control studies, one study was graded as high quality (nine points), and two studies were graded as low quality (six points). The details of the quality assessment are shown in Tables 2 and 3.

The association between oral health and risk of GC

Before the data analysis, we adjusted the RR or HR to the OR. The information on the participants' adjustment is shown in Table 4. According to the data analysis, poor oral health could increase the risk of GC (OR = 1.15, 95%CI: 1.02-1.29; I^2 = 59.47%, P = 0.00 < 0.01) (Figure 2).

Subgroup analysis

We classified oral health into five subgroups and analysed their respective effects on the risk of GC. We found that patients with periodontitis (OR = 1.13, 95%CI: 1.04-1.23; I^2 = 0.00%, P < 0.01) had a greater risk of GC. However, tooth loss (OR = 1.12, 95%CI: 0.94-1.29; I^2 = 6.01%, P > 0.01), gingivitis (OR = 1.19, 95%CI: 0.71-1.67; I^2 = 0.00%, P > 0.01), dentures (OR = 1.27, 95%CI: 0.63-1.19, I^2 = 68.79%, P > 0.01), and tooth brushing (OR = 1.25, 95%CI: 0.78-1.71, I^2 = 88.87%, P > 0.01) had no effect on the risk of GC (Figure 3).

Table 1 Baseline characteristics of included studies

Ref.	Country	Study date	Patients	Study type	Follow-up	Diagnosis of gastric cancer	Definition of oral health	NOS
Watabe <i>et al</i> [32], 1998	Japan	October 1996 to September 1997	242	Retrospective cohort study	NA	Gastric cancer	Brush teeth, decayed teeth, gingivitis, bad occlusion, dentures (partial and full), and lack of teeth ≥ 10	6
Abnet <i>et al</i> [27], 2001	China	March 1986 to May 1991	28868	Prospective cohort study	5.25 yr	Gastric cardia tumor and non-cardia tumor	Tooth loss	8
Hujoel <i>et al</i> [20], 2003	United States	1971 to 1992	11328	Prospective cohort study	Until 1992	Gastric cancer (ICD-9 151.0-151.9)	Periodontitis, gingivitis, and edentulism	7
Abnet <i>et al</i> [28], 2005	Finland	1985 to 1999	29124	Prospective cohort study	April 1993 to April 1999	Gastric non-cardia adenocarcinoma	Tooth loss included 0-10 teeth lost, 11-31 teeth lost, and edentulous	8
Michaud <i>et al</i> [29], 2008	United States	1986 to January 2004	49375	Prospective cohort study	Median of 17.7 yr	Gastric cancer	Periodontal disease and tooth loss	9
Hiraki <i>et al</i> [19], 2008	Japan	2000 to 2005	15720	Case-control study	NA	Gastric cancer (ICD-10 C16)	Remaining teeth	7
Shakeri <i>et al</i> [31], 2013	Iran	January 2004 to June 2008	922	Case-control study	December 2004 to December 2011	Gastric adenocarcinoma included non-cardia, cardia, and mixed-locations	Tooth loss, decayed, missing, filled teeth score, and frequency of tooth brushing	6
Ndegwa <i>et al</i> [30], 2018	Sweden	1973 to 1974	19831	Prospective cohort study	569233 person-years	Gastric cancer was divided into cardia (ICD 151.1) and non-cardia gastric cancer (all ICD-7 151 codes except ICD 151.1)	Number of teeth, dental plaque status, and presence of any oral mucosal lesions	7
Yano <i>et al</i> [33], 2021	Iran	January 2004 to June 2008	50045	Prospective cohort study	Until December 31, 2019	Gastric cancer cases were limited to adenocarcinomas (cardia and non-cardia)	Frequency of tooth brushing, tooth loss, and the sum of decayed, missing, or filled teeth	9
Zhang <i>et al</i> [8], 2022	China	October 2010 to September 2013	2873	Case-control	NA	Gastric cancer was divided into esophago-gastric junction cancer and total gastric cancer	Tooth loss after 20 yr, number of tooth loss after age 20 yr, age of first tooth loss after age 20 yr, denture wearing, number of filled teeth, missing and filled teeth, frequency of toothbrushing, frequency of oral discomfort while eating, avoidance of some foods because of oral problems	6
Zhang <i>et al</i> [34], 2022	China	2004 to 2008	510148	Prospective cohort study	Median of 9.17 yr and range of 0.1 to 11.5 yr	Gastric cancer (ICD-10 C16)	Gum bleeding and rarely or never brush teeth	9
Kim <i>et al</i> [21], 2022	South Korea	January 2003 to December 2015	713201	Retrospective cohort study	Up to 10 yr	Gastric cancer (ICD-10 C16)	Periodontitis (who visited a dental clinic two or more than two times within one year and were diagnosed with periodontitis under those ICD-10 codes (K05.2, K05.3, K05.4, K05.5, and K05.6)	9

ICD: International Classification of Diseases; NA: Not available; NOS: Newcastle-Ottawa Scale.

Publication bias

According to the data analysis, the funnel plot was relatively symmetrical, indicating low publication bias (Figure 4).

Sensitivity analysis

Each study was excluded each time the sensitivity was assessed. There were no significant differences in the results after each analysis was performed.

Table 2 Results of quality assessment using the Newcastle-Ottawa Scale for cohort studies

Ref.	Selection				Comparability	Outcome			Scores
	Representativeness of exposure	Selection of the non-exposure	Ascertainment of exposure	Demonstration that outcome was not present at start	Cohorts on the basis of the design or analysis	Assessment	Long follow-up for outcomes to occur	Adequacy of follow-up	
Abnet <i>et al</i> [28], 2005	★	★	★	★	★★	☆	★	★	8
Abnet <i>et al</i> [27], 2001	☆	★	★	★	★★	★	★	★	8
Hiraki <i>et al</i> [19], 2008	★	★	★	★	★★	★	☆	☆	7
Hujoel <i>et al</i> [20], 2003	★	★	★	★	★★	★	☆	☆	7
Kim <i>et al</i> [21], 2022	★	★	★	★	★★	★	★	★	9
Michaud <i>et al</i> [29], 2008	★	★	★	★	★★	★	★	★	9
Ndegwa <i>et al</i> [30], 2018	☆	☆	★	★	★★	★	★	★	7
Watabe <i>et al</i> [32], 1998	☆	★	★	★	★★	★	☆	☆	6
Yano <i>et al</i> [33], 2021	★	★	★	★	★★	★	★	★	9

Table 3 Results of quality assessment using the Newcastle-Ottawa Scale for case-control studies

Ref.	Selection				Comparability	Exposure			Scores
	Adequate definition of cases	Representativeness of the cases	Selection of controls	Definition of controls	Control for important factor ¹	Ascertainment of exposure	Same method of ascertainment for cases and controls	Non-response rate	
Shakeri <i>et al</i> [31], 2013	★	☆	☆	★	★★	☆	★	★	6
Zhang <i>et al</i> [8], 2022	★	☆	☆	★	★★	★	★	☆	6
Zhang <i>et al</i> [34], 2022	★	★	★	★	★★	★	★	★	9

¹A maximum of 2 stars can be allotted in this category, one for age, the other for other controlled factors.




Figure 1 Flowchart of study selection.




Figure 2 The association between oral health and risk of gastric cancer. CI: Confidence interval.

DISCUSSION

A total of 1431808 patients were enrolled from twelve studies in the present study. After the data analysis, the outcomes showed that poor oral health was associated with a greater risk of GC. We classified oral health into five subgroups: Tooth loss, periodontitis, gingivitis, dentures, and tooth brushing. After subgroup analysis, the outcomes showed that

Table 4 The analyses were adjusted for the following variables

Ref.	Variables of adjustment
Abnet <i>et al</i> [28], 2005	Age and education
Abnet <i>et al</i> [27], 2001	Age, sex, tobacco use, and alcohol use
Hiraki <i>et al</i> [19], 2008	Age, sex, smoking and drinking status (never, former, current), vegetable and fruit intake, BMI, and regular exercise
Hujoel <i>et al</i> [20], 2003	Age and gender
Kim <i>et al</i> [21], 2022	Age
Michaud <i>et al</i> [29], 2008	Age (continuous), ethnic origin (white, Asian, black), physical activity (quintiles), history of diabetes (yes or no), alcohol (quartiles), BMI (< 22, 22-24.9, 25-29.9, 30 +), geographical location (south, west, northeast, mid-west), height (quintiles), calcium intake (quintiles), total calorific intake (quintiles), red-meat intake (quintiles), fruit and vegetable intake (quintiles), vitamin D score (deciles), smoking history (never, past quit ≤ 10 yr, past quit > 10 yr, current 1-14 cigarettes per day, 15-24 cigarettes per day, 25 + cigarettes per day), and pack-years (continuous)
Ndegwa <i>et al</i> [30], 2018	Age as time-scale, age at entry, sex, area of residence (rural, small-town or urban), tobacco use status (non-tobacco use, smoking only, snus only or mixed usage), and alcohol consumption (less than once a week <i>versus</i> once a week or more)
Shakeri <i>et al</i> [31], 2013	Age, ethnicity, education, fruit and vegetable use, socioeconomic status, ever opium or tobacco use, and denture use
Watabe <i>et al</i> [32], 1998	NA
Yano <i>et al</i> [33], 2021	Age, sex, socioeconomic score, ethnicity, residence, education, cigarette use, and opium use
Zhang <i>et al</i> [8], 2022	Age (continuous), sex, education (illiteracy, primary school, junior school, high school and above), marital status (single, married, divorced or widowed), job type (farmer, worker, others), wealth score (five levels), BMI 10 years ago (< 18.5 kg/m ² , 18.5 to 24.0 kg/m ² , 24.0 to 28.0 kg/m ² , ≥ 28.0 kg/m ²), tobacco smoking (never, ≤ 30 pack-years, > 30 pack-years), alcohol drinking (never, ≤ 80 g/d, > 80 g/d), <i>H. pylori</i> seropositivity (yes/no), and family history of GC (yes/no)
Zhang <i>et al</i> [34], 2022	Age (continuous), sex (male, female), BMI (continuous), study sites (10 sites), education level (no formal school, primary or middle school, high school and above), marital status (married, other), household income per year (< 10000, < 10000-19999, < 20000-34999, or < 35000), alcohol consumption (non-drinker, occasional drinker, former drinker, or regular drinker), smoking status (never smoker, occasional smoker, former smoker, or regular smoker), physical activity in MET hours a day (continuous), aspirin prescription for CVD (no, yes, or missing), menopausal status (pre-menopausal or post-menopausal, women only), personal history of diabetes (no, yes), and family history of cancer (no, yes)

BMI: Body mass index; MET: Metabolic equivalent tasks; CVD: Cardiovascular disease; NA: Not applicable; *H. pylori*: *Helicobacter pylori*; GC: Gastric cancer.

patients with periodontitis had a greater risk of GC. However, tooth loss, gingivitis, dentures, and tooth brushing had no effect on the risk of GC.

In recent years, a growing number of researchers have focused on the relationship between oral health and cancer[27, 28,35,36]. Periodontitis, a common disease that affects oral health, is a chronic inflammatory disease caused by bacteria that carry the risk of supporting tissue breakdown and tooth loss[37]. Moreover, periodontitis was reported to be a predictive factor for GC[19]. With the increase in the number of GC patients in China[6], the association between oral health and the risk of GC needs more attention in the future.

However, previous studies on oral health and the incidence of GC have been controversial. Several studies reported that tooth loss was not a predictive factor for increased risk of GC[19,29,30,32]. In contrast, some studies have reported that tooth loss could increase the risk of GC[27,33]. Previous studies have shown that periodontitis increases the risk of GC[19]. However, Hujoel *et al*[20] and Michaud *et al*[29] showed that there was no association between periodontitis and the risk of GC. Therefore, it was necessary to explore the real association between oral health and the risk of GC.

In our study, we found that poor oral health, especially periodontitis, was associated with a greater risk of GC. However, the mechanism by which poor oral health increases the risk of GC is unclear. There are several hypotheses that might explain the association between oral health and the risk of GC. First, the oral cavity provides passage between the external environment and the gastrointestinal tract, which is involved in the intake and digestion of food. Oral hygiene might affect the gastrointestinal flora and nutritional status, therefore resulting in the development of chronic diseases [19]. Second, periodontal disease and poor oral hygiene could lead to tooth loss[38]. However, in our study, we found that there was no significant difference between tooth loss and the risk of GC. Tooth loss is often accompanied by chronic infection and inflammation, such as periodontitis[39]. Moreover, tooth loss leads to a decrease in the ability to chew and might alter the patient's eating patterns[40-42]. These inflammatory conditions and changes in dietary habits associated with tooth loss might be the cause of the increased risk of GC. Third, patients with periodontal disease and poor oral hygiene had significantly greater levels of oral bacteria, while nitrosamine levels were significantly greater in the oral cavity due to the presence of nitrate-reducing bacteria; moreover, it is widely known that nitrites are recognized carcinogens[43]. Tooth brushing could also affect oral health; however, we did not find an association between tooth brushing




Figure 3 Subgroups analysis. CI: Confidence interval.

and GC. Shakeri *et al*[31] discussed the relationship between toothbrushing frequency and GC incidence in their study. They found that those who never brushed their teeth had significantly greater rates of GC, while those who brushed their teeth every day or less than daily had no significant change in their rates of GC. In our study, we explored the effect of toothbrushing on the incidence of GC only. This might have contributed to our results.

To the best of our knowledge, the present study was the first to pool the risk of GC in patients who had oral health problems. Our study had a large sample size, and subgroup analysis was conducted. Moreover, the publication bias of the included studies was low. Thus, the outcomes were relatively reliable. This study has several limitations. In our study, there were more cohort studies and fewer case-control studies. Second, because of insufficient data, we lacked information on the effect of different numbers of missing teeth on the incidence of GC. Therefore, further case-control studies need to be performed in the future.




Figure 4 Funnel plot. CI: Confidence interval.

CONCLUSION

In conclusion, patients with poor oral health, especially those with periodontitis, had a higher risk of GC. Thus, patients should be concerned about their oral health. Improving oral health might reduce the risk of GC.

ARTICLE HIGHLIGHTS

Research background

Gastric cancer (GC) is one of the most common tumours worldwide and the forth leading cause of cancer death. Prevention of GC has become a focal point because of these worrisome numbers. Prevention of GC can be divided into primary prevention (reducing the incidence of GC) and secondary prevention (early detection and treatment). Primary prevention includes smoking cessation, reducing salt intake, increasing fruit and vegetable intake, and other health behaviours, such as oral health behaviours.

Research motivation

The aim of present study is to assess whether there is a relationship between oral health and the risk of GC.

Research objectives

The research objective was to explore the relationship between oral health and GC risk.

Research methods

This study searched five databases to find eligible studies from inception to April 10, 2023. Newcastle-Ottawa Scale score was used to assess the quality of included studies. The quality of cohort studies and case-control studies were evaluated separately in this study. Incidence of GC were described by odds ratio (OR) and 95% confidence interval (CI). Funnel plot was used to represent the publication bias of included studies. We performed the data analysis by StataSE 16.

Research results

A total of 1431677 patients from twelve included studies were enrolled for data analysis in this study. According to our analysis, we found that poor oral health was associated with a high risk of GC (OR = 1.15, 95%CI: 1.02-1.29; $I^2 = 59.47\%$, $P = 0.00 < 0.01$), particularly periodontitis (OR = 1.13, 95%CI: 1.04-1.23; $I^2 = 0.00\%$, $P < 0.01$). Moreover, after subgroup analysis, tooth loss (OR = 1.12, 95%CI: 0.94-1.29; $I^2 = 6.01\%$, $P > 0.01$), gingivitis (OR = 1.19, 95%CI: 0.71-1.67; $I^2 = 0.00\%$, $P > 0.01$), dentures (OR = 1.27, 95%CI: 0.63-1.19; $I^2 = 68.79\%$, $P > 0.01$), or tooth brushing (OR = 1.25, 95%CI: 0.78-1.71; $I^2 = 88.87\%$, $P > 0.01$) had no influence on the risk of GC.

Research conclusions

Oral health status associated with GC risk. People should focus on oral health as it might reduce the incidence of GC.

Research perspectives

This study was extended to a multi-center study.

FOOTNOTES

Co-first authors: Fei Liu and Shi-Jun Tang.

Author contributions: Liu F and Tang SJ contributed equally to this work. All authors contributed to data collection and analysis, drafting, or revising the manuscript, have agreed on the journal to which the manuscript will be submitted, gave final approval of the version to be published, and agree to be accountable for all aspects of the work.

Conflict-of-interest statement: All the authors report no relevant conflicts of interest for this article.

PRISMA 2009 Checklist statement: The authors have read the PRISMA 2009 Checklist, and the manuscript was prepared and revised according to the PRISMA 2009 Checklist.

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <https://creativecommons.org/licenses/by-nc/4.0/>

Country/Territory of origin: China

ORCID number: Fei Liu 0000-0002-4022-0732; Shi-Jun Tang 0000-0003-0136-3658; Zi-Wei Li 0000-0001-9759-4535; Xu-Rui Liu 0000-0002-6069-2104; Quan Lv 0009-0005-8861-0181; Wei Zhang 0000-0002-5822-9970; Dong Peng 0000-0003-4050-4337.

S-Editor: Wang JJ

L-Editor: A

P-Editor: ZhangYL

REFERENCES

- 1 Sung H, Ferlay J, Siegel RL, Laversanne M, Soerjomataram I, Jemal A, Bray F. Global Cancer Statistics 2020: GLOBOCAN Estimates of Incidence and Mortality Worldwide for 36 Cancers in 185 Countries. *CA Cancer J Clin* 2021; **71**: 209-249 [PMID: 33538338 DOI: 10.3322/caac.21660]
- 2 Cheng YX, Tao W, Liu XY, Zhang H, Yuan C, Zhang B, Zhang W, Peng D. Does Chronic Kidney Disease Affect the Surgical Outcome and Prognosis of Patients with Gastric Cancer? A Meta-Analysis. *Nutr Cancer* 2022; **74**: 2059-2066 [PMID: 34693829 DOI: 10.1080/01635581.2021.1993277]
- 3 Bray F, Ferlay J, Soerjomataram I, Siegel RL, Torre LA, Jemal A. Global cancer statistics 2018: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. *CA Cancer J Clin* 2018; **68**: 394-424 [PMID: 30207593 DOI: 10.3322/caac.21492]
- 4 Peng D, Zou YY, Cheng YX, Tao W, Zhang W. Effect of Time (Season, Surgical Starting Time, Waiting Time) on Patients with Gastric Cancer. *Risk Manag Healthc Policy* 2021; **14**: 1327-1333 [PMID: 33824610 DOI: 10.2147/RMHP.S294141]
- 5 Ferlay J, Soerjomataram I, Dikshit R, Eser S, Mathers C, Rebelo M, Parkin DM, Forman D, Bray F. Cancer incidence and mortality worldwide: sources, methods and major patterns in GLOBOCAN 2012. *Int J Cancer* 2015; **136**: E359-E386 [PMID: 25220842 DOI: 10.1002/ijc.29210]
- 6 Kang B, Liu XY, Cheng YX, Tao W, Peng D. Factors associated with hypertension remission after gastrectomy for gastric cancer patients. *World J Gastrointest Surg* 2022; **14**: 743-753 [PMID: 36157372 DOI: 10.4240/wjgs.v14.i8.743]
- 7 Chen W, Zheng R, Baade PD, Zhang S, Zeng H, Bray F, Jemal A, Yu XQ, He J. Cancer statistics in China, 2015. *CA Cancer J Clin* 2016; **66**: 115-132 [PMID: 26808342 DOI: 10.3322/caac.21338]
- 8 Zhang T, Yang X, Yin X, Yuan Z, Chen H, Jin L, Chen X, Lu M, Ye W. Poor oral hygiene behavior is associated with an increased risk of gastric cancer: A population-based case-control study in China. *J Periodontol* 2022; **93**: 988-1002 [PMID: 34599526 DOI: 10.1002/JPER.21-0301]
- 9 Karimi P, Islami F, Anandasabapathy S, Freedman ND, Kamangar F. Gastric cancer: descriptive epidemiology, risk factors, screening, and prevention. *Cancer Epidemiol Biomarkers Prev* 2014; **23**: 700-713 [PMID: 24618998 DOI: 10.1158/1055-9965.EPI-13-1057]
- 10 Tang X, Zhang M, He Q, Sun G, Wang C, Gao P, Qu H. Histological Differentiated/Undifferentiated Mixed Type Should Not Be Considered as a Non-Curative Factor of Endoscopic Resection for Patients With Early Gastric Cancer. *Front Oncol* 2020; **10**: 1743 [PMID: 33014861 DOI: 10.3389/fonc.2020.01743]
- 11 Chelimo C, Elwood JM. Sociodemographic differences in the incidence of oropharyngeal and oral cavity squamous cell cancers in New Zealand. *Aust N Z J Public Health* 2015; **39**: 162-167 [PMID: 25827186 DOI: 10.1111/1753-6405.12352]
- 12 Eksheden I, Yang X, Chen H, Chen X, Yuan Z, Jin L, Lu M, Ye W. Associations Between Gastric Atrophy and Its Interaction With Poor Oral Health and the Risk for Esophageal Squamous Cell Carcinoma in a High-Risk Region of China: A Population-Based Case-Control Study. *Am J Epidemiol* 2020; **189**: 931-941 [PMID: 31899792 DOI: 10.1093/aje/kwz283]
- 13 Printz C. African American women with gum disease and tooth loss face higher pancreatic cancer risk. *Cancer* 2019; **125**: 2719 [PMID: 31355935 DOI: 10.1002/ncr.32413]
- 14 Pizzo G, Guiglia R, Lo Russo L, Campisi G. Dentistry and internal medicine: from the focal infection theory to the periodontal medicine concept. *Eur J Intern Med* 2010; **21**: 496-502 [PMID: 21111933 DOI: 10.1016/j.ejim.2010.07.011]
- 15 Kekschull M, Demmer RT, Papapanou PN. "Gum bug, leave my heart alone!"--epidemiologic and mechanistic evidence linking periodontal infections and atherosclerosis. *J Dent Res* 2010; **89**: 879-902 [PMID: 20639510 DOI: 10.1177/0022034510375281]
- 16 Wada T, Kunisaki C, Ono HA, Makino H, Akiyama H, Endo I. Implications of BMI for the Prognosis of Gastric Cancer among the Japanese Population. *Dig Surg* 2015; **32**: 480-486 [PMID: 26529523 DOI: 10.1159/000440654]

- 17 **Gao Z**, Ni J, Ding H, Yan C, Ren C, Li G, Pan F, Jin G. A nomogram for prediction of stage III/IV gastric cancer outcome after surgery: A multicenter population-based study. *Cancer Med* 2020; **9**: 5490-5499 [PMID: 32543092 DOI: 10.1002/cam4.3215]
- 18 **Pietiläinen M**, Liljestrand JM, Kopra E, Pussinen PJ. Mediators between oral dysbiosis and cardiovascular diseases. *Eur J Oral Sci* 2018; **126** Suppl 1: 26-36 [PMID: 30178551 DOI: 10.1111/eos.12423]
- 19 **Hiraki A**, Matsuo K, Suzuki T, Kawase T, Tajima K. Teeth loss and risk of cancer at 14 common sites in Japanese. *Cancer Epidemiol Biomarkers Prev* 2008; **17**: 1222-1227 [PMID: 18483345 DOI: 10.1158/1055-9965.EPI-07-2761]
- 20 **Hujoel PP**, Drangsholt M, Spiekerman C, Weiss NS. An exploration of the periodontitis-cancer association. *Ann Epidemiol* 2003; **13**: 312-316 [PMID: 12821269 DOI: 10.1016/s1047-2797(02)00425-8]
- 21 **Kim EH**, Nam S, Park CH, Kim Y, Lee M, Ahn JB, Shin SJ, Park YR, Jung HI, Kim BI, Jung I, Kim HS. Periodontal disease and cancer risk: A nationwide population-based cohort study. *Front Oncol* 2022; **12**: 901098 [PMID: 36081548 DOI: 10.3389/fonc.2022.901098]
- 22 **Page MJ**, McKenzie JE, Bossuyt PM, Boutron I, Hoffmann TC, Mulrow CD, Shamseer L, Tetzlaff JM, Akl EA, Brennan SE, Chou R, Glanville J, Grimshaw JM, Hróbjartsson A, Lalu MM, Li T, Loder EW, Mayo-Wilson E, McDonald S, McGuinness LA, Stewart LA, Thomas J, Tricco AC, Welch VA, Whiting P, Moher D. The PRISMA 2020 statement: An updated guideline for reporting systematic reviews. *J Clin Epidemiol* 2021; **134**: 178-189 [PMID: 33789819 DOI: 10.1016/j.jclinepi.2021.03.001]
- 23 **Stang A**. Critical evaluation of the Newcastle-Ottawa scale for the assessment of the quality of nonrandomized studies in meta-analyses. *Eur J Epidemiol* 2010; **25**: 603-605 [PMID: 20652370 DOI: 10.1007/s10654-010-9491-z]
- 24 **Zhang J**, Yu KF. What's the relative risk? A method of correcting the odds ratio in cohort studies of common outcomes. *JAMA* 1998; **280**: 1690-1691 [PMID: 9832001 DOI: 10.1001/jama.280.19.1690]
- 25 **Ioannidis JP**. Interpretation of tests of heterogeneity and bias in meta-analysis. *J Eval Clin Pract* 2008; **14**: 951-957 [PMID: 19018930 DOI: 10.1111/j.1365-2753.2008.00986.x]
- 26 **Higgins JP**, Thompson SG, Deeks JJ, Altman DG. Measuring inconsistency in meta-analyses. *BMJ* 2003; **327**: 557-560 [PMID: 12958120 DOI: 10.1136/bmj.327.7414.557]
- 27 **Abnet CC**, Qiao YL, Mark SD, Dong ZW, Taylor PR, Dawsey SM. Prospective study of tooth loss and incident esophageal and gastric cancers in China. *Cancer Causes Control* 2001; **12**: 847-854 [PMID: 11714113 DOI: 10.1023/a:1012290009545]
- 28 **Abnet CC**, Kamangar F, Dawsey SM, Stolzenberg-Solomon RZ, Albanes D, Pietinen P, Virtamo J, Taylor PR. Tooth loss is associated with increased risk of gastric non-cardia adenocarcinoma in a cohort of Finnish smokers. *Scand J Gastroenterol* 2005; **40**: 681-687 [PMID: 16036528 DOI: 10.1080/00365520510015430]
- 29 **Michaud DS**, Liu Y, Meyer M, Giovannucci E, Joshipura K. Periodontal disease, tooth loss, and cancer risk in male health professionals: a prospective cohort study. *Lancet Oncol* 2008; **9**: 550-558 [PMID: 18462995 DOI: 10.1016/S1470-2045(08)70106-2]
- 30 **Ndegwa N**, Ploner A, Liu Z, Roosaar A, Axéll T, Ye W. Association between poor oral health and gastric cancer: A prospective cohort study. *Int J Cancer* 2018; **143**: 2281-2288 [PMID: 29873081 DOI: 10.1002/ijc.31614]
- 31 **Shakeri R**, Malekzadeh R, Etemadi A, Nasrollahzadeh D, Abedi-Ardekani B, Khoshnia M, Islami F, Pourshams A, Pawlita M, Boffetta P, Dawsey SM, Kamangar F, Abnet CC. Association of tooth loss and oral hygiene with risk of gastric adenocarcinoma. *Cancer Prev Res (Phila)* 2013; **6**: 477-482 [PMID: 23503651 DOI: 10.1158/1940-6207.CAPR-12-0491]
- 32 **Watabe K**, Nishi M, Miyake H, Hirata K. Lifestyle and gastric cancer: a case-control study. *Oncol Rep* 1998; **5**: 1191-1194 [PMID: 9683833 DOI: 10.3892/or.5.5.1191]
- 33 **Yano Y**, Abnet CC, Poustchi H, Roshandel G, Pourshams A, Islami F, Khoshnia M, Amirani T, Norouzi A, Kamangar F, Boffetta P, Brennan P, Dawsey SM, Vogtman E, Malekzadeh R, Etemadi A. Oral Health and Risk of Upper Gastrointestinal Cancers in a Large Prospective Study from a High-risk Region: Golestan Cohort Study. *Cancer Prev Res (Phila)* 2021; **14**: 709-718 [PMID: 33731409 DOI: 10.1158/1940-6207.CAPR-20-0577]
- 34 **Zhang X**, Liu B, Lynn HS, Chen K, Dai H. Poor oral health and risks of total and site-specific cancers in China: A prospective cohort study of 0.5 million adults. *EClinicalMedicine* 2022; **45**: 101330 [PMID: 35274091 DOI: 10.1016/j.eclinm.2022.101330]
- 35 **Mello FW**, Melo G, Pasetto JJ, Silva CAB, Warnakulasuriya S, Rivero ERC. The synergistic effect of tobacco and alcohol consumption on oral squamous cell carcinoma: a systematic review and meta-analysis. *Clin Oral Investig* 2019; **23**: 2849-2859 [PMID: 3111280 DOI: 10.1007/s00784-019-02958-1]
- 36 **Wang YP**, Han XY, Su W, Wang YL, Zhu YW, Sasaba T, Nakachi K, Hoshiyama Y, Tagashira Y. Esophageal cancer in Shanxi Province, People's Republic of China: a case-control study in high and moderate risk areas. *Cancer Causes Control* 1992; **3**: 107-113 [PMID: 1562700 DOI: 10.1007/BF00051650]
- 37 **Fu MM**, Chien WC, Chung CH, Lee WC, Tu HP, Fu E. Is periodontitis a risk factor of benign or malignant colorectal tumor? A population-based cohort study. *J Periodontol Res* 2022; **57**: 284-293 [PMID: 34854493 DOI: 10.1111/jre.12955]
- 38 **Shimazaki Y**, Soh I, Koga T, Miyazaki H, Takehara T. Risk factors for tooth loss in the institutionalised elderly; a six-year cohort study. *Community Dent Health* 2003; **20**: 123-127 [PMID: 12828274 DOI: 10.1111/j.1600-0528.2011.00648.x]
- 39 **Coussens LM**, Werb Z. Inflammation and cancer. *Nature* 2002; **420**: 860-867 [PMID: 12490959 DOI: 10.1038/nature01322]
- 40 **Kim S**, Doh RM, Yoo L, Jeong SA, Jung BY. Assessment of Age-Related Changes on Masticatory Function in a Population with Normal Dentition. *Int J Environ Res Public Health* 2021; **18** [PMID: 34199065 DOI: 10.3390/ijerph18136899]
- 41 **Feldman RS**, Kapur KK, Alman JE, Chauncey HH. Aging and mastication: changes in performance and in the swallowing threshold with natural dentition. *J Am Geriatr Soc* 1980; **28**: 97-103 [PMID: 7354209 DOI: 10.1111/j.1532-5415.1980.tb00240.x]
- 42 **Batisse C**, Bonnet G, Eschevins C, Hennequin M, Nicolas E. The influence of oral health on patients' food perception: a systematic review. *J Oral Rehabil* 2017; **44**: 996-1003 [PMID: 28600840 DOI: 10.1111/joor.12535]
- 43 **Nair J**, Ohshima H, Nair UJ, Bartsch H. Endogenous formation of nitrosamines and oxidative DNA-damaging agents in tobacco users. *Crit Rev Toxicol* 1996; **26**: 149-161 [PMID: 8688158 DOI: 10.3109/10408449609017928]



Published by **Baishideng Publishing Group Inc**
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA

Telephone: +1-925-3991568

E-mail: office@baishideng.com

Help Desk: <https://www.f6publishing.com/helpdesk>

<https://www.wjgnet.com>

