



Retrospective Study

Comparison of mismatch repair and immune checkpoint protein profile with histopathological parameters in pancreatic, periampullary/ampullary, and choledochal adenocarcinomas

Arzu Hazal Aydin, Nesrin Turhan

Specialty type: Oncology

Provenance and peer review:

Unsolicited article; Externally peer reviewed.

Peer-review model: Single blind

Peer-review report's scientific quality classification

Grade A (Excellent): A

Grade B (Very good): 0

Grade C (Good): C

Grade D (Fair): 0

Grade E (Poor): 0

P-Reviewer: Haddadi S, Algeria; Mei LC, China

Received: November 3, 2023

Peer-review started: November 3, 2023

First decision: December 1, 2023

Revised: December 12, 2023

Accepted: January 10, 2024

Article in press: January 10, 2024

Published online: March 15, 2024



Arzu Hazal Aydin, Department of Pathology, Aksaray University Aksaray Training and Research Hospital, Aksaray 68200, Turkey

Nesrin Turhan, Department of Pathology, University of Health Sciences Ankara City Hospital, Ankara 06800, Turkey

Corresponding author: Arzu Hazal Aydin, MD, Doctor, Department of Pathology, Aksaray University Aksaray Training and Research Hospital, Sifahane District, No. 3152 Street, Aksaray 68200, Turkey. arzu.hazal.aydin@gmail.com

Abstract

BACKGROUND

Pancreatic, periampullary/ampullary, and choledochal adenocarcinomas are aggressive malignancies with a poor prognosis. Immune checkpoint blockade is a promising treatment option for several tumor types. H long terminal repeat-associated 2 (HHLA2), which is analogous to programmed death-ligand 1 (PD-L1), is a recently discovered member of the B7/cluster of differentiation 28 family and is expressed in many malignancies.

AIM

To analyze the expression of HHLA2 and its association with the pathologic biomarkers that predict sensitivity to immunotherapy.

METHODS

Ninety-two adenocarcinoma cases located in the pancreas, ampulla, and distal common bile duct were identified. This study assessed 106 pancreaticoduodenectomy and distal/total pancreatectomy samples that were delivered to Ankara City Hospital between 2019 and 2021. Immunohistochemistry was conducted to examine the expression of DNA mismatch repair (MMR), PD-L1, and HHLA2 proteins.

RESULTS

Patients with high HHLA2 expression had a higher mean age than those with low expression. Low HHLA2 expression was associated with high perineural invasion. HHLA2 expression was low in pathological stage T3 (pT) 3 cases and high in pathological stage T1, T2, and T4 cases. There was no correlation between HHLA2

expression and the expression of MMR proteins and PD-L1.

CONCLUSION

Evaluation of HHLA2 expression in microsatellite stable and PD-L1-negative tumors may be useful for predicting the response of individuals to immunotherapy and may serve as a novel therapeutic target for immunotherapy in advanced-stage disease.

Key Words: H long terminal repeat-associating 2; Programmed death-ligand 1; Adenocarcinoma; Pancreas; Ampulla of Vater; Distal common bile duct

©The Author(s) 2024. Published by Baishideng Publishing Group Inc. All rights reserved.

Core Tip: Pancreatic, periampullary/ampullary, and choledochal adenocarcinomas are aggressive malignancies with recurrences and metastases occurring in a short period of time. Complete resection is possible in only a small proportion of patients; thus, targeted therapies are needed. H long terminal repeat-associating 2 (HHLA2), which is analogous to programmed death-ligand 1 (PD-L1), is expressed in many malignancies. Evaluation of HHLA2 expression in microsatellite stable and PD-L1-negative tumors may be useful for predicting the response of individuals to immunotherapy, and may serve as a novel therapeutic target in patients with advanced disease who do not respond to classical chemotherapy and have unresectable cancer.

Citation: Aydın AH, Turhan N. Comparison of mismatch repair and immune checkpoint protein profile with histopathological parameters in pancreatic, periampullary/ampullary, and choledochal adenocarcinomas. *World J Gastrointest Oncol* 2024; 16(3): 875-882

URL: <https://www.wjgnet.com/1948-5204/full/v16/i3/875.htm>

DOI: <https://dx.doi.org/10.4251/wjgo.v16.i3.875>

INTRODUCTION

Pancreatic, ampullary, and distal common bile duct (DCBD) carcinomas are characterized by late presentation, advanced disease at the time of diagnosis, and the inability to completely resect because of the location. While the 5-year disease-specific survival (DSS) is 42% in resectable DCBD carcinomas, in ampullary adenocarcinoma (AAC), the DSS rate ranges from 20% to 80% according to the histological type, location, and stage[1]. In pancreatic ductal adenocarcinoma (PDAC), the 5-year survival rate is much lower, between 2% and 9%[1]. Complete resection is possible in only a small proportion of patients, and recurrences and metastases occur in a short period of time; thus, targeted therapies are needed for this disease[2].

Immunotherapy has been used to treat patients with advanced stage disease or unresectable tumor to improve cancer survival. However, most patients do not respond to immunotherapy; hence, there is a need for specific biomarkers to improve the selection of patients who will best respond to therapy.

Mismatch repair (MMR) proteins play an important role in the detection and correction of errors that occur during DNA replication and genetic recombination. MMR deficiency can lead to microsatellite instability (MSI). Consequently, the accumulation of tumor mutational burden and generation of neoantigens stimulate the host immune response. MSI-high (MSI-H) solid tumors are more sensitive to immunotherapy agents and associated with a better prognosis[3].

Immunogenic proteins, also called immune checkpoint proteins, are produced by tumor cells. Programmed cell death protein 1 (PD-1) is an immune checkpoint protein that promotes immune evasion and tumor progression *via* interaction with its ligands [programmed death-ligand (PD-L)1 and PD-L2]. Immune checkpoint inhibitors (iCPIs) allow T cells to destroy tumor cells by blocking this interaction[4].

Clinical success with monoclonal antibodies such as pembrolizumab, which blocks the PD-1/PD-L1 axis, has opened new doors in cancer immunology. Pembrolizumab is approved for the treatment of patients with unresectable or metastatic MSI-H or defective MMR (dMMR) solid tumors. The success of pembrolizumab in treating dMMR/MSI-H tumors has elicited excitement and encouraged the exploration of new discoveries in immunotherapy[4].

PD-L1 is expressed in many tumor types. The relationship between cancer and the immune system is not fully elucidated, and it is unclear if there is an association between histopathological features and the response to iCPIs. The success of PD-1/PD-L1 blocking therapies has led to the identification of other members of this pathway as targets for immunotherapy and the use of PD-1/PD-L1 expression as a prognostic factor[2].

Despite the promising antitumor activity of PD-1/PD-L1 inhibitors, only a small group of patients show marked responses to single-agent antibody therapy. The need to elucidate the mechanism underlying the resistance to immune checkpoint blockade in some patients has been the focus of many studies[3,5-7].

The success of PD-1/PD-L1 blockade in cancer immunotherapy has raised the possibility of other members of the B7/cluster of differentiation 28 (CD28) family serving as targets for cancer immunotherapy. H long terminal repeat-associating 2 (HHLA2), which is a recently discovered member of the B7/CD28 family, is analogous to PD-L1 and regulates T

cell functions[8-10]. In addition, HHLA2 has been shown to be expressed in both nonneoplastic tissue and many malignancies[9]. In limited studies, the high expression of HHLA2 in PDAC and periampullary adenocarcinoma has been associated with the better overall survival (OS) of patients[9,11]; however, there is a need for more studies on this subject.

Herein, we describe our assessments of the association of histological and clinical parameters [age, sex, histological type, histological grade, tumor-node-metastasis (TNM) stage, presence of tumor infiltrating lymphocytes (TILs), tumor stroma ratio (TSR), tumor budding (TB)] with the expression of MMR, PD-L1, and HHLA2 proteins in patients diagnosed with adenocarcinoma located in the pancreas, periampulla/ampulla, and DCBD.

MATERIALS AND METHODS

Study design

Ninety-two cases of adenocarcinoma located in the pancreas, periampulla/ampulla, and DCBD were identified. This study assessed pancreaticoduodenectomy and distal/total pancreatectomy samples that were delivered to Ankara City Hospital (Ankara, Turkey) between 2019 and 2021. All protocols were reviewed and approved by the Ethics Committee of Ankara City Hospital.

Demographic information and pathology reports were obtained from the hospital database (HICAMP). Information on sex, age, tumor macroscopic location, presence/absence of lymphovascular invasion (LVI)/perineural invasion (PNI), and lymph node metastasis were obtained from the pathology reports.

All tumor sections stained with hematoxylin and eosin (H&E) were removed from the slide archive and examined under a microscope by two observers. The parameters related to the histological type and grade of the cases were reevaluated based on the 5th edition of the World Health Organization Digestive System Tumor Classification published in 2019 and the 8th edition of the American Joint Committee on Cancer staging system based on the TNM concept.

Assessment of TB, TILs, and the TSR

To evaluate and grade TB, a three-tier system proposed at the 2016 International Tumor Budding Consensus Conference was used. TILs were evaluated separately as stromal and intraepithelial TILs. To evaluate stromal TILs, standardized criteria prepared in 2014 by the International TIL Study Group were used. At a low magnification, tumor slides were scanned and focused on TS to determine the type of inflammatory infiltrate. Mononuclear infiltrate was accepted as stromal TIL, and three groups were formed based on their densities. Stromal TIL rate of 0%-10% was included in group A, 10%-40% was included in group B, and 40%-90% was included in group C. The number of intraepithelial lymphocytes (IELs) were counted and averaged over five high-power fields for each case; two groups were created: < 2 and ≥ 2 IELs. For TSR evaluation, the most invasive area of the tumor was selected among the H&E-stained slides. Then, the slides were digitized with a 40 × objective using the Aperio AT Turbo digital whole slide scanning system (Leica Biosystems Imaging, Vista, CA, United States). The preparations were evaluated using the semi-quantitative image analysis program Virapath-3.4.4 (Virasoft Software, Istanbul, Turkey). Tissue presegmentation analysis was run in the hotspot regions corresponding to the 10 × magnification area in each evaluated case. The tumoral region area was calculated and the ratio was calculated by dividing it by the area of the stroma regions. Two groups (low and high) were formed using the 50% cutoff value according to the literature[12].

Immunohistochemistry

Immunohistochemistry was conducted with mouse monoclonal antibodies against the four main MMR proteins: MLH-1 (clone M1), MSH-2 (clone G219-1129), MSH-6 (clone SP93), and PMS-2 (clone A16-4); a mouse monoclonal antibody against PD-L1 (clone 22C3; Dako Products, Santa Clara, CA, United States); and a rabbit polyclonal antibody against HHLA2 (clone ab214327). For MMR and PD-L1, sections were prepared from formalin-fixed paraffin blocks; for HHLA2, sections were prepared from tissue microarray blocks.

Immunohistochemistry staining of MMR proteins was evaluated for the presence of nuclear staining in the tumor cells. Stromal cells and lymphocytes adjacent to each tumor served as internal controls. In the presence of an optimal internal control, nuclear staining in tumor cells was considered preserved expression. Tumor cells with membranous staining of PD-L1 were considered positive. At least 100 tumor cells were evaluated for each case, and the number of PD-L1-positive tumor cells was divided by the total number of tumor cells. A rate < 1% was considered negative, rates from 1% to 49% were classified as low, and a rate ≥ 50% was classified as high (Figure 1). HHLA2 cytoplasmic staining and the ratio of stained cells were determined for each case. Tumors with no staining were considered negative. The cases with staining were grouped as low and high expression according to the 50% cutoff value (Figure 2).

Statistical analyses

Statistical analyses were performed using SPSS version 17.0 (IBM, Armonk, NY, United States). The conformity of the variables to a normal distribution was examined by histogram graphs and the Kolmogorov-Smirnov test. Mean, standard deviation, and median values were used to present the descriptive analyses. Categorical variables were compared with Pearson's χ^2 test. In cases where the data were not normally distributed, groups of two were evaluated with the Mann-Whitney *U* test and groups of more than two were evaluated with the Kruskal-Wallis test. Spearman's correlation test was used to analyze the measurement data. *P* < 0.05 was considered statistically significant.




Figure 1 Membranous staining of tumor cells expressing programmed death-ligand 1. All panels show programmed death-ligand 1 expression in tumor cells from different cases (20 ×).




Figure 2 Cytoplasmic expression of H long terminal repeat-associating 2 in tumor cells. Examples of different cases stained for H long terminal repeat-associating 2.

RESULTS

In our study, the mean patient age was 65.51 years, and 64 (69.57%) cases were male and 28 (30.43%) were female. Fifty-eight of the ninety-two cases (63.04%) were located in the pancreas. The location of the other cases was as follows: 21 (22.83%) in the ampulla and 11 (11.96%) in the DCBD. According to histological subtype, 81 (87.79%) cases were ductal adenocarcinoma (DAC), 4 (4.44%) were intraductal papillary mucinous neoplasm-associated invasive carcinoma, 3 (3.33%) were intraductal tubulopapillary neoplasm-associated invasive carcinoma, 3 (3.33%) were classified as adeno-squamous carcinoma, and 1 (1.11%) as signet ring cell carcinoma. In total, 29 of these cases were pathological stage T3 (pT3), 20 were pathological stage T2 (pT2), 2 were pathological stage T4 (pT4), and 1 was pathological stage T1 (pT1). TB was lower in the well-differentiated cases than in the moderately and poorly differentiated cases. There was more LVI and PNI in cases with high TB.

In our study, the cases were classified as < 2 and ≥ 2 according to the presence of IELs. A significant correlation was found between stromal TIL groups and IEL density. It was determined that 52 of the 87 cases with IELs < 2 showed the presence of group B stromal TILs. The rate of group C stromal TILs in cases with IELs ≥ 2 was found to be higher than the rate of group A and B stromal TILs. The cases were divided into two groups (low and high) according to the TSR. The TSR was not significantly associated with histopathological features, TB, or stromal TIL groups. HHLA2 expression was compared with PD-L1 and MMR protein expression. No meaningful correlation was found between the MMR protein expression profile and the expression of PD-L1 and HHLA2. In addition, no significant relationship was found between

the absence/presence of HHLA2 expression and TB, stromal TIL groups, IEL density, or TSR. Patients with high HHLA2 expression had a higher mean age than those with low expression. Low HHLA2 expression was associated with high PNI. HHLA2 expression was low in pT3 cases and high in pT1, pT2, and pT4 cases (Table 1).

DISCUSSION

Many studies have revealed the relationship between high-grade TB and aggressive histopathologic features (LVI, advanced pT stage) in PDAC[13,14]. Fewer studies have evaluated TB in ampullary and DCBD carcinomas. For both tumor types, high-grade TB was determined to be associated with features that reflect the aggressive tumor characteristics including poor histological differentiation, advanced stage, LVI and PNI, and lymph node metastasis[9,11].

Corresponding with the studies in the literature, the TB was lower in well-differentiated tumors than in moderately and poorly differentiated tumors in our study. TB was found to be significantly higher in tumors with LVI and PNI. The aggressive nature of tumors is a multifactorial process that is considerably affected by the tumor microenvironment (TME). The stroma, which is the main component of the TME, promotes the growth, invasion, and metastasis of tumors[12].

A low TSR has been shown to be associated with poor prognostic parameters such as increased tumor size and advanced tumor stage, as well as decreased OS and disease-free survival[12]. However, no study has investigated the relationship between the TSR and histopathologic features for adenocarcinomas located in the ampulla and DCBD. Our study did not find a significant correlation between the TSR and histopathologic features of the adenocarcinomas examined, possibly because the TSR was evaluated in the densest area; thus, the fact that the entire tumor area was not included in the evaluation may have affected the result. TILs and secreted cytokines are major components of the TME and play important roles in the immune regulation of cancer[15]. TIL density has come to the forefront as a predictive biomarker of the immunotherapy response[16].

The distribution of intratumoral, peritumoral, stromal, and intraepithelial TILs have been investigated in relation to antitumoral immune reactions in the TME[15]. There are few studies in the literature comparing the different localization of TILs in tumors located in the pancreas, ampulla, and DCBD. In a study by Zhang *et al*[15], the authors assessed the presence of stromal and intraepithelial TILs in PDAC and investigated the effects on prognosis. A significant T cell response due to intratumoral TILs was observed in PDAC tissues, whereas the response due to intraepithelial attack was very low. It has been reported that CD8+ T intraepithelial infiltration is an independent positive prognostic factor in OS and is negatively correlated with vascular invasion. However, high intratumoral CD8+ T cell infiltration without intraepithelial attack is associated with advanced tumor stage and is considered an adverse prognostic factor[14].

In our study, the relationship between the density of IEL and the stromal TIL groups was analyzed. The rate of having group C stromal TILs in cases with IELs ≥ 2 and the rate of being group A and B in cases with IELs < 2 were found to be significantly higher than in group C with IELs < 2 . It has been suggested that tumor mutation load is related to the response rate to PD-1 inhibitors[3,17]. Assessment of the presence of MSI in tumors is important for predicting the effectiveness of PD-1/PD-L1 blockade[2]. The median number of somatic mutations is higher in MSI-H tumors than in MSS tumors[18]. MSI is found in 1% of PDAC cases, 9.5% of AAC cases, and approximately 4% of DCBD adenocarcinoma cases[2].

In this study, the expression of MMR proteins was preserved in 90 cases, a loss of MSH6 and PMS2 expression was observed in 1 case each; both cases were pancreatic cancer and belonged to the DAC histomorphologic subtype. A significant relationship was not found between MMR protein expression and the histopathological features or immunohistochemical markers assessed in our study. According to the available information in the literature, when the three localizations (pancreas, ampulla, and DCBD) are compared within themselves, MSI is least in pancreatic adenocarcinoma, while the most common in AAC. These results may be explained by the fact that there were more pancreatic carcinoma cases included in our study than AACs.

Satisfaction with immunotherapy outcomes has led to investigations into the relationship between PD-L1 expression and prognosis in many solid tumors. Studies have evaluated the expression of PD-L1 in adenocarcinomas located in the pancreas, ampulla, and DCBD and shown that the high expression of PD-L1 is associated with a poor prognosis in PDAC and AAC[19,20]. However expression of PD-L1 in TILs is associated with increased survival[21]. In our study, the expression of PD-L1 was evaluated only in tumor cells according to Tumor Proportion Score. No significant association was found between PD-L1 expression and histopathologic features. HHLA2, which is analogous to PD-1, is a newly discovered member of the B7/CD28 family and is expressed in many malignancies[9].

In a limited number of studies, it has been proposed that the expression of HHLA2 is independent from the other members of the B7/CD28 family, and its high expression is associated with long-term survival[9,10,22]. Nevertheless, Chen *et al*[22] reported that the high expression of HHLA2 is related to decreased survival. HHLA2 expression has been investigated in many solid tumors[23,24]; however, there has been no study on HHLA2 expression in adenocarcinomas located in the DCBD.

Byers *et al*[11] detected the decreased expression/Loss of expression of HHLA2 in PDAC compared to normal pancreatic ductal cells. Similar to that study, we found higher HHLA2 expression in the nonneoplastic adjacent tissue of tumors including the pancreas, TILs, and stromal cells. In addition, in cases where cytoplasmic staining was not observed in tumor cells, expression was preserved in nonneoplastic tissue.

In our study, patients with high HHLA2 expression had a higher mean age than those with low expression. At the same time, the rate of PNI was lower in cases with high HHLA2 expression. Low HHLA2 expression was correlated with advanced pathological stage. High expression of HHLA2 was observed in pT1, pT2, and pT4 cases, while low expression was found in pT3 cases. Only one pT4 case had high HHLA2 expression, which did not provide sufficient statistics for a

Table 1 Comparison of H long terminal repeat-associating 2 expression and clinicopathological parameters

		Cytoplasmic HHLA2 expression			P value ¹
		Negative, n (%)	Low, n (%)	High, n (%)	
Gender	Male	7 (63.64)	48 (75.00)	9 (52.94)	0.193
	Female	4 (36.36)	16 (25.00)	8 (47.06)	
Pathologic stage of tumor	T1	0 (0.00)	0(.00)	3(17.65)	0.011
	T2	4 (36.36)	23(35.94)	8(47.06)	
	T3	7 (63.64)	38(59.38)	5(29.41)	
	T4	0 (0.00)	3(4.69)	1(5.88)	
Lymphovascular invasion	Present	1 (9.09)	8(12.50)	2(11.76)	0.949
	Absent	10 (90.91)	56(87.50)	15(88.24)	
Perineural invasion	Present	0 (0.00)	2 (3.13)	3 (17.65)	0.044
	Absent	11 (100.00)	62 (96.88)	14 (82.35)	
Histologic grade	Poorly	2 (18.18)	11 (17.19)	3 (17.65)	0.901
	Moderately	7 (63.64)	41 (64.06)	9 (52.94)	
	Well	2 (18.18)	12 (18.75)	5 (29.41)	
Tumor stroma ratio	Low	4 (36.36)	33 (51.56)	9 (52.94)	0.625
	High	7 (63.64)	31 (48.44)	8 (47.06)	
Diameter of tumor (cm) ³		3.11 ± 1.25 (3.00)	3.65 ± 1.24 (4.00)	2.94 ± 1.03 (3.00)	0.088 ²
Tumor budding degree		2.45 ± 0.82 (3.00)	2.48 ± 0.80 (3.00)	1.88 ± 1.05 (2.00)	0.053 ²
Age ⁴		59.91 ± 6.33 (60.00)	65.38 ± 8.28 (66.00)	69.65 ± 8.77 (72.00)	0.005²

¹Chi-Square test.²Kruskal Wallis test.³mean ± SD is given instead of *n*.⁴mean ± SD is given instead of %.

meaningful inference.

In various tumors, expression of HHLA2 may be consequence of a complex process that is affected by many histopathologic features. Additional studies are needed to analyze the expression of HHLA2 in solid tumors and investigate the relationship between HHLA2 expression and survival.

In a limited number of studies, it has been proposed that both the expression of HHLA2 and co-expression of HHLA2 and PD-L1 are related to high density of TILs[22]. However, there have been no studies on the co-expression of HHLA2/PD-L1 in adenocarcinomas located in the pancreas, ampulla, and DCBD. In our study, the co-expression of PD-L1 and HHLA2 and its relationship with the presence of TILs were compared and no significant correlation was found. While no statistically significant correlation was found between the MMR and PD-L1 protein expression profiles and histopathologic features, a significant correlation was found between HHLA2 cytoplasmic expression and age, pT, and the presence of PNI irrespective of other immunophenotypic features.

Our study had some limitations. First, since we designed this research as a retrospective study, the data of some of the patients could not be accessed, so the number of patients included in the study did not reach the desired level. Second, more PDAC cases were included in the study compared to other cancer types and the examples were not homogenous. However, in light of our findings, which need to be supported by larger case studies, HHLA2 may serve as a novel therapeutic target for immunotherapy.

CONCLUSION

The results of this study suggest that HHLA2 expression in MSS and PD-L1-negative tumors may be useful for predicting the response of individuals to immunotherapy, and may serve as a therapeutic target for immunotherapy in patients with advanced-staged disease who do not respond to classical chemotherapy and have unresectable cancer.

ARTICLE HIGHLIGHTS

Research background

Despite advanced techniques in surgical methods and chemotherapy protocols, pancreatic, periampullary/ampullary, and choledochal adenocarcinomas still have high mortality rates. Thus, targeted therapies are needed.

Research motivation

Immunotherapy has opened a new era in cancer treatment. Despite the positive results obtained in treatment, it is necessary to investigate the patient group that does not respond to treatment, and clarify mechanisms and molecules involved in the resistance to immunotherapy. H long terminal repeat-associating 2 (HHLA2) is a novel immune checkpoint molecule. Therefore, the evaluation of HHLA2 expression may be useful in predicting the response to immunotherapy.

Research objectives

To evaluate the relationship of HHLA2 expression with other immunophenotypic markers.

Research methods

The expression of DNA mismatch repair, programmed death-ligand 1, and HHLA2 proteins was examined by immunohistochemistry. All tumor slides stained with hematoxylin and eosin were screened to evaluate other immunophenotypic features such as intraepithelial tumor-infiltrating lymphocytes.

Research results

Low HHLA2 expression was associated with high perineural invasion (PNI). HHLA2 expression was low in pathological stage T3 cases and high in pathological stage T1, T2, and T4 cases. Patients with high HHLA2 expression had a higher mean age than those with low expression.

Research conclusions

We found that HHLA2 expression was correlated with age, pathological stage and the presence of PNI irrespective of other immunophenotypic features. Thus, HHLA2 may be a useful biomarker for predicting the response to immunotherapy.

Research perspectives

In light of our findings, which will be supported by larger case studies, HHLA2 may serve as a novel therapeutic target for immunotherapy in advanced-stage cancer.

FOOTNOTES

Author contributions: Aydın AH and Turhan N participated in the design, execution, and analysis of the article and approved the final version.

Institutional review board statement: Prior to the study, approval was obtained from T.C. Health Sciences University Ankara City Hospital Ethics Committee (Numbered E2-2021-601; Dated 16/06/2021).

Informed consent statement: Since the study was retrospectively designed, informed consent was not obtained from the patients.

Conflict-of-interest statement: The authors have no conflicts of interest to declare.

Data sharing statement: No additional data are available.

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <https://creativecommons.org/licenses/by-nc/4.0/>

Country/Territory of origin: Turkey

ORCID number: Arzu Hazal Aydın 0000-0003-2591-7037; Nesrin Turhan 0000-0001-6566-2695.

S-Editor: Lin C

L-Editor: A

P-Editor: Xu ZH

REFERENCES

- 1 **McGuigan A**, Kelly P, Turkington RC, Jones C, Coleman HG, McCain RS. Pancreatic cancer: A review of clinical diagnosis, epidemiology, treatment and outcomes. *World J Gastroenterol* 2018; **24**: 4846-4861 [PMID: 30487695 DOI: 10.3748/wjg.v24.i43.4846]
- 2 **Qian Y**, Gong Y, Fan Z, Luo G, Huang Q, Deng S, Cheng H, Jin K, Ni Q, Yu X, Liu C. Molecular alterations and targeted therapy in pancreatic ductal adenocarcinoma. *J Hematol Oncol* 2020; **13**: 130 [PMID: 33008426 DOI: 10.1186/s13045-020-00958-3]
- 3 **Zhao P**, Li L, Jiang X, Li Q. Mismatch repair deficiency/microsatellite instability-high as a predictor for anti-PD-1/PD-L1 immunotherapy efficacy. *J Hematol Oncol* 2019; **12**: 54 [PMID: 31151482 DOI: 10.1186/s13045-019-0738-1]
- 4 **Seliger B**. Basis of PD1/PD-L1 Therapies. *J Clin Med* 2019; **8** [PMID: 31817953 DOI: 10.3390/jcm8122168]
- 5 **Liu S**, Gönen M, Stadler ZK, Weiser MR, Hechtman JF, Vakiani E, Wang T, Vyas M, Joneja U, Al-Bayati M, Segal NH, Smith JJ, King S, Guercio S, Ntiamoah P, Markowitz AJ, Zhang L, Cercek A, Garcia-Aguilar J, Saltz LB, Diaz LA, Klimstra DS, Shia J. Cellular localization of PD-L1 expression in mismatch-repair-deficient and proficient colorectal carcinomas. *Mod Pathol* 2019; **32**: 110-121 [PMID: 30166615 DOI: 10.1038/s41379-018-0114-7]
- 6 **Henriksen A**, Dyhl-Polk A, Chen I, Nielsen D. Checkpoint inhibitors in pancreatic cancer. *Cancer Treat Rev* 2019; **78**: 17-30 [PMID: 31325788 DOI: 10.1016/j.ctrv.2019.06.005]
- 7 **Tanigawa M**, Naito Y, Akiba J, Kawahara A, Okabe Y, Ishida Y, Ishikawa H, Hisaka T, Fujita F, Yasunaga M, Shigaki T, Sudo T, Mihara Y, Nakayama M, Kondo R, Kusano H, Shimamatsu K, Okuda K, Akagi Y, Yano H. PD-L1 expression in pancreatic adenosquamous carcinoma: PD-L1 expression is limited to the squamous component. *Pathol Res Pract* 2018; **214**: 2069-2074 [PMID: 30477643 DOI: 10.1016/j.prp.2018.10.006]
- 8 **Wang B**, Ran Z, Liu M, Ou Y. Prognostic Significance of Potential Immune Checkpoint Member HHLA2 in Human Tumors: A Comprehensive Analysis. *Front Immunol* 2019; **10**: 1573 [PMID: 31379814 DOI: 10.3389/fimmu.2019.01573]
- 9 **Yan H**, Qiu W, Koehne de Gonzalez AK, Wei JS, Tu M, Xi CH, Yang YR, Peng YP, Tsai WY, Remotti HE, Miao Y, Su GH. HHLA2 is a novel immune checkpoint protein in pancreatic ductal adenocarcinoma and predicts post-surgical survival. *Cancer Lett* 2019; **442**: 333-340 [PMID: 30447255 DOI: 10.1016/j.canlet.2018.11.007]
- 10 **Boor PPC**, Sideras K, Biermann K, Hosein Aziz M, Levink IJM, Mancham S, Erler NS, Tang X, van Eijck CH, Bruno MJ, Sprengers D, Zang X, Kwekkeboom J. HHLA2 is expressed in pancreatic and ampullary cancers and increased expression is associated with better post-surgical prognosis. *Br J Cancer* 2020; **122**: 1211-1218 [PMID: 32071413 DOI: 10.1038/s41416-020-0755-4]
- 11 **Byers JT**, Paniccia A, Kaplan J, Koenig M, Kahn N, Wilson L, Chen L, Schulick RD, Edil BH, Zhu Y. Expression of the Novel Costimulatory Molecule B7-H5 in Pancreatic Cancer. *Ann Surg Oncol* 2015; **22** Suppl 3: S1574-S1579 [PMID: 25519928 DOI: 10.1245/s10434-014-4293-2]
- 12 **Wu J**, Liang C, Chen M, Su W. Association between tumor-stroma ratio and prognosis in solid tumor patients: a systematic review and meta-analysis. *Oncotarget* 2016; **7**: 68954-68965 [PMID: 27661111 DOI: 10.18632/oncotarget.12135]
- 13 **Karamitopoulou E**, Zlobec I, Born D, Kondi-Pafiti A, Lykoudis P, Mellou A, Gennatas K, Gloor B, Lugli A. Tumour budding is a strong and independent prognostic factor in pancreatic cancer. *Eur J Cancer* 2013; **49**: 1032-1039 [PMID: 23177090 DOI: 10.1016/j.ejca.2012.10.022]
- 14 **Petrova E**, Zielinski V, Bolm L, Schreiber C, Knief J, Thorns C, Bronsert P, Timme-Bronsert S, Bausch D, Perner S, Keck T, Wellner U. Tumor budding as a prognostic factor in pancreatic ductal adenocarcinoma. *Virchows Arch* 2020; **476**: 561-568 [PMID: 31786688 DOI: 10.1007/s00428-019-02719-1]
- 15 **Zhang J**, Wang YF, Wu B, Zhong ZX, Wang KX, Yang LQ, Wang YQ, Li YQ, Gao J, Li ZS. Intraepithelial Attack Rather than Intratumorally Infiltration of CD8+T Lymphocytes is a Favorable Prognostic Indicator in Pancreatic Ductal Adenocarcinoma. *Curr Mol Med* 2017; **17**: 689-698 [PMID: 29521231 DOI: 10.2174/1566524018666180308115705]
- 16 **Santoemma PP**, Powell DJ Jr. Tumor infiltrating lymphocytes in ovarian cancer. *Cancer Biol Ther* 2015; **16**: 807-820 [PMID: 25894333 DOI: 10.1080/15384047.2015.1040960]
- 17 **Morrison AH**, Byrne KT, Vonderheide RH. Immunotherapy and Prevention of Pancreatic Cancer. *Trends Cancer* 2018; **4**: 418-428 [PMID: 29860986 DOI: 10.1016/j.trecan.2018.04.001]
- 18 **Greenman C**, Stephens P, Smith R, Dalgleish GL, Hunter C, Bignell G, Davies H, Teague J, Butler A, Stevens C, Edkins S, O'Meara S, Vastrik I, Schmidt EE, Avis T, Barthorpe S, Bhamra G, Buck G, Choudhury B, Clements J, Cole J, Dicks E, Forbes S, Gray K, Halliday K, Harrison R, Hills K, Hinton J, Jenkinson A, Jones D, Menzies A, Mironenko T, Perry J, Raine K, Richardson D, Shepherd R, Small A, Tofts C, Varian J, Webb T, West S, Widaa S, Yates A, Cahill DP, Louis DN, Goldstraw P, Nicholson AG, Brasseur F, Looijenga L, Weber BL, Chiew YE, DeFazio A, Greaves MF, Green AR, Campbell P, Birney E, Easton DF, Chenevix-Trench G, Tan MH, Khoo SK, Teh BT, Yuen ST, Leung SY, Wooster R, Futreal PA, Stratton MR. Patterns of somatic mutation in human cancer genomes. *Nature* 2007; **446**: 153-158 [PMID: 17344846 DOI: 10.1038/nature05610]
- 19 **Kim MH**, Jang M, Kim H, Lee WJ, Kang CM, Choi HJ. Distinct immunological properties of the two histological subtypes of adenocarcinoma of the ampulla of Vater. *Cancer Immunol Immunother* 2019; **68**: 443-454 [PMID: 30604042 DOI: 10.1007/s00262-018-02293-6]
- 20 **Wang L**, Ma Q, Chen X, Guo K, Li J, Zhang M. Clinical significance of B7-H1 and B7-1 expressions in pancreatic carcinoma. *World J Surg* 2010; **34**: 1059-1065 [PMID: 20145927 DOI: 10.1007/s00268-010-0448-x]
- 21 **Badalamenti G**, Fanale D, Incorvaia L, Barraco N, Listi A, Maragliano R, Vincenzi B, Calò V, Iovanna JL, Bazan V, Russo A. Role of tumor-infiltrating lymphocytes in patients with solid tumors: Can a drop dig a stone? *Cell Immunol* 2019; **343**: 103753 [PMID: 29395859 DOI: 10.1016/j.cellimm.2018.01.013]
- 22 **Chen L**, Zhu D, Feng J, Zhou Y, Wang Q, Feng H, Zhang J, Jiang J. Overexpression of HHLA2 in human clear cell renal cell carcinoma is significantly associated with poor survival of the patients. *Cancer Cell Int* 2019; **19**: 101 [PMID: 31015801 DOI: 10.1186/s12935-019-0813-2]
- 23 **Xiao Y**, Freeman GJ. A New B7:CD28 Family Checkpoint Target for Cancer Immunotherapy: HHLA2. *Clin Cancer Res* 2015; **21**: 2201-2203 [PMID: 25869386 DOI: 10.1158/1078-0432.CCR-14-2658]
- 24 **Zhu Z**, Dong W. Overexpression of HHLA2, a member of the B7 family, is associated with worse survival in human colorectal carcinoma. *Onco Targets Ther* 2018; **11**: 1563-1570 [PMID: 29593422 DOI: 10.2147/OTT.S160493]



Published by **Baishideng Publishing Group Inc**
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA

Telephone: +1-925-3991568

E-mail: office@baishideng.com

Help Desk: <https://www.f6publishing.com/helpdesk>

<https://www.wjgnet.com>

