



Metabolic dysfunction-associated steatotic liver disease: A silent pandemic

Arghya Samanta, Moinak Sen Sarma

Specialty type: Gastroenterology and hepatology

Provenance and peer review: Invited article; Externally peer reviewed.

Peer-review model: Single blind

Peer-review report's classification

Scientific Quality: Grade B

Novelty: Grade B

Creativity or Innovation: Grade A

Scientific Significance: Grade B

P-Reviewer: Nashwan AJ, Qatar

Received: January 23, 2024

Revised: March 5, 2024

Accepted: April 7, 2024

Published online: April 27, 2024



Arghya Samanta, Moinak Sen Sarma, Department of Pediatric Gastroenterology, Sanjay Gandhi Postgraduate Institute of Medical Sciences, Lucknow 226014, India

Corresponding author: Moinak Sen Sarma, MBBS, MD, Adjunct Associate Professor, Doctor, Department of Pediatric Gastroenterology, Sanjay Gandhi Postgraduate Institute of Medical Sciences, Raebareli Road, Lucknow 226014, India. moinaksen@yahoo.com

Abstract

The worldwide epidemiology of non-alcoholic fatty liver disease (NAFLD) is showing an upward trend, parallel to the rising trend of metabolic syndrome, owing to lifestyle changes. The pathogenesis of NAFLD has not been fully understood yet. Therefore, NAFLD has emerged as a public health concern in the field of hepatology and metabolisms worldwide. Recent changes in the nomenclature from NAFLD to metabolic dysfunction-associated steatotic liver disease have brought a positive outlook changes in the understanding of the disease process and doctor-patient communication. Lifestyle changes are the main treatment modality. Recently, clinical trial using drugs that target 'insulin resistance' which is the driving force behind NAFLD, have shown promising results. Further translational research is needed to better understand the underlying pathophysiological mechanism of NAFLD which may open newer avenues of therapeutic targets. The role of gut dysbiosis in etiopathogenesis and use of fecal microbiota modification in the treatment should be studied extensively. Prevention of this silent epidemic by spreading awareness and early intervention should be our priority.

Key Words: Metabolic dysfunction; Fatty liver; Obesity; Insulin resistance

©The Author(s) 2024. Published by Baishideng Publishing Group Inc. All rights reserved.

Core Tip: Non-alcoholic fatty liver disease is often considered the hepatic manifestation of metabolic syndrome. The new nomenclature of "metabolic dysfunction associated steatotic liver disease" emphasizes the role of disordered metabolism in the pathogenesis. Weight reduction by lifestyle changes is the mainstay of treatment.

Citation: Samanta A, Sen Sarma M. Metabolic dysfunction-associated steatotic liver disease: A silent pandemic. *World J Hepatol* 2024; 16(4): 511-516

URL: <https://www.wjgnet.com/1948-5182/full/v16/i4/511.htm>

DOI: <https://dx.doi.org/10.4254/wjh.v16.i4.511>

INTRODUCTION

Non-alcoholic fatty liver disease (NAFLD) is a broad spectrum of liver disorders related to dysmetabolic conditions. It is characterized by macrovesicular steatosis with or without hepatocellular ballooning, lobular inflammation, and hepatic fibrosis[1]. NAFLD is the leading cause of hepatic morbidity and mortality worldwide and is now the most common indication of liver transplantation (LT)[2-5]. Thus, NAFLD is associated with exorbitant healthcare costs[6,7]. It affects nearly one third of the adult population[8-11] and 9%-12% of the pediatric population[12-14]. The latest meta-analysis by Younossi *et al*[15], showed a global rise in NAFLD prevalence at an alarming rate- from 25.26% (21.59-29.33) in 1990-2006 to 38% (33.71-42.49) in 2016-2019 ($P < 0.001$)[15]. The prevalence in Asia is following a trajectory similar to that in the western countries[16-18].

THE DEBATE OVER THE NEW NOMENCLATURE: NAFLD VS MASLD

NAFLD is usually defined as the presence of steatosis in $>5\%$ hepatocytes, detected by imaging or histopathology after exclusion of secondary causes for hepatic steatosis[19]. Pathologically, it is strongly linked to metabolic syndrome, which is a constellation of obesity, hypertension, hyperlipidemia, type 2 diabetes mellitus (T2DM)[20]. Patients with NAFLD are at higher risk of liver-related complications as well as cardiovascular complications and mortality[20,21]. The most common cause of mortality in patients with NAFLD is cardiovascular complications, followed by extrahepatic malignancies and hepatic complications, highlighting the fact that NAFLD is a multisystemic disease[22,23]. Recently, an international expert group proposed to change the existing nomenclature “Non-Alcoholic Fatty Liver Disease” and adopt the acronym MASLD, or “Metabolic dysfunction-Associated Steatotic Liver Disease”, thus emphasizing the role of systemic metabolic dysfunction in the etiopathogenesis[24]. The shift in the nomenclature introduces a “positive” diagnostic criteria and highlights the cardiovascular risk profile of these individuals. The new nomenclature thus aims for a better understanding of the disease and patient-physician communication.

The diagnosis of MASLD is based on the detection of steatosis of hepatocytes (diagnosed by imaging, biomarkers, or histology) and at least one feature among the following three - overweight/obesity, type 2 diabetes mellitus and metabolic dysregulation. The criterion of metabolic dysregulation is fulfilled when atleast two features among the following are found: increased waist circumference, hypertension, hyperlipidemia, low level of high-density lipoprotein-C (HDL-C), prediabetes, insulin resistance, and subclinical inflammation. These criteria will ensure the identification of a more homogenous disease condition than NAFLD, overcoming the dilemmas and controversies in defining alcohol intake, thereby encouraging new pathophysiological developments and augmenting clinical studies (as elegantly reviewed by Vargas *et al*[25] in this present issue).

LEAN NAFLD

The prevalence of NAFLD showed a rising trend similar to the rising burden of obesity[26,27]. In contrast, lean patients with NAFLD were also detected. In the meta-analysis by Young *et al*[28], 11% and 25% of the general and NAFLD populations, were identified to be “lean NAFLD” respectively[29]. Metabolic profile was more deranged in lean NAFLD patients than healthy controls. These patients also had a higher prevalence of insulin resistance, metabolic syndrome and higher levels of pro-inflammatory mediators[30,31]. On the contrary, lean NAFLD patients have more favorable histologic features than obese NAFLD patients[31].

The ethnicity of the study population should be considered for correctly defining lean NAFLD patients. Body mass index (BMI) cutoffs depending on the ethnicity of the individual have been recommended to define “lean NAFLD”. The cutoffs for defining lean NAFLD are BMI $< 25 \text{ kg/m}^2$ for Caucasians and $< 23 \text{ kg/m}^2$ for Asians[32]. The prevalence of lean NAFLD has been found to be 5%–45% in the Asian population and 5%–20% among Europeans[33]. Further studies are needed to better characterize the newly-defined lean NAFLD patients.

T2DM AND NAFLD

Presence of concomitant T2DM accelerates the disease progression in NAFLD, as patients with concomitant T2DM and NAFLD had higher rates of advanced fibrosis and adverse outcomes compared to NAFLD without T2DM[10]. Furthermore, the concomitant NAFLD and T2DM causes increased liver-related, cardiovascular complications as well as overall mortalities[10]. Other complications of T2DM like diabetic retinopathy, nephropathy, and polyneuropathy, have

been detected more frequently in diabetes patients with coexisting NAFLD[34-36]. Therefore, a novel diagnostic score has been recommended for T2DM patients with NAFLD[37]. Those with an FIB-4 score of more than 1.3 have a higher risk of developing severe disease[38]. Recent NAFLD guidelines recommend that T2DM populations be screened for NAFLD [24].

NAFLD AND METABOLIC SYNDROME

NAFLD is considered as the hepatic manifestation of metabolic syndrome[39]. In the meta-analysis by Ballestri *et al*[40], NAFLD was associated with incident metabolic syndrome in 5-year follow-up. On the other hand, another study by Ma *et al*[41] demonstrated that patients with metabolic syndrome had a higher risk of developing NAFLD. While comparing the new term 'MASLD' with the traditional definition of 'NAFLD', Lin *et al*[42] found higher proportions of metabolic comorbidities in patients with MASLD, emphasizing the impact of positive diagnostic criteria.

DIAGNOSTIC EVALUATION

Liver biopsy is the gold standard to assess disease activity and severity. The severity of liver fibrosis has been identified as the most important prognostic factor and is independently linked with hepatic outcomes in NAFLD patients. Sanyal *et al*[2], in an elegantly done prospective study of 1773 adult patients with NAFLD, found that F3, and F4 fibrosis were associated with increased risk of hepatic complications and death, after adjustment for age, sex, race and diabetes status.

Several non-invasive tests have been developed as diagnostic and prognostic tools in patients of NAFLD as liver biopsy is invasive and less preferred for disease monitoring. Imaging to detect and quantify hepatic steatosis has gained prominence with advances of computerized tomography (liver attenuation index) and magnetic resonance imaging (magnetic resonance imaging proton density fat fraction-MRI-PDFF)[43-45]. Multiparametric MRI, which consists of MR spectroscopy, MR elastography and T1 mapping, has demonstrated high diagnostic accuracy, comparable to liver histology[45,46]. Similar multiparametric CT sequences that can evaluate the hepatic attenuation, liver segmental volume ratio, splenic volume, and liver surface nodularity score, have shown encouraging results as an alternative diagnostic tool to identify advanced fibrosis in NAFLD patients[47]. MRI-PDFF response has been studied as a potential surrogate for histologic improvement after treatment of NAFLD. Several studies have shown a clear correlation between a reduction in MRI-PDFF (usually taken as a $\geq 30\%$ relative reduction) and improvement in the NAFLD activity score, resolution of NASH, and fibrosis[48,49]. Boursier *et al*[50], in a large cohort of 1097 patients, compared the prognostic efficacy of fibrosis index based on 4-factors (FIB4), transient elastography (TE) and liver biopsy. The results showed that FIB4 and TE showed good accuracy for the prediction of liver-related events (LRE), with Harrell's C-indexes > 0.80 [0.817 (0.768-0.866) *vs.* 0.878 (0.835-0.921), respectively, $P = 0.059$], as compared to liver biopsy. The authors proposed a stepwise algorithm to accurately stratify NAFLD patients based on their risk for LRE: compared to patients with "FIB4 < 1.30 ", those with "FIB4 ≥ 1.30 then TE < 8.0 kPa" had a similar risk of LREs [adjusted hazard ratio (aHR) 1.3; 95%CI 0.3-6.8], whereas the risk of LREs significantly increased in patients with "FIB4 ≥ 1.30 then TE 8.0-12.0 kPa" (aHR 3.8; 95%CI 1.3-10.9), and even more for those with "FIB4 ≥ 1.30 then TE > 12.0 kPa" (aHR 12.4; 95%CI 5.1-30.2).

However, we need to keep in mind a major limitation of using these non-invasive methods to diagnose and monitor hepatic steatosis is that it provides no information on the underlying etiology or associated risk factors.

TREATMENT

As NAFLD is a systemic disease of disordered metabolism, a multi-disciplinary approach is of utmost importance for the treatment. Weight reduction by lifestyle changes and dietary interventions is the cornerstone of treatment in obese and lean NAFLD patients[19,51]. The beneficial effects of lifestyle modifications have been consistently found to be helpful in the resolution of hepatic steatosis in both lean Asian and Caucasian NAFLD patients[51-53]. Medical treatment with glucagon-like peptide receptor agonists, sodium-glucose cotransporter-2 inhibitors, and peroxisome proliferator-activated receptor- γ agonists was able to improve inflammation and fibrosis, as well as reduction in blood pressure, better glycemic control and lipid profile[54-56]. Bariatric surgery is an effective treatment for a select group of patients who are non-responsive to dietary interventions and exercise or unable to lose weight through lifestyle changes. It can improve both histological characteristics of NASH as well as mortality due to cardiovascular complications[57]. Lim *et al*[58] studied the usefulness of endoscopic bariatric therapies such as intragastric balloon, endoscopic sleeve gastropasty, and duodeno-jejunal bypass liner in reducing weight and found better results as compared to standard medical therapy. With extensive research into the therapeutic options in the pipeline, treatment strategies for NAFLD treatment are promising[59].

FUTURE PERSPECTIVE

Further prospective studies are the need of the hour to develop more accurate diagnostic tools for advanced fibrosis in NAFLD and to explore the underlying pathophysiological mechanisms linking NAFLD with other conditions. More in-depth research on gut microbiota in the etiopathogenesis of NAFLD and its role in the therapeutics is warranted. Most

randomized clinical trials of available drugs do not reflect the proper scenario, due to the limitations of therapeutic targets, drug safety, and other factors. This current review by Vargas in this issue serves as a valuable resource for researchers seeking a comprehensive understanding of NAFLD and tries to address these issues[25].

CONCLUSION

NAFLD must be evaluated as a multisystemic metabolic disorder. It may lead to liver-related complications, thus the need for multidisciplinary screening and disease management cannot be over emphasized. Routine screening for NAFLD is recommended in patients with metabolic syndrome. Lifestyle intervention remains the most important treatment modality. The global pandemic of NAFLD poses significant social and economic burden; thus it is of utmost importance to create widespread awareness in order to make early interventions and achieve better outcome.

FOOTNOTES

Author contributions: Samanta A did the literature review and wrote the original manuscript; Sen Sarma M reviewed and revised the manuscript, did critical analysis; All authors approved the final draft of the manuscript.

Conflict-of-interest statement: All authors declare no conflict of interest.

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <https://creativecommons.org/licenses/by-nc/4.0/>

Country/Territory of origin: India

ORCID number: Arghya Samanta 0000-0002-1768-0263; Moinak Sen Sarma 0000-0003-2015-4069.

S-Editor: Gong ZM

L-Editor: A

P-Editor: Cai YX

REFERENCES

- 1 Brunt EM, Kleiner DE, Carpenter DH, Rinella M, Harrison SA, Loomba R, Younossi Z, Neuschwander-Tetri BA, Sanyal AJ; American Association for the Study of Liver Diseases NASH Task Force. NAFLD: Reporting Histologic Findings in Clinical Practice. *Hepatology* 2021; **73**: 2028-2038 [PMID: 33111374 DOI: 10.1002/hep.31599]
- 2 Sanyal AJ, Van Natta ML, Clark J, Neuschwander-Tetri BA, Diehl A, Dasarathy S, Loomba R, Chalasani N, Kowdley K, Hameed B, Wilson LA, Yates KP, Belt P, Lazo M, Kleiner DE, Behling C, Tonascia J; NASH Clinical Research Network (CRN). Prospective Study of Outcomes in Adults with Nonalcoholic Fatty Liver Disease. *N Engl J Med* 2021; **385**: 1559-1569 [PMID: 34670043 DOI: 10.1056/NEJMoa2029349]
- 3 Horn P, Newsome PN. Nonalcoholic Fatty Liver Disease. *N Engl J Med* 2022; **386**: 294 [PMID: 35045235 DOI: 10.1056/NEJMc2118255]
- 4 Pais R, Barritt AS 4th, Calmus Y, Scatton O, Runge T, Lebray P, Poynard T, Ratziu V, Conti F. NAFLD and liver transplantation: Current burden and expected challenges. *J Hepatol* 2016; **65**: 1245-1257 [PMID: 27486010 DOI: 10.1016/j.jhep.2016.07.033]
- 5 Battistella S, D'Arcangelo F, Grasso M, Zanetto A, Gambato M, Germani G, Senzolo M, Russo FP, Burra P. Liver transplantation for non-alcoholic fatty liver disease: indications and post-transplant management. *Clin Mol Hepatol* 2023; **29**: S286-S301 [PMID: 36577425 DOI: 10.3350/cmh.2022.0392]
- 6 Allen AM, Lazarus JV, Younossi ZM. Healthcare and socioeconomic costs of NAFLD: A global framework to navigate the uncertainties. *J Hepatol* 2023; **79**: 209-217 [PMID: 36740046 DOI: 10.1016/j.jhep.2023.01.026]
- 7 Premkumar M, Anand AC. Overview of Complications in Cirrhosis. *J Clin Exp Hepatol* 2022; **12**: 1150-1174 [PMID: 35814522 DOI: 10.1016/j.jceh.2022.04.021]
- 8 Younossi Z, Anstee QM, Marietti M, Hardy T, Henry L, Eslam M, George J, Bugianesi E. Global burden of NAFLD and NASH: trends, predictions, risk factors and prevention. *Nat Rev Gastroenterol Hepatol* 2018; **15**: 11-20 [PMID: 28930295 DOI: 10.1038/nrgastro.2017.109]
- 9 Riazi K, Azhari H, Charette JH, Underwood FE, King JA, Afshar EE, Swain MG, Congly SE, Kaplan GG, Shaheen AA. The prevalence and incidence of NAFLD worldwide: a systematic review and meta-analysis. *Lancet Gastroenterol Hepatol* 2022; **7**: 851-861 [PMID: 35798021 DOI: 10.1016/S2468-1253(22)00165-0]
- 10 Younossi ZM, Golabi P, de Avila L, Paik JM, Srishord M, Fukui N, Qiu Y, Burns L, Afendy A, Nader F. The global epidemiology of NAFLD and NASH in patients with type 2 diabetes: A systematic review and meta-analysis. *J Hepatol* 2019; **71**: 793-801 [PMID: 31279902 DOI: 10.1016/j.jhep.2019.06.021]
- 11 Younossi ZM, Koenig AB, Abdelatif D, Fazel Y, Henry L, Wymer M. Global epidemiology of nonalcoholic fatty liver disease-Meta-analytic assessment of prevalence, incidence, and outcomes. *Hepatology* 2016; **64**: 73-84 [PMID: 26707365 DOI: 10.1002/hep.28431]
- 12 Quek J, Chan KE, Wong ZY, Tan C, Tan B, Lim WH, Tan DJH, Tang ASP, Tay P, Xiao J, Yong JN, Zeng RW, Chew NWS, Nah B, Kulkarni A, Siddiqui MS, Dan YY, Wong VW, Sanyal AJ, Noureddin M, Muthiah M, Ng CH. Global prevalence of non-alcoholic fatty liver disease and non-alcoholic steatohepatitis in the overweight and obese population: a systematic review and meta-analysis. *Lancet Gastroenterol Hepatol*

- 2023; **8**: 20-30 [PMID: 36400097 DOI: 10.1016/S2468-1253(22)00317-X]
- 13 **Anderson EL**, Howe LD, Jones HE, Higgins JP, Lawlor DA, Fraser A. The Prevalence of Non-Alcoholic Fatty Liver Disease in Children and Adolescents: A Systematic Review and Meta-Analysis. *PLoS One* 2015; **10**: e0140908 [PMID: 26512983 DOI: 10.1371/journal.pone.0140908]
 - 14 **Nobili V**, Alisi A, Valenti L, Miele L, Feldstein AE, Alkhouri N. NAFLD in children: new genes, new diagnostic modalities and new drugs. *Nat Rev Gastroenterol Hepatol* 2019; **16**: 517-530 [PMID: 31278377 DOI: 10.1038/s41575-019-0169-z]
 - 15 **Younossi ZM**, Golabi P, Paik JM, Henry A, Van Dongen C, Henry L. The global epidemiology of nonalcoholic fatty liver disease (NAFLD) and nonalcoholic steatohepatitis (NASH): a systematic review. *Hepatology* 2023; **77**: 1335-1347 [PMID: 36626630 DOI: 10.1097/HEP.0000000000000004]
 - 16 **Zhou J**, Zhou F, Wang W, Zhang XJ, Ji YX, Zhang P, She ZG, Zhu L, Cai J, Li H. Epidemiological Features of NAFLD From 1999 to 2018 in China. *Hepatology* 2020; **71**: 1851-1864 [PMID: 32012320 DOI: 10.1002/hep.31150]
 - 17 **Fan JG**, Kim SU, Wong VW. New trends on obesity and NAFLD in Asia. *J Hepatol* 2017; **67**: 862-873 [PMID: 28642059 DOI: 10.1016/j.jhep.2017.06.003]
 - 18 **Li J**, Zou B, Yeo YH, Feng Y, Xie X, Lee DH, Fujii H, Wu Y, Kam LY, Ji F, Li X, Chien N, Wei M, Ogawa E, Zhao C, Wu X, Stave CD, Henry L, Barnett S, Takahashi H, Furusyo N, Eguchi Y, Hsu YC, Lee TY, Ren W, Qin C, Jun DW, Toyoda H, Wong VW, Cheung R, Zhu Q, Nguyen MH. Prevalence, incidence, and outcome of non-alcoholic fatty liver disease in Asia, 1999-2019: a systematic review and meta-analysis. *Lancet Gastroenterol Hepatol* 2019; **4**: 389-398 [PMID: 30902670 DOI: 10.1016/S2468-1253(19)30039-1]
 - 19 **Chalasani N**, Younossi Z, Lavine JE, Charlton M, Cusi K, Rinella M, Harrison SA, Brunt EM, Sanyal AJ. The diagnosis and management of nonalcoholic fatty liver disease: Practice guidance from the American Association for the Study of Liver Diseases. *Hepatology* 2018; **67**: 328-357 [PMID: 28714183 DOI: 10.1002/hep.29367]
 - 20 **Ng CH**, Chan KE, Chin YH, Zeng RW, Tsai PC, Lim WH, Tan DJH, Khoo CM, Goh LH, Ling ZJ, Kulkarni A, Mak LL, Huang DQ, Chan M, Chew NW, Siddiqui MS, Sanyal AJ, Muthiah M. The effect of diabetes and prediabetes on the prevalence, complications and mortality in nonalcoholic fatty liver disease. *Clin Mol Hepatol* 2022; **28**: 565-574 [PMID: 35585687 DOI: 10.3350/cmh.2022.0096]
 - 21 **Sanyal AJ**, Campbell-Sargent C, Mirshahi F, Rizzo WB, Contos MJ, Sterling RK, Luketic VA, Shiffman ML, Clore JN. Nonalcoholic steatohepatitis: association of insulin resistance and mitochondrial abnormalities. *Gastroenterology* 2001; **120**: 1183-1192 [PMID: 11266382 DOI: 10.1053/gast.2001.23256]
 - 22 **Mantovani A**, Scorletti E, Mosca A, Alisi A, Byrne CD, Targher G. Complications, morbidity and mortality of nonalcoholic fatty liver disease. *Metabolism* 2020; **111S**: 154170 [PMID: 32006558 DOI: 10.1016/j.metabol.2020.154170]
 - 23 **Muzurović E**, Peng CC, Belanger MJ, Sanoudou D, Mikhailidis DP, Mantzoros CS. Nonalcoholic Fatty Liver Disease and Cardiovascular Disease: a Review of Shared Cardiometabolic Risk Factors. *Hypertension* 2022; **79**: 1319-1326 [PMID: 35465684 DOI: 10.1161/HYPERTENSIONAHA.122.17982]
 - 24 **Eslam M**, Sanyal AJ, George J; International Consensus Panel. MAFLD: A Consensus-Driven Proposed Nomenclature for Metabolic Associated Fatty Liver Disease. *Gastroenterology* 2020; **158**: 1999-2014.e1 [PMID: 32044314 DOI: 10.1053/j.gastro.2019.11.312]
 - 25 **Vargas M**, Cardoso Toniasso SC, Riedel PG, Baldin CP, Dos Reis FL, Pereira RM, Brum MCB, Joveleviths D, Alvares-da-Silva MR. Metabolic disease and the liver: A review. *World J Hepatol* 2024; **16**: 33-40 [PMID: 38313243 DOI: 10.4254/wjh.v16.i1.33]
 - 26 **Li L**, Liu DW, Yan HY, Wang ZY, Zhao SH, Wang B. Obesity is an independent risk factor for non-alcoholic fatty liver disease: evidence from a meta-analysis of 21 cohort studies. *Obes Rev* 2016; **17**: 510-519 [PMID: 27020692 DOI: 10.1111/obr.12407]
 - 27 **Finucane MM**, Stevens GA, Cowan MJ, Danaei G, Lin JK, Paciorek CJ, Singh GM, Gutierrez HR, Lu Y, Bahalim AN, Farzadfar F, Riley LM, Ezzati M; Global Burden of Metabolic Risk Factors of Chronic Diseases Collaborating Group (Body Mass Index). National, regional, and global trends in body-mass index since 1980: systematic analysis of health examination surveys and epidemiological studies with 960 country-years and 9.1 million participants. *Lancet* 2011; **377**: 557-567 [PMID: 21295846 DOI: 10.1016/S0140-6736(10)62037-5]
 - 28 **Young S**, Tariq R, Provenza J, Satapathy SK, Faisal K, Choudhry A, Friedman SL, Singal AK. Prevalence and Profile of Nonalcoholic Fatty Liver Disease in Lean Adults: Systematic Review and Meta-Analysis. *Hepatol Commun* 2020; **4**: 953-972 [PMID: 32626829 DOI: 10.1002/hep4.1519]
 - 29 **Kumar R**, Rastogi A, Sharma MK, Bhatia V, Garg H, Bihari C, Sarin SK. Clinicopathological characteristics and metabolic profiles of non-alcoholic fatty liver disease in Indian patients with normal body mass index: Do they differ from obese or overweight non-alcoholic fatty liver disease? *Indian J Endocrinol Metab* 2013; **17**: 665-671 [PMID: 23961483 DOI: 10.4103/2230-8210.113758]
 - 30 **Fracanzani AL**, Petta S, Lombardi R, Pisano G, Russello M, Consonni D, Di Marco V, Cammà C, Mensi L, Dongiovanni P, Valenti L, Craxi A, Fargion S. Liver and Cardiovascular Damage in Patients With Lean Nonalcoholic Fatty Liver Disease, and Association With Visceral Obesity. *Clin Gastroenterol Hepatol* 2017; **15**: 1604-1611.e1 [PMID: 28554682 DOI: 10.1016/j.cgh.2017.04.045]
 - 31 **Chakrabarty M**, Jha AN, Sharma DJ. Clinical Characteristics and Metabolic Profiles of Non-Alcoholic Fatty Liver Disease (NAFLD) in Lean Patients and Their Comparison with Obese and Overweight NAFLD. *J Assoc Physicians India* 2022; **70**: 11-12 [PMID: 35443531]
 - 32 Obesity: preventing and managing the global epidemic. Report of a WHO consultation. *World Health Organ Tech Rep Ser* 2000; **894**: i-xii, 1 [PMID: 11234459]
 - 33 **Eslam M**, Chen F, George J. NAFLD in Lean Asians. *Clin Liver Dis (Hoboken)* 2020; **16**: 240-243 [PMID: 33489095 DOI: 10.1002/cld.930]
 - 34 **Mantovani A**, Byrne CD, Bonora E, Targher G. Nonalcoholic Fatty Liver Disease and Risk of Incident Type 2 Diabetes: A Meta-analysis. *Diabetes Care* 2018; **41**: 372-382 [PMID: 29358469 DOI: 10.2337/dc17-1902]
 - 35 **Hazlehurst JM**, Woods C, Marjot T, Cobbold JF, Tomlinson JW. Non-alcoholic fatty liver disease and diabetes. *Metabolism* 2016; **65**: 1096-1108 [PMID: 26856933 DOI: 10.1016/j.metabol.2016.01.001]
 - 36 **Targher G**, Lonardo A, Byrne CD. Nonalcoholic fatty liver disease and chronic vascular complications of diabetes mellitus. *Nat Rev Endocrinol* 2018; **14**: 99-114 [PMID: 29286050 DOI: 10.1038/nrendo.2017.173]
 - 37 **Younossi ZM**, Corey KE, Alkhouri N, Noureddin M, Jacobson I, Lam B, Clement S, Basu R, Gordon SC, Ravendhra N, Puri P, Rinella M, Scudera P, Singal AK, Henry L; US Members of the Global Nash Council. Clinical assessment for high-risk patients with non-alcoholic fatty liver disease in primary care and diabetology practices. *Aliment Pharmacol Ther* 2020; **52**: 513-526 [PMID: 32598051 DOI: 10.1111/apt.15830]
 - 38 **Yilmaz Y**, Kaya E, Eren F. Letter: the use of Fibrosis-4 score in primary care and diabetology practices-Occam's razor applied to advanced fibrosis screening. *Aliment Pharmacol Ther* 2020; **52**: 1759-1760 [PMID: 33205873 DOI: 10.1111/apt.16034]
 - 39 **Kim D**, Tournas A, Kim WR. Nonalcoholic Fatty Liver Disease and Metabolic Syndrome. *Clin Liver Dis* 2018; **22**: 133-140 [PMID: 29128053 DOI: 10.1016/j.cld.2017.08.010]

- 40 **Ballestri S**, Zona S, Targher G, Romagnoli D, Baldelli E, Nascimbeni F, Roverato A, Guaraldi G, Lonardo A. Nonalcoholic fatty liver disease is associated with an almost twofold increased risk of incident type 2 diabetes and metabolic syndrome. Evidence from a systematic review and meta-analysis. *J Gastroenterol Hepatol* 2016; **31**: 936-944 [PMID: 26667191 DOI: 10.1111/jgh.13264]
- 41 **Ma J**, Hwang SJ, Pedley A, Massaro JM, Hoffmann U, Chung RT, Benjamin EJ, Levy D, Fox CS, Long MT. Bi-directional analysis between fatty liver and cardiovascular disease risk factors. *J Hepatol* 2017; **66**: 390-397 [PMID: 27729222 DOI: 10.1016/j.jhep.2016.09.022]
- 42 **Lin S**, Huang J, Wang M, Kumar R, Liu Y, Liu S, Wu Y, Wang X, Zhu Y. Comparison of MAFLD and NAFLD diagnostic criteria in real world. *Liver Int* 2020; **40**: 2082-2089 [PMID: 32478487 DOI: 10.1111/liv.14548]
- 43 **Schaapman JJ**, Tushuizen ME, Coenraad MJ, Lamb HJ. Multiparametric MRI in Patients With Nonalcoholic Fatty Liver Disease. *J Magn Reson Imaging* 2021; **53**: 1623-1631 [PMID: 32822095 DOI: 10.1002/jmri.27292]
- 44 **van Werven JR**, Marsman HA, Nederveen AJ, Smits NJ, ten Kate FJ, van Gulik TM, Stoker J. Assessment of hepatic steatosis in patients undergoing liver resection: comparison of US, CT, T1-weighted dual-echo MR imaging, and point-resolved 1H MR spectroscopy. *Radiology* 2010; **256**: 159-168 [PMID: 20574093 DOI: 10.1148/radiol.10091790]
- 45 **Park CC**, Nguyen P, Hernandez C, Bettencourt R, Ramirez K, Fortney L, Hooker J, Sy E, Savides MT, Alquirraish MH, Valasek MA, Rizo E, Richards L, Brenner D, Sirlin CB, Loomba R. Magnetic Resonance Elastography vs Transient Elastography in Detection of Fibrosis and Noninvasive Measurement of Steatosis in Patients With Biopsy-Proven Nonalcoholic Fatty Liver Disease. *Gastroenterology* 2017; **152**: 598-607.e2 [PMID: 27911262 DOI: 10.1053/j.gastro.2016.10.026]
- 46 **Bannas P**, Kramer H, Hernando D, Agni R, Cunningham AM, Mandal R, Motosugi U, Sharma SD, Munoz del Rio A, Fernandez L, Reeder SB. Quantitative magnetic resonance imaging of hepatic steatosis: Validation in *ex vivo* human livers. *Hepatology* 2015; **62**: 1444-1455 [PMID: 26224591 DOI: 10.1002/hep.28012]
- 47 **Lubner MG**, Graffy PM, Said A, Watson R, Zea R, Malecki KM, Pickhardt PJ. Utility of Multiparametric CT for Identification of High-Risk NAFLD. *AJR Am J Roentgenol* 2021; **216**: 659-668 [PMID: 33474981 DOI: 10.2214/AJR.20.22842]
- 48 **Stine JG**, Munaganuru N, Barnard A, Wang JL, Kaulback K, Argo CK, Singh S, Fowler KJ, Sirlin CB, Loomba R. Change in MRI-PDFF and Histologic Response in Patients With Nonalcoholic Steatohepatitis: A Systematic Review and Meta-Analysis. *Clin Gastroenterol Hepatol* 2021; **19**: 2274-2283.e5 [PMID: 32882428 DOI: 10.1016/j.cgh.2020.08.061]
- 49 **Tamaki N**, Munaganuru N, Jung J, Yonan AQ, Loomba RR, Bettencourt R, Ajmera V, Valasek MA, Behling C, Sirlin CB, Loomba R. Clinical utility of 30% relative decline in MRI-PDFF in predicting fibrosis regression in non-alcoholic fatty liver disease. *Gut* 2022; **71**: 983-990 [PMID: 33883248 DOI: 10.1136/gutjnl-2021-324264]
- 50 **Boursier J**, Hagström H, Ekstedt M, Moreau C, Bonacci M, Cure S, Ampuero J, Nasr P, Tallab L, Canivet CM, Kechagias S, Sánchez Y, Dincuff E, Lucena A, Roux M, Riou J, Trylesinski A, Romero-Gomez M. Non-invasive tests accurately stratify patients with NAFLD based on their risk of liver-related events. *J Hepatol* 2022; **76**: 1013-1020 [PMID: 35063601 DOI: 10.1016/j.jhep.2021.12.031]
- 51 **Risi R**, Tozzi R, Watanabe M. Beyond weight loss in nonalcoholic fatty liver disease: the role of carbohydrate restriction. *Curr Opin Clin Nutr Metab Care* 2021; **24**: 349-353 [PMID: 33871418 DOI: 10.1097/MCO.0000000000000762]
- 52 **Hamurcu Varol P**, Kaya E, Alphan E, Yilmaz Y. Role of intensive dietary and lifestyle interventions in the treatment of lean nonalcoholic fatty liver disease patients. *Eur J Gastroenterol Hepatol* 2020; **32**: 1352-1357 [PMID: 32092046 DOI: 10.1097/MEG.0000000000001656]
- 53 **Brunner KT**, Henneberg CJ, Wilechansky RM, Long MT. Nonalcoholic Fatty Liver Disease and Obesity Treatment. *Curr Opin Clin Nutr Metab Rep* 2019; **8**: 220-228 [PMID: 30945129 DOI: 10.1007/s13679-019-00345-1]
- 54 **Ng CH**, Lin SY, Chin YH, Lee MH, Syn N, Goh XL, Koh JH, Quek J, Hao Tan DJ, Mok SF, Tan E, Dan YY, Chew N, Khoo CM, Siddiqui MS, Muthiah M. Antidiabetic Medications for Type 2 Diabetics with Nonalcoholic Fatty Liver Disease: Evidence From a Network Meta-Analysis of Randomized Controlled Trials. *Endocr Pract* 2022; **28**: 223-230 [PMID: 34606980 DOI: 10.1016/j.eprac.2021.09.013]
- 55 **Feng W**, Gao C, Bi Y, Wu M, Li P, Shen S, Chen W, Yin T, Zhu D. Randomized trial comparing the effects of gliclazide, liraglutide, and metformin on diabetes with non-alcoholic fatty liver disease. *J Diabetes* 2017; **9**: 800-809 [PMID: 28332301 DOI: 10.1111/1753-0407.12555]
- 56 **Dwinata M**, Putera DD, Hasan I, Raharjo M. SGLT2 inhibitors for improving hepatic fibrosis and steatosis in non-alcoholic fatty liver disease complicated with type 2 diabetes mellitus: a systematic review. *Clin Exp Hepatol* 2020; **6**: 339-346 [PMID: 33511282 DOI: 10.5114/ceh.2020.102173]
- 57 **Chauhan M**, Singh K, Thuluvath PJ. Bariatric Surgery in NAFLD. *Dig Dis Sci* 2022; **67**: 408-422 [PMID: 34981313 DOI: 10.1007/s10620-021-07317-3]
- 58 **Lim WH**, Lin SY, Ng CH, Tan DJH, Xiao J, Yong JN, Tay PWL, Syn N, Chin YH, Chan KE, Khoo CM, Chew N, Foo RSY, Shabbir A, Tan EX, Huang DQ, Nouredin M, Sanyal AJ, Siddiqui MS, Muthiah MD. Foregut bypass vs. restrictive bariatric procedures for nonalcoholic fatty liver disease: a meta-analysis of 3,355 individuals. *Hepatobiliary Surg Nutr* 2023; **12**: 658-670 [PMID: 37886204 DOI: 10.21037/hbsn-21-520]
- 59 **Chew NWS**, Ng CH, Truong E, Nouredin M, Kowdley KV. Nonalcoholic Steatohepatitis Drug Development Pipeline: An Update. *Semin Liver Dis* 2022; **42**: 379-400 [PMID: 35709720 DOI: 10.1055/a-1877-9656]



Published by **Baishideng Publishing Group Inc**
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA

Telephone: +1-925-3991568

E-mail: office@baishideng.com

Help Desk: <https://www.f6publishing.com/helpdesk>

<https://www.wjgnet.com>

