



Retrospective Study

Sepsis during short bowel syndrome hospitalizations: Identifying trends, disparities, and clinical outcomes in the United States

Dushyant Singh Dahiya, Jennifer Wachala, Shantanu Solanki, Dhanshree Solanki, Asim Kichloo, Samantha Holcomb, Uvesh Mansuri, Khwaja Saad Haq, Hassam Ali, Manesh Kumar Gangwani, Yash R Shah, Teresa Varghese, Hafiz Muzaffar Akbar Khan, Simon Peter Horslen, Thomas D Schiano, Syed-Mohammed Jafri

Specialty type: Gastroenterology and hepatology

Provenance and peer review: Invited article; Externally peer reviewed.

Peer-review model: Single blind

Peer-review report's scientific quality classification

Grade A (Excellent): 0
Grade B (Very good): B
Grade C (Good): 0
Grade D (Fair): 0
Grade E (Poor): 0

P-Reviewer: Sanayeh EB, United States

Received: January 15, 2024

Peer-review started: January 15, 2024

First decision: February 3, 2024

Revised: February 10, 2024

Accepted: March 25, 2024

Article in press: March 25, 2024

Published online: April 22, 2024



Dushyant Singh Dahiya, Division of Gastroenterology, Hepatology & Motility, The University of Kansas School of Medicine, Kansas City, KS 66160, United States

Jennifer Wachala, Asim Kichloo, Samantha Holcomb, Department of Internal Medicine, Samaritan Medical Center, Watertown, NY 13601, United States

Shantanu Solanki, Division of Gastroenterology Hepatology & Nutrition, The University of Texas MD Anderson Cancer Center, Houston, TX 77030, United States

Dhanshree Solanki, Department of Medicine, Institute for Foreign Medical Graduate Education, Houston, TX 77030, United States

Uvesh Mansuri, Department of Internal Medicine, MedStar Harbor Hospital, Baltimore, MD 21225, United States

Khwaja Saad Haq, Teresa Varghese, Department of Internal Medicine, WellStar Spalding Regional Hospital, Griffin, GA 30224, United States

Hassam Ali, Division of Gastroenterology, Hepatology and Nutrition, East Carolina University/Brody School of Medicine, Greenville, NC 27858, United States

Manesh Kumar Gangwani, Department of Gastroenterology and Hepatology, University of Arkansas for Medical Sciences, Little Rock, AR 72205, United States

Yash R Shah, Department of Internal Medicine, Trinity Health Oakland/Wayne State University, Pontiac, MI 48341, United States

Hafiz Muzaffar Akbar Khan, Division of Gastroenterology and Hepatology, SUNY Upstate Medical University, Syracuse, NY 13210, United States

Simon Peter Horslen, Department of Pediatrics, School of Medicine and UPMC Children's Hospital of Pittsburgh, University of Pittsburgh, Pittsburgh, PA 15219, United States

Thomas D Schiano, Division of Liver Diseases, Icahn School of Medicine at Mount Sinai, New York, NY 10029, United States

Syed-Mohammed Jafri, Division of Gastroenterology and Hepatology, Henry Ford Hospital, Detroit, MI 48202, United States

Corresponding author: Dushyant Singh Dahiya, MD, Doctor, Division of Gastroenterology, Hepatology & Motility, The University of Kansas School of Medicine, 2000 Olathe Blvd, Kansas City, KS 66160, United States. dush.dahiya@gmail.com

Abstract

BACKGROUND

Short bowel syndrome (SBS) hospitalizations are often complicated with sepsis. There is a significant paucity of data on adult SBS hospitalizations in the United States and across the globe.

AIM

To assess trends and outcomes of SBS hospitalizations complicated by sepsis in the United States.

METHODS

The National Inpatient Sample was utilized to identify all adult SBS hospitalizations between 2005-2014. The study cohort was further divided based on the presence or absence of sepsis. Trends were identified, and hospitalization characteristics and clinical outcomes were compared. Predictors of mortality for SBS hospitalizations complicated with sepsis were assessed.

RESULTS

Of 247097 SBS hospitalizations, 21.7% were complicated by sepsis. Septic SBS hospitalizations had a rising trend of hospitalizations from 20.8% in 2005 to 23.5% in 2014 (P trend < 0.0001). Compared to non-septic SBS hospitalizations, septic SBS hospitalizations had a higher proportion of males (32.8% *vs* 29.3%, $P < 0.0001$), patients in the 35-49 (45.9% *vs* 42.5%, $P < 0.0001$) and 50-64 (32.1% *vs* 31.1%, $P < 0.0001$) age groups, and ethnic minorities, *i.e.*, Blacks (12.4% *vs* 11.3%, $P < 0.0001$) and Hispanics (6.7% *vs* 5.5%, $P < 0.0001$). Furthermore, septic SBS hospitalizations had a higher proportion of patients with intestinal transplantation (0.33% *vs* 0.22%, $P < 0.0001$), inpatient mortality (8.5% *vs* 1.4%, $P < 0.0001$), and mean length of stay (16.1 d *vs* 7.7 d, $P < 0.0001$) compared to the non-sepsis cohort. A younger age, female gender, White race, and presence of comorbidities such as anemia and depression were identified to be independent predictors of inpatient mortality for septic SBS hospitalizations.

CONCLUSION

Septic SBS hospitalizations had a rising trend between 2005-2014 and were associated with higher inpatient mortality compared to non-septic SBS hospitalizations.

Key Words: Short bowel syndrome; Sepsis; Outcomes; Mortality; Trends

©The Author(s) 2024. Published by Baishideng Publishing Group Inc. All rights reserved.

Core Tip: Short bowel syndrome (SBS) is a well-known complication of small bowel surgical resection. Sepsis is a well-documented complication of SBS, particularly in infants and children. However, there is limited data on adult SBS hospitalizations complicated by sepsis in the United States. In this study, we noted that about one-fifth of SBS hospitalizations were complicated by sepsis. There was a higher proportion of men, individuals in the 35-64 age group, and ethnic minorities (Blacks and Hispanics) in the septic SBS cohort compared to the non-sepsis cohort. Septic SBS hospitalizations also had a higher length of stay and inpatient mortality compared to the non-sepsis cohort. Furthermore, younger age, female gender, White race, anemia, and depression were identified to be independent predictors of inpatient mortality for septic SBS hospitalizations.

Citation: Dahiya DS, Wachala J, Solanki S, Solanki D, Kichloo A, Holcomb S, Mansuri U, Haq KS, Ali H, Gangwani MK, Shah YR, Varghese T, Khan HMA, Horslen SP, Schiano TD, Jafri SM. Sepsis during short bowel syndrome hospitalizations: Identifying trends, disparities, and clinical outcomes in the United States. *World J Gastrointest Pathophysiol* 2024; 15(1): 92085

URL: <https://www.wjgnet.com/2150-5330/full/v15/i1/92085.htm>

DOI: <https://dx.doi.org/10.4291/wjgp.v15.i1.92085>

INTRODUCTION

Short bowel syndrome (SBS) is a well-known complication of surgical resection of the small bowel[1]. Nonsurgical causes include inflammatory bowel disease, cancer of the intestine, or ischemic and hemorrhagic vascular diseases of the gut[1, 2]. Sepsis is a well-documented complication of SBS in infants and children, and these recurrent bloodstream infections (BSI) have been associated with higher rates of childhood morbidity and mortality[3,4]. The primary pathophysiologic mechanism implicated in the development of BSI is bacterial translocation from the gut to the bloodstream during enteral

feeding in children with SBS[5].

Most patients with SBS derive their nutrition *via* parenteral routes using indwelling venous catheters. Hence, these patients are at a greater-than-average risk of BSI from skin flora, especially if the indwelling catheter has been placed for a prolonged duration[4]. Moreover, parenteral nutrition leads to the impairment of immunological barriers (altered inflammatory responses) and physical barriers (secondary to villous atrophy) which may increase the risk of small bowel bacterial overgrowth (SBBO)[4,6].

SBS has established itself to be one of the strongest predictors of BSI in the pediatric population. However, there is a significant paucity of data on adult SBS hospitalizations complicated by sepsis, both in the United States and across the globe. Hence, this study was designed to investigate trends, hospitalization characteristics, predictors, racial disparities of Elixhauser co-morbidities, and gender disparities of Elixhauser co-morbidities for septic SBS hospitalizations in the United States. Furthermore, we also performed a comparative analysis for trends, hospitalization characteristics, and predictors between septic SBS and non-septic SBS hospitalizations.

MATERIALS AND METHODS

Data source

The National Inpatient Sample (NIS) is the largest, publicly available, multi-ethnic, all-payer inpatient database which is a part of the healthcare cost and utilization project (HCUP)[7,8]. HCUP is a family of healthcare databases and related software tools developed through a unique Federal-State-Industry partnership and sponsored by the Agency for Healthcare Research and Quality[9]. The NIS, when weighted, estimates more than 35 million hospitalizations nationally [7]. HCUP databases are limited data sets, which can be used to generate United States regional and national estimates[7].

Study population, design, and outcomes

We used the International Classification of Diseases, Clinical Modification (ICD-9-CM) codes to identify all adults (≥ 18 years of age) with SBS from the NIS database for the 2005-2014 period. The precedence of utilization of ICD-9-CM codes for SBS hospitalizations has already been established in previous studies[10,11]. Individuals < 18 years of age were excluded from the analysis. This study population was further divided based on the presence and absence of sepsis using the Clinical Classification Software diagnosis code "2", which has been used previously in multiple NIS-based studies[12, 13]. We then compared hospitalization characteristics (age, race, and gender), hospital-level characteristics (bed size, location, and admission type), and clinical outcomes [length of stay (LOS), all-cause inpatient mortality, and disposition status] between septic and non-septic SBS hospitalizations. Furthermore, the rates of comorbidities were calculated using Elixhauser Comorbidity Index codes provided by HCUP[14].

Statistical analysis

Statistical Analysis Software 9.3 (SAS Institute, Cary, NC, United States) was used for univariate and multivariate analyses. We used weighted values provided by HCUP to produce nationally representative estimates for all variables [15]. Categorical variables like gender, race, and comorbidities were compared using the Chi-squared (χ^2) test, and continuous variables like age and LOS were compared using the Wilcoxon rank-sum test. We also created a multivariate logistic regression model to determine predictors of inpatient mortality for SBS hospitalizations complicated by sepsis. P value ≤ 0.05 was considered statistically significant.

Ethical considerations

A review by our institutional review board was not required as the NIS database is Health Insurance Portability and Accountability Act protected and does not contain identifiable patient and hospital-level data[10].

RESULTS

Trends of septic SBS hospitalizations

Between 2005-2014, there were 247097 adult SBS hospitalizations in the United States. Of these, 53550 (21.7%) were complicated by sepsis. We noted a rising trend of SBS hospitalizations complicated by sepsis from 20.8% in 2005 to 23.5% in 2014 ($P < 0.0001$) (Table 1).

Hospitalization characteristics and clinical outcomes

Compared to non-septic SBS hospitalizations, septic SBS hospitalizations had a higher proportion of males (32.8% *vs* 29.3%, $P < 0.0001$), and patients in the 35-49 (45.9% *vs* 42.5%, $P < 0.0001$), and 50-64 (32.1% *vs* 31.1%, $P < 0.0001$) age groups. However, within the septic SBS cohort, a female predominance was noted (Table 2).

Racial disparities were also prevalent as we noted a higher proportion of ethnic minorities such as Blacks (12.4% *vs* 11.3%, $P < 0.0001$) and Hispanics (6.7% *vs* 5.5%, $P < 0.0001$) in the septic SBS cohort, while there was a higher proportion of Whites (80.0% *vs* 77.3%, $P < 0.0001$) in the non-septic cohort (Table 2).

From a hospital perspective, large (69.4% *vs* 64.6%, $P < 0.0001$), urban teaching (57.3% *vs* 51.8%, $P < 0.0001$) hospitals had a higher proportion of septic SBS hospitalizations compared to the non-septic cohort. Furthermore, there was a

Table 1 Trends for short bowel syndrome hospitalizations and septic short bowel syndrome hospitalizations in the United States from 2005-2014

Years	Short bowel syndrome hospitalizations	Septic short bowel syndrome hospitalizations	% of sepsis in short bowel syndrome hospitalizations
2005	4199	20198	20.8
2006	3908	20206	19.3
2007	4083	20206	20.2
2008	5762	26411	21.8
2009	5869	25810	22.7
2010	5640	25423	22.2
2011	5805	27647	21.0
2012	5780	27015	21.4
2013	6105	26900	22.7
2014	6400	27280	23.5
Total	53550	247097	21.7

higher proportion of emergent or urgent (19.3 *vs* 18.9%, $P < 0.0001$) septic SBS hospitalizations compared to the non-septic SBS cohort (Table 2).

The all-cause inpatient mortality was significantly higher for septic SBS hospitalizations (8.5% *vs* 1.4%, $P < 0.0001$) compared to the non-sepsis cohort. Additionally, we noted a longer mean LOS for septic SBS hospitalizations (16.1 d \pm 0.4 d *vs* 7.7 d \pm 0.1 d, $P < 0.0001$) compared to the non-septic cohort (Table 2).

Predictors of inpatient mortality for septic SBS hospitalizations

For septic SBS hospitalizations, the 18-44 age group had a 5.85 times higher risk of inpatient mortality compared to the ≥ 85 age group [odds ratio (OR): 5.85; 95% confidence interval (95%CI): 3.95-8.66, $P < 0.0001$; Table 3]. Hence, the risk of inpatient mortality decreased with increasing age. Additionally, women had a higher risk of inpatient mortality as compared to men (OR: 1.18; 95%CI: 1.02-1.38, $P = 0.03$). Race was also identified to be an independent predictor of all-cause inpatient mortality. Whites were noted to have a higher mortality risk (OR: 1.65; 95%CI: 1.17-2.33, $P = 0.005$) compared to other races. Furthermore, the presence of comorbidities such as deficiency anemias and depression were associated with a significantly higher risk of inpatient mortality for septic SBS hospitalizations (Table 3).

Intestinal transplantation and septic SBS hospitalizations

We noted a higher proportion of patients who had transplant of the intestine (TOI) for septic SBS hospitalizations (0.33% *vs* 0.22%, $P < 0.0001$) compared to the non-septic cohort (Table 1). However, due to limitations of the NIS database, we were unable to ascertain whether these hospitalizations took place within a year of undergoing TOI or later.

Racial disparities in exhauster comorbidities for septic SBS hospitalizations

A significantly higher proportion of septic SBS hospitalizations for Whites had comorbidities such as congestive heart failure, chronic pulmonary diseases, hypothyroidism, depression, and other psychiatric disorders (Table 4). Meanwhile, Blacks had significantly higher rates of comorbidities such as deficiency anemias, drug abuse, hypertension, obesity, and renal failure. Furthermore, Hispanics had the highest rates of comorbidities like coagulopathy, uncomplicated diabetes, and liver disease (Table 4).

Gender disparities in exhauster comorbidities for septic SBS hospitalizations

Septic SBS hospitalizations for women were noted to have higher rates of comorbidities such as deficiency anemias, depression, rheumatic disorders, chronic pulmonary disease, hypothyroidism, and obesity; however, men had higher rates of diabetes, drug abuse, liver disease, and renal failure (Table 5).

DISCUSSION

After surgical resection of the bowel in adults, there is a significant alteration of the nutritional, fluid, and electrolyte homeostasis, along with significant changes in the gut microbiome. These changes may lead to malabsorptive diarrhea, micronutrient deficiency, malnutrition, and SBBO[16]. Disruption of the intestinal microbiome seen in patients with SBBO impacts the production of antimicrobial peptides and immunomodulatory cells[17]. Furthermore, commensal bacteria dysbiosis may disrupt intestinal permeability leading to bacterial translocation in surrounding areas, thereby increasing

Table 2 Baseline hospitalization characteristics and clinical outcomes for septic and non-septic short bowel syndrome hospitalizations in the United States from 2005-2014

Variable	Non-septic short bowel syndrome hospitalizations	%	Septic short bowel syndrome hospitalizations	%	P value
Number of obs. (n)	193547	78.30	53550	21.70	
Age, yr (mean \pm SE)	58.0 \pm 0.3	-	57.9 \pm 0.2	-	
Age, yr					< 0.0001
18-34	41979	21.70	10000	18.70	
35-49	82308	42.50	24561	45.90	
50-64	60113	31.10	17333	32.40	
≥ 65	9149	4.70	1655	3.10	
Gender					< 0.0001
Male	56601	29.30	17545	32.80	
Female	136916	70.80	36005	67.20	
Race					< 0.0001
White	130004	80.00	35660	77.30	
Black	18436	11.30	5730	12.40	
Hispanic	8866	5.50	3091	6.70	
Others	5274	3.20	1651	3.60	
Hospital location					< 0.0001
Rural	22155	11.50	4576	8.60	
Urban nonteaching	70692	36.70	18187	34.10	
Urban teaching	99780	51.80	30509	57.30	
Hospital bed size					< 0.0001
Small	22978	11.90	5107	9.60	
Medium	45289	23.50	11219	21.10	
Large	124360	64.60	36946	69.40	
Admission type					0.0100
Elective	157070	81.20	43201	80.70	
Emergent or Urgent	36477	18.90	10349	19.30	
Intestinal Transplant	430	0.22	178	0.33	< 0.0001
Disposition status					< 0.0001
Home	156802	81.10	34439	64.40	
Facility	33898	17.50	14497	27.10	
Inpatient mortality	2724	1.40	4537	8.50	< 0.0001
LOS, d (mean \pm SE)	7.7 \pm 0.1	-	16.1 \pm 0.4	-	< 0.0001

LOS: Length of stay.

the risk of sepsis[17]. Despite known alteration of the intestine, current literature lacks data to support the routine use of antibiotics in SBS patients to prevent inflammatory gut changes[16]. Management for these patients is primarily focused on nutrition. However, patients with SBS who rely on parenteral nutrition are at increased risk of sepsis from catheter-associated infections, leading to increased hospitalizations[16]. This finding was highlighted in our study as we noted a rising trend of septic SBS hospitalizations from 20.8% in 2005 to 23.5% in 2014.

The utilization of prebiotics, probiotics, or antibiotics in SBS patients is controversial[16]. Intestinal transplantation is indicated in SBS patients with recurrent sepsis or those who are unable to receive total parenteral nutrition due to end-stage liver disease or end-stage loss of venous access[18]. Intestinal transplant significantly improves intestinal transit

Table 3 Predictors of mortality for septic short bowel syndrome hospitalizations in the United States from 2005-2014

Characteristics/co-morbidities	Odds ratio (95% confidence interval)	P value
Age (yr)		
18-44	5.85 (3.95-8.66)	< 0.0001
45-64	3.30 (2.37-4.59)	< 0.0001
65-84	1.75 (1.27-2.41)	0.001
≥ 85	Reference	
Gender		
Male	Reference	
Female	1.18 (1.02-1.38)	0.030
Race		
White	1.65 (1.17-2.33)	0.005
Black	1.28 (0.86-1.90)	0.220
Hispanic	1.22 (0.79-1.88)	0.360
Others	Reference	
Hospital bed size		
Small	1.16 (0.90-1.51)	0.250
Medium	1.10 (0.92-1.31)	0.320
Large	Reference	
Hospital type		
Rural	1.05 (0.79-1.38)	0.360
Urban non-teaching	0.94 (0.80-1.10)	0.420
Teaching	Reference	
Median household income		
Quartile 1	Reference	
Quartile 2	0.96 (0.78-1.18)	0.690
Quartile 3	1.03 (0.83-1.27)	0.800
Quartile 4	1.18 (0.94-1.47)	0.150
Co-morbidities		
Deficiency anemias	1.58 (1.35-1.85)	< 0.0001
Rheumatic disorders	1.20 (0.80-1.79)	0.380
Depression	1.66 (1.30-2.13)	< 0.0001
Drug abuse	1.51 (0.95-2.40)	0.090
Hypertension	1.16 (0.98-1.36)	0.080
Hypothyroidism	1.06 (0.83-1.34)	0.650
Lymphoma	2.48 (0.99-6.21)	0.050
Valvular disease	1.02 (0.73-1.41)	0.920

time, peristalsis, and the absorptive functions of the gut[19]. However, like any transplant, post-operative care for these patients requires lifelong immunosuppression, which imminently increases the risk for subsequent infections and sepsis. Expectedly, in our study, we noted a higher proportion of patients who had TOI in septic SBS hospitalizations compared to the non-sepsis cohort (Table 1).

Traditionally, sepsis tends to affect the elderly. However, a multicenter longitudinal cohort study in California from 2008-2015 noted that the highest overall increase in rates of sepsis and severe sepsis were for patients 18-44 years of age [20]. Although there were higher incidence rates of sepsis in the elderly, there was a notable increase in the relative risk of sepsis among young adults[20]. This was consistent with the findings in our study. In the septic SBS cohort, the 35-49 age

Table 4 Racial distribution of Elixhauser co-morbidities for septic short bowel syndrome hospitalizations in the United States from 2005-2014

Elixhauser co-morbidity	Whites	Blacks	Hispanics	Others	P value
Acquired immunodeficiency syndrome	0.0	1.0	0.3	0.0	< 0.0001
Alcohol abuse	1.6	2.4	0.8	1.4	< 0.0001
Deficiency anemias	38.8	49.0	45.4	37.6	< 0.0001
Rheumatic disorders	4.3	4.1	2.9	4.0	0.0040
Chronic blood loss anemia	2.1	2.1	2.3	3.2	0.0200
Congestive heart failure	11.2	10.8	8.6	9.8	< 0.0001
Chronic pulmonary disease	19.7	14.6	10.6	18.4	< 0.0001
Coagulopathy	15.4	16.9	19.1	16.9	< 0.0001
Depression	18.0	10.9	14.2	11.5	< 0.0001
Uncomplicated diabetes	11.4	14.4	17.7	17.0	< 0.0001
Diabetes with chronic complications	2.8	5.8	5.4	5.0	< 0.0001
Drug abuse	4.8	6.6	4.4	5.0	< 0.0001
Hypertension	32.4	42.5	34.5	34.6	< 0.0001
Hypothyroidism	13.5	10.4	8.8	8.6	< 0.0001
Liver disease	5.8	7.0	9.9	6.9	< 0.0001
Lymphoma	1.1	1.1	1.2	0.9	0.7500
Fluid and electrolyte disorders	57.5	62.5	61.2	58.2	< 0.0001
Metastatic cancer	4.5	4.4	5.2	8.6	< 0.0001
Neurological disorders	8.8	8.3	6.7	7.9	0.001
Obesity	5.6	6.3	5.7	6.9	0.0300
Paralysis	2.3	3.3	4.2	4.9	< 0.0001
Peripheral vascular disorders	6.8	7.4	7.0	6.5	0.4300
Psychiatric disorder	6.1	5.5	5.6	4.1	0.0030
Pulmonary circulation disorders	4.0	5.5	2.3	3.4	< 0.0001
Renal failure	21.2	30.7	16.7	18.8	< 0.0001
Solid tumor without metastasis	2.6	2.8	2.9	5.2	< 0.0001
Peptic ulcer disease excluding bleeding	0.1	0.1	0.1	0.0	0.3400
Valvular disease	5.2	4.8	3.8	5.3	0.0050
Weight loss	40.4	40.9	39.3	47.9	< 0.0001

group had the highest proportion of patients, while the ≥ 65 age group had the lowest proportion of patients (Table 2). On comparative analysis, septic SBS hospitalizations had a higher proportion of patients between the ages of 35-64 compared to the non-septic cohort. The exact reasons for the higher hospitalization rates of septic SBS younger patients are currently unknown, but may, in part, be due to a greater degree of awareness of sepsis in this subset population prompting them to seek immediate care or due to complications from comorbidities not previously common in this age group[20]. Nonetheless, additional prospective studies are needed to further investigate these findings.

A prospective observational cohort study of critically ill patients from 2011-2014 showed similar rates of sepsis and mortality between men and women[21]. However, there was a greater degree of endothelial cell activation in young women compared to men[21]. Increased gut permeability, often seen in patients with SBS, coupled with increased endothelial disruption of the vasculature may lead to the transportation of antigens and commensal gut microbiota from the intestine to the blood, making females even more prone to sepsis[22]. The findings of our study aligned with this current literature as females made up more than two-thirds of the total septic SBS hospitalizations in the United States.

The study by Siddiqui *et al*[2] reported that Whites made up 78% of all SBS hospitalizations in the United States. We report similar findings as septic SBS hospitalizations had a higher proportion of Whites (77.3%) compared to other ethnic minorities such as Blacks or Hispanics (Table 2). However, on comparative analysis, septic SBS hospitalizations had a higher proportion of Blacks (12.4% *vs* 11.3%, $P < 0.0001$) and Hispanics (6.7% *vs* 5.5%, $P < 0.0001$) compared to the non-

Table 5 Gender distribution of elixhauser co-morbidities for septic short bowel syndrome hospitalizations in the United States from 2005-2014

Elixhauser co-morbidity	Male	Female	P value
Acquired immunodeficiency syndrome	0.3	0.1	< 0.0001
Alcohol abuse	2.9	1.0	< 0.0001
Deficiency anemias	36.8	39.7	< 0.0001
Rheumatic disorders	2.0	5.1	< 0.0001
Chronic blood loss anemia	1.9	2.1	0.0800
Congestive heart failure	10.7	10.8	0.9000
Chronic pulmonary disease	16.7	19.0	< 0.0001
Coagulopathy	16.8	15.0	< 0.0001
Depression	12.4	18.1	< 0.0001
Uncomplicated diabetes	12.6	11.6	0.0010
Diabetes with chronic complications	3.9	2.7	< 0.0001
Drug abuse	5.1	4.5	0.0010
Hypertension	33.6	32.1	0.0010
Hypothyroidism	6.0	14.9	< 0.0001
Liver disease	7.4	5.3	< 0.0001
Lymphoma	1.4	0.9	< 0.0001
Fluid and electrolyte disorders	56.6	57.9	0.0030
Metastatic cancer	4.9	4.4	0.0020
Neurological disorders	7.0	8.8	< 0.0001
Obesity	3.8	6.4	< 0.0001
Paralysis	3.5	2.1	< 0.0001
Peripheral vascular disorders	8.1	5.9	< 0.0001
Psychiatric disorder	4.7	6.3	< 0.0001
Pulmonary circulation disorders	4.2	3.8	0.0400
Renal failure	26.3	18.8	< 0.0001
Solid tumor without metastasis	2.6	2.7	0.5900
Peptic ulcer disease excluding bleeding	0.0	0.1	< 0.0001
Valvular disease	5.2	4.7	0.0100
Weight loss	39.1	40.9	< 0.0001

septic cohort. This may, in part, be due to lack of healthcare facilities, and awareness about SBS among ethnic minorities leading to a progression of their disease and further complications by sepsis. Hence, we advocate for the need for urgent interventions to improve healthcare access, increase awareness about sepsis in SBS, and improve outpatient follow-up for these high-risk populations.

The all-cause inpatient mortality for SBS hospitalizations was found to be 1.4%. However, when these hospitalizations were complicated by sepsis, the all-cause inpatient mortality increased to 8.5%. This was in line with current literature which reports increased inpatient mortality for septic SBS hospitalizations[2]. Moreover, the presence of a greater number of comorbidities is also associated with higher rates of mortality in patients with sepsis[23]. A cohort study evaluating adult sepsis survivors identified age, sex, race, severe comorbidities, and site of initial infection to be long-term risk factors for mortality[24]. Similarly, in our study, independent risk factors that increased inpatient mortality for septic SBS hospitalizations are as follows.

Age

In our study, the risk of inpatient mortality was almost six times higher in the 18-44 age group compared to individuals ≥ 85 years of age. In 2013, a Quality and Cost of Primary Care study reported that younger patients were more likely to visit a healthcare provider for both mental and physical health conditions compared to the elderly population[25]. The

increased health awareness, highly accurate provider evaluation in younger adults without non-specific baseline symptoms, and presence of additional comorbidities which were previously not common in this age group may have led to increased diagnosis and associated inpatient mortality.

Gender

In 2011, a prospective clinical trial of intensive care unit patients reported that females with sepsis had higher mortality rates than males with sepsis[26]. Another study by Wilcox *et al*[27] noted similar gender outcomes. Similarly, in our study, septic SBS female patients had higher rates of all-cause inpatient mortality compared to males. Furthermore, we noted higher rates of deficiency anemias and depression in females compared to males, which was also independently associated with higher mortality rates in septic SBS hospitalizations.

Race

We noted that White septic SBS hospitalizations had a 1.65 times greater risk of inpatient mortality compared to other races. A study of Caucasians with acute respiratory distress syndrome (ARDS) secondary to pneumonia evaluated the FER rs4957796 TT genotype as a means of determining 90-d mortality risk[28]. Although this study specifically assessed ARDS patients and their survival in the setting of pneumonia, interestingly the FER gene is known to play a key role in the regulation of intestinal barrier function[29]. White septic SBS hospitalizations may have a higher mortality risk due to the absence of FER protein which may exacerbate intestinal dysfunction and increase bacterial translocation into the bloodstream leading to a greater severity of sepsis. However, additional prospective studies are needed to further validate our findings.

Comorbidities

In our study, deficiency anemias and depression were found to coincide with significantly greater mortality risk in septic SBS hospitalizations. Deficiency anemias are highly prevalent in SBS patients due to altered anatomy[30-33]. In septic SBS hospitalizations, deficiency anemia can negatively impact the host's defense and immunomodulatory response, increasing the severity of sepsis and overall mortality risk. Furthermore, severe depression has a known association with increased BSI, leading to sepsis and higher mortality rates[34,35]. Depression ultimately leads to decreased immune function and disruption of the brain-gut-microbiome axis, which may increase the host's risk for sepsis and adverse clinical outcomes.

Limitations

A key strength of this study is the study population which has been derived from one of the biggest, national, diverse, multi-ethnic databases in the United States. Through our analysis over 10 years, we are able to provide meaningful information on the trends of septic SBS hospitalizations. Furthermore, we also perform a unique comparative analysis between septic and non-septic SBS hospitalizations and identify predictors of inpatient mortality for septic SBS hospitalizations to give gastroenterologists real-world data on the patients at the highest risk of adverse clinical outcomes. However, we do acknowledge all the limitations associated with our study. We were unable to perform a detailed analysis after the 2014 study period as the NIS changed from ICD-9-CM to ICD-10-CM coding at the beginning of October 1, 2015. Converting ICD-9 to ICD-10 is an extremely challenging process as there are differences in the structure and granularity of codes. Hence, it would be impossible to find the exact matches for the codes with a high level of confidence. Additionally, the NIS database lacks information on the time from hospitalization to diagnosis of sepsis, hospital course, treatment aspects, inpatient procedures, and pharmacological aspects of management. Lastly, the NIS is an administrative database. Therefore, the possibility of coding errors cannot be excluded. Despite these limitations, we believe that our study helps fill the current knowledge gaps for SBS hospitalizations complicated by sepsis in the US as this entity has not been studied extensively. We hope that the findings of our study can serve as a foundation for future prospective studies and randomized controlled trials.

CONCLUSION

In conclusion, SBS is a well-known complication of surgical resection of the intestine. Due to their dependence on parenteral nutrition using indwelling venous catheters, these patients are at increased risk of BSI and sepsis from bacterial translocation from the gut to the bloodstream during enteral feeding. In the United States from 2005-2014, we noted a rising trend of septic SBS hospitalizations with a significant female predominance. Compared to the non-septic cohort, septic SBS hospitalizations had a higher proportion of patients with TOI, higher all-cause inpatient mortality, and longer mean LOS. Independent predictors of mortality for septic SBS hospitalizations included White race, female gender, younger age, and those with associated comorbidities such as deficiency anemias and depression.

FOOTNOTES

Author contributions: Dahiya DS, Wachala J, Solanki S, and Jafri SM contributed to the conception and design; Dahiya DS, Solanki S, Kichloo A, and Jafri SM contributed to the administrative support; Dahiya DS, Wachala J, Solanki S, Solanki D, Kichloo A, Holcomb S, Mansuri U, Haq KS, Ali H, Gangwani MK, and Shah YR contributed to the provision, collection, and assembly of data; Dahiya DS,

Wachala J, Solanki S, Solanki D, Kichloo A, Holcomb S, Mansuri U, Haq KS, Ali H, Gangwani MK, Shah YR, Varghese T, Khan HMA, Horslen S, Schiano TD, and Jafri SM contributed to the review of literature and drafting the manuscript; Dahiya DS, Wachala J, Solanki S, Solanki D, Kichloo A, Holcomb S, Mansuri U, Haq KS, Ali H, Gangwani MK, Shah YR, Varghese T, Khan HMA, Horslen S, Schiano TD, and Jafri SM contributed to the revision of key components of the manuscript and final approval of manuscript; Dahiya DS, Wachala J, Solanki S, Solanki D, Kichloo A, Holcomb S, Mansuri U, Haq KS, Ali H, Gangwani MK, Shah YR, Varghese T, Khan HMA, Horslen S, Schiano TD, and Jafri SM are accountable for all aspects of the work.

Institutional review board statement: The National Inpatient Sample lacks specific patient and hospital identifiers. Hence, our analysis did not require institutional review board (IRB) approval as per guidelines put forth by our institutional IRB for the analysis of national databases.

Informed consent statement: The National Inpatient Sample lacks specific patient identifiers. Hence, informed consent was not required.

Conflict-of-interest statement: Authors declare no conflict of interests for this article.

Data sharing statement: The National Inpatient Sample (NIS) is a large, publicly available, all-payer inpatient care database in the United States containing data on millions of hospital stays per year. The large sample size derived from the NIS database provides sufficient data for analysis of common diseases, uncommon disorders, and procedures. The NIS is publicly available at: <https://www.hcup-us.ahrq.gov/>

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <https://creativecommons.org/licenses/by-nc/4.0/>

Country/Territory of origin: United States

ORCID number: Dushyant Singh Dahiya 0000-0002-8544-9039; Shantanu Solanki 0000-0001-9563-2470; Hassam Ali 0000-0001-5546-9197; Manesh Kumar Gangwani 0000-0002-3931-6163; Thomas D Schiano 0000-0003-1878-5101; Syed-Mohammed Jafri 0000-0001-8108-7408.

S-Editor: Chen YL

L-Editor: A

P-Editor: Zhao YQ

REFERENCES

- Massironi S, Cavalcoti F, Rausa E, Invernizzi P, Braga M, Vecchi M. Understanding short bowel syndrome: Current status and future perspectives. *Dig Liver Dis* 2020; **52**: 253-261 [PMID: 31892505 DOI: 10.1016/j.dld.2019.11.013]
- Siddiqui MT, Al-Yaman W, Singh A, Kirby DF. Short-Bowel Syndrome: Epidemiology, Hospitalization Trends, In-Hospital Mortality, and Healthcare Utilization. *JPEN J Parenter Enteral Nutr* 2021; **45**: 1441-1455 [PMID: 33233017 DOI: 10.1002/jpen.2051]
- Kurkchubasche AG, Smith SD, Rowe MI. Catheter sepsis in short-bowel syndrome. *Arch Surg* 1992; **127**: 21-4; discussion 24 [PMID: 1734846 DOI: 10.1001/archsurg.1992.01420010027004]
- Miko BA, Kamath SS, Cohen BA, Jeon C, Jia H, Larson EL. Epidemiologic Associations Between Short-Bowel Syndrome and Bloodstream Infection Among Hospitalized Children. *J Pediatric Infect Dis Soc* 2015; **4**: 192-197 [PMID: 26336089 DOI: 10.1093/jpids/piu079]
- Weber TR. Enteral feeding increases sepsis in infants with short bowel syndrome. *J Pediatr Surg* 1995; **30**: 1086-8; discussion 1088 [PMID: 7472938 DOI: 10.1016/0022-3468(95)90347-x]
- Terra RM, Plopper C, Waitzberg DL, Cukier C, Santoro S, Martins JR, Song RJ, Gama-Rodrigues J. Remaining small bowel length: association with catheter sepsis in patients receiving home total parenteral nutrition: evidence of bacterial translocation. *World J Surg* 2000; **24**: 1537-1541 [PMID: 11193720 DOI: 10.1007/s002680010274]
- Healthcare Cost and Utilization Project (HCUP). Overview of National (Nationwide) Inpatient Sample (NIS). [cited 1 March 2024]. Available from: <https://hcup-us.ahrq.gov/nisoverview.jsp>
- Healthcare Cost and Utilization Project (HCUP). NIS Database Documentation. Volume 2020: Healthcare Cost and Utilization Project (HCUP). [cited 1 March 2024]. Available from: <https://hcup-us.ahrq.gov/nisoverview.jsp>
- Hoffman RD, Danos DM, Lau FH. National health disparities in incisional hernia repair outcomes: An analysis of the Healthcare Cost and Utilization Project National Inpatient Sample (HCUP-NIS) 2012-2014. *Surgery* 2021; **169**: 1393-1399 [PMID: 33422347 DOI: 10.1016/j.surg.2020.11.028]
- Healthcare Cost and Utilization Project (HCUP). Healthcare Cost and Utilization Project Data Use Agreement Course. [cited 1 March 2024]. Available from: https://hcup-us.ahrq.gov/tech_assist/dua.jsp
- Limketkai BN, Parian AM, Chen PH, Colombel JF. Treatment With Biologic Agents Has Not Reduced Surgeries Among Patients With Crohn's Disease With Short Bowel Syndrome. *Clin Gastroenterol Hepatol* 2017; **15**: 1908-1914.e2 [PMID: 28666947 DOI: 10.1016/j.cgh.2017.06.040]
- Papalos A, Narula J, Bavishi C, Chaudhry FA, Sengupta PP. U.S. Hospital Use of Echocardiography: Insights From the Nationwide Inpatient Sample. *J Am Coll Cardiol* 2016; **67**: 502-511 [PMID: 26846948 DOI: 10.1016/j.jacc.2015.10.090]
- Mansuri U, Patel A, Shah H, Chauhan K, Poojary P, Saha A, Dave M, Hazra A, Mishra T, Annapureddy N, Nadkarni G, Chan L. Trends and outcomes of sepsis hospitalizations complicated by acute kidney injury requiring hemodialysis. *J Crit Care* 2017; **38**: 353-355 [PMID: 27986342 DOI: 10.1016/j.jcrc.2016.10.012]

- 14 **Healthcare Cost and Utilization Project (HCUP).** Elixhauser Comorbidity Software, Version 3.7. Volume 2021. [cited 1 March 2024]. Available from: <https://hcup-us.ahrq.gov/toolssoftware/comorbidity/comorbidity.jsp>
- 15 **Healthcare Cost and Utilization Project (HCUP).** Trend Weights for HCUP NIS Data. Volume 2020. [cited 1 March 2024]. Available from: <https://hcup-us.ahrq.gov/db/nation/nis/trendwghts.jsp>
- 16 **Parrish CR, DiBaise JK.** Managing the Adult Patient With Short Bowel Syndrome. *Gastroenterol Hepatol (N Y)* 2017; **13**: 600-608 [PMID: 29230136]
- 17 **Fay KT, Ford ML, Coopersmith CM.** The intestinal microenvironment in sepsis. *Biochim Biophys Acta Mol Basis Dis* 2017; **1863**: 2574-2583 [PMID: 28286161 DOI: 10.1016/j.bbadis.2017.03.005]
- 18 **Höllwarth ME.** Surgical strategies in short bowel syndrome. *Pediatr Surg Int* 2017; **33**: 413-419 [PMID: 28039510 DOI: 10.1007/s00383-016-4043-6]
- 19 **Kaufman SS, Avitzur Y, Beath SV, Ceulemans LJ, Gondolesi GE, Mazariegos GV, Pironi L.** New Insights Into the Indications for Intestinal Transplantation: Consensus in the Year 2019. *Transplantation* 2020; **104**: 937-946 [PMID: 31815899 DOI: 10.1097/TP.0000000000003065]
- 20 **Wardi G, Tainter CR, Ramnath VR, Brennan JJ, Tolia V, Castillo EM, Hsia RY, Malhotra A, Schmidt U, Meier A.** Age-related incidence and outcomes of sepsis in California, 2008-2015. *J Crit Care* 2021; **62**: 212-217 [PMID: 33429114 DOI: 10.1016/j.jcrc.2020.12.015]
- 21 **van Vught LA, Scicluna BP, Wiewel MA, Hoogendijk AJ, Klein Klouwenberg PMC, Ong DSY, Cremer OL, Horn J, Frantza M, Toliat MR, Nürnberg P, Bonten MMJ, Schultz MJ, van der Poll T; MARS Consortium.** Association of Gender With Outcome and Host Response in Critically Ill Sepsis Patients. *Crit Care Med* 2017; **45**: 1854-1862 [PMID: 28806220 DOI: 10.1097/CCM.0000000000002649]
- 22 **Assimakopoulos SF, Triantos C, Thomopoulos K, Fligou F, Maroulis I, Marangos M, Gogos CA.** Gut-origin sepsis in the critically ill patient: pathophysiology and treatment. *Infection* 2018; **46**: 751-760 [PMID: 30003491 DOI: 10.1007/s15010-018-1178-5]
- 23 **Rhee C, Jones TM, Hamad Y, Pande A, Varon J, O'Brien C, Anderson DJ, Warren DK, Dantes RB, Epstein L, Klompas M; Centers for Disease Control and Prevention (CDC) Prevention Epicenters Program.** Prevalence, Underlying Causes, and Preventability of Sepsis-Associated Mortality in US Acute Care Hospitals. *JAMA Netw Open* 2019; **2**: e187571 [PMID: 30768188 DOI: 10.1001/jamanetworkopen.2018.7571]
- 24 **Shankar-Hari M, Harrison DA, Ferrando-Vivas P, Rubenfeld GD, Rowan K.** Risk Factors at Index Hospitalization Associated With Longer-term Mortality in Adult Sepsis Survivors. *JAMA Netw Open* 2019; **2**: e194900 [PMID: 31150081 DOI: 10.1001/jamanetworkopen.2019.4900]
- 25 **Thompson AE, Anisimowicz Y, Miedema B, Hogg W, Wodchis WP, Aubrey-Bassler K.** The influence of gender and other patient characteristics on health care-seeking behaviour: a QUALICOPC study. *BMC Fam Pract* 2016; **17**: 38 [PMID: 27036116 DOI: 10.1186/s12875-016-0440-0]
- 26 **Nachtigall I, Tafelski S, Rothbart A, Kaufner L, Schmidt M, Tamarkin A, Kartachov M, Zebedies D, Trefzer T, Wernecke KD, Spies C.** Gender-related outcome difference is related to course of sepsis on mixed ICUs: a prospective, observational clinical study. *Crit Care* 2011; **15**: R151 [PMID: 21693012 DOI: 10.1186/cc10277]
- 27 **Elizabeth Wilcox M, Donnelly JP, Lone NI.** Understanding gender disparities in outcomes after sepsis. *Intensive Care Med* 2020; **46**: 796-798 [PMID: 32072302 DOI: 10.1007/s00134-020-05961-3]
- 28 **Hinz J, Büttner B, Kriesel F, Steinau M, Frederik Popov A, Ghadimi M, Beissbarth T, Tzvetkov M, Bergmann I, Mansur A.** The FER rs4957796 TT genotype is associated with unfavorable 90-day survival in Caucasian patients with severe ARDS due to pneumonia. *Sci Rep* 2017; **7**: 9887 [PMID: 28851893 DOI: 10.1038/s41598-017-08540-7]
- 29 **Qi W, Ebbert KV, Craig AW, Greer PA, McCafferty DM.** Absence of Fer protein tyrosine kinase exacerbates endotoxin induced intestinal epithelial barrier dysfunction in vivo. *Gut* 2005; **54**: 1091-1097 [PMID: 16009680 DOI: 10.1136/gut.2004.061887]
- 30 **Liu Q, Wu J, Zhang X, Wu X, Zhao Y, Ren J.** Iron homeostasis and disorders revisited in the sepsis. *Free Radic Biol Med* 2021; **165**: 1-13 [PMID: 33486088 DOI: 10.1016/j.freeradbiomed.2021.01.025]
- 31 **Sun H, Weaver CM.** Decreased Iron Intake Parallels Rising Iron Deficiency Anemia and Related Mortality Rates in the US Population. *J Nutr* 2021; **151**: 1947-1955 [PMID: 33834234 DOI: 10.1093/jn/nxab064]
- 32 **Alker W, Haase H.** Zinc and Sepsis. *Nutrients* 2018; **10** [PMID: 30060473 DOI: 10.3390/nu10080976]
- 33 **Wolffenbuttel BHR, Heiner-Fokkema MR, Green R, Gans ROB.** Relationship between serum B12 concentrations and mortality: experience in NHANES. *BMC Med* 2020; **18**: 307 [PMID: 33032595 DOI: 10.1186/s12916-020-01771-y]
- 34 **Askim Å, Gustad LT, Paulsen J, Reitan SK, Mehl A, Mohus RM, Dewan A, Damås JK, Solligård E, Åsvold BO.** Anxiety and Depression Symptoms in a General Population and Future Risk of Bloodstream Infection: The HUNT Study. *Psychosom Med* 2018; **80**: 673-679 [PMID: 29923889 DOI: 10.1097/PSY.0000000000000619]
- 35 **Thorkildsen MS, Laugsand LE, Nilsen TIL, Mohus RM, Høvik LH, Rogne T, Solligård E, Damås JK, Gustad LT.** Insomnia symptoms and risk of bloodstream infections: prospective data from the prospective population-based Nord-Trøndelag Health Study (HUNT), Norway. *J Sleep Res* 2023; **32**: e13696 [PMID: 36068650 DOI: 10.1111/jsr.13696]



Published by **Baishideng Publishing Group Inc**
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA

Telephone: +1-925-3991568

E-mail: office@baishideng.com

Help Desk: <https://www.f6publishing.com/helpdesk>

<https://www.wjgnet.com>

