



Pivotal role of exosomes in diagnosis and treatment of esophageal cancer in a new era of precision medicine

Grigorios Christodoulidis, Konstantinos Eleftherios Koumarelas, Marina Nektaria Kouliou

Specialty type: Oncology

Provenance and peer review:

Invited article; Externally peer reviewed.

Peer-review model: Single blind

Peer-review report's scientific quality classification

Grade A (Excellent): 0
Grade B (Very good): 0
Grade C (Good): 0
Grade D (Fair): 0
Grade E (Poor): 0

P-Reviewer: Zhang X, China

Received: December 8, 2023

Peer-review started: December 8, 2023

First decision: December 21, 2023

Revised: December 23, 2023

Accepted: January 17, 2024

Article in press: January 17, 2024

Published online: March 20, 2024



Grigorios Christodoulidis, Konstantinos Eleftherios Koumarelas, Marina Nektaria Kouliou,
Department of General Surgery, University Hospital of Larissa, Larissa 41110, Greece

Corresponding author: Grigorios Christodoulidis, MD, PhD, Surgeon, Department of General Surgery, University Hospital of Larissa, Mezourlo, Larissa 41110, Greece. gregsurg@yahoo.gr

Abstract

In this editorial we comment on the article published by Ning *et al*, "Role of exosomes in metastasis and therapeutic resistance in esophageal cancer". Esophageal cancer (EC) represents a significant global health concern, being the seventh most common and sixth in terms of mortality worldwide. Despite the advances in therapeutic modalities, the management of patients with EC remains challenging, with a 5-year survival rate of only 25% and a limited eligibility for curative surgery due to its late diagnosis. Conventional screening methods are impractical for the early detection of EC, given their either invasive or insensitive nature. The advent of liquid biopsy, with a focus on circulating tumor cells, circulating tumor DNA, and exosomes, heralds a non-invasive avenue for cancer detection. Exosomes, small vesicles involved in intercellular communication, are highlighted as potential biomarkers for EC diagnosis and prognosis. Along with a diverse cargo encompassing various types of RNA, DNA molecules, proteins, and metabolites, exosomes emerge as key players in tumorigenesis, tumor development, and metastasis. Their significance extends to carrying distinctive biomarkers, including microRNAs (miRNAs), long non-coding RNAs, and circular RNAs, underscoring their potential diagnostic and prognostic value. Furthermore, exosomes may be utilized for therapeutic purposes in the context of EC treatment, serving as efficient delivery vehicles for therapeutic agents such as chemotherapeutic medicines and miRNAs. In this editorial we delve into the applications of exosomes for the early detection and treatment of EC, as well as the future perspectives.

Key Words: Exosomes; Esophageal cancer; Diagnostic methods; Novel therapies

©The Author(s) 2024. Published by Baishideng Publishing Group Inc. All rights reserved.

Core Tip: Esophageal cancer (EC) is a global health concern ranking seventh in incidence and sixth in mortality worldwide, with over 604000 new cases and 544000 deaths in 2020. Despite advancements in therapies, nearly half of EC patients experience distant metastasis or therapeutic resistance, resulting in a 5-year survival rate below 25%. Traditional screening methods are limited, necessitating the need for non-invasive alternatives. Liquid biopsy, particularly focusing on exosomes, emerges as a promising option for early EC detection. Exosomes, small vesicles for intercellular communication, carry diverse biomarkers and play a crucial role in tumorigenesis. Notably, exosomal microRNAs, long non-coding RNAs, and circular RNAs show potential as diagnostic and prognostic markers for esophageal squamous cell carcinoma. Beyond diagnosis, exosomes serve as effective delivery tools for therapeutic agents, exhibiting advantages in immunotherapy, gene therapy, and drug delivery, presenting a multifaceted approach in the battle against EC.

Citation: Christodoulidis G, Koumarelas KE, Kouliou MN. Pivotal role of exosomes in diagnosis and treatment of esophageal cancer in a new era of precision medicine. *World J Methodol* 2024; 14(1): 90624

URL: <https://www.wjgnet.com/2222-0682/full/v14/i1/90624.htm>

DOI: <https://dx.doi.org/10.5662/wjm.v14.i1.90624>

INTRODUCTION

Esophageal cancer represents a prevalent malignancy in the digestive system, ranking seventh globally in incidence and sixth in mortality, accounting for over 604000 new cases and more than 544000 deaths worldwide in 2020[1-5]. This malignancy predominantly manifests in two histological subtypes: Esophageal adenocarcinoma (EAC) and esophageal squamous cell carcinoma (ESCC). EAC typically arises in the lower part of the esophagus and is often associated with chronic gastroesophageal reflux disease (GERD). It is characterized by changes in the lining of the esophagus, known as Barrett's esophagus. On the other hand, ESCC typically originates in the upper or middle part of the esophagus and is closely linked to factors such as tobacco and alcohol consumption[1,4]. ESCC primarily affects people in East and Central Asia, while EAC is presented more in Western Europe and North America. Notably, China with its high prevalence, contributes to over 50% of newly diagnosed EC cases worldwide[1-3].

Despite advancements in surgical and systemic drug therapies, EC poses considerable challenges, with nearly half of patients experiencing distant metastasis or therapeutic resistance post-treatment. The 5-year survival rate for patients remains below 25%, with a median survival time ranging from 13.6 to 19.3 mo. Surgery stands as the cornerstone treatment, yet only around 30% of newly diagnosed patients are eligible for curative resection[1-3,5]. Presently, there is a scarcity of effective biomarkers for early-stage ESCC detection. Moreover, a significant percentage of patients experience loco-regional recurrence following surgical resection aimed at cure. While adjuvant radiotherapy and chemotherapy hold importance for ESCC, their clinical efficacy remains a subject of debate, emphasizing the critical need for earlier diagnosis [1,3,5].

Endoscopic examination with biopsy and imaging studies, while being two of the main screening methods for EC, are invasive and insensitive, respectively. Emerging minimally invasive technologies like cytosponge or transnasal endoscopy face barriers related to cost and discomfort, limiting their widespread acceptance as screening methods for ESCC[5].

Liquid biopsy, which focuses on circulating tumor cells (CTCs), circulating tumor DNA (ctDNA), and extracellular vesicles (EVs), particularly exosomes, is a potential early detection method[2,5]. Exosomes, microvesicles, and apoptotic bodies are three types of EVs, with exosomes having dimensions ranging from 30 to 150 nm and playing an important role in intercellular communication. Exosomes carry a diverse cargo of RNA types (miRNAs, regulatory mRNAs, piwi-interacting RNA, small nucleolar RNA, tRNA-derived small RNAs (tsRNAs), long noncoding RNAs (lncRNAs), and unidentified small RNAs), DNA sequences, proteins, lipids, glycoconjugates, and metabolites, and are present in multiple biological fluids, such as blood, saliva, urine, malignant ascites, and cerebrospinal fluid, offering a non-invasive avenue for cancer detection compared to methods requiring phlebotomy, like ctDNA and CTCs[1-6].

Recent studies underscore the significant role of exosomes in tumorigenesis, development, and metastasis, and they participate in processes such as angiogenesis, extracellular matrix remodeling, and miRNA transfer[1,3]. Furthermore, exosomes harbor lncRNAs, pivotal in intercellular material exchange, signal transduction, and regulation of crucial oncological behaviors. The dysregulation of exosomal lncRNAs shows promise as diagnostic and prognostic biomarkers across various cancers[3].

In-depth comprehension of esophageal cancer epidemiology, subtypes, and associated biomarkers, with a particular focus on the role of exosomes, presents invaluable insights for advancing early detection, prognostication, and therapeutic strategies, aiming for enhanced outcomes among EC patients.

EXOSOMES AND DIAGNOSIS

Exosomes play pivotal roles in various stages of the tumor development cascade, encompassing tumor proliferation, angiogenesis, epithelial-mesenchymal transition, migration, microenvironment remodeling, and therapeutic resistance[1, 2,7,8]. The early diagnosis and accurate prediction of therapy outcomes using exosomes have the potential to significantly

enhance the prognosis of patients with ESCC.

Exosomes emerge as potent cancer biomarkers due to their ease of extraction from most biofluids, exceptional stability, and capacity to convey dynamic information regarding the tumor state[1,2]. Zhao *et al*[3] observed a notable upregulation in the levels of circulating exosomes (CEs) from ESCC patients, which demonstrated a sensitivity of 75% and specificity of 85% in distinguishing ESCC patients from healthy individuals. The elevated level of CE independently serves as a prognostic marker for ESCC patients.

Exosomal miRNAs, including miR-652-5p, miRNA-21, miR-766-3p, and miR-182, exhibit potential as diagnostic and prognostic biomarkers for ESCC. Particularly, miR-182 depletion in the postoperative period renders it a promising biomarker post-surgery. The serum miR-25/miR-203 ratio is significantly higher in ESCC patients, presenting potential for monitoring the effect of tumor resection and recurrence (sensitivity = 71.9%, specificity = 96.6%). Following surgery, the expression level of miR-25/miR-203 significantly decreases, as indicated by the receiver operating characteristic curve (sensitivity = 97.4%, specificity = 65.8%)[2,5,9-11].

Exosomal Dicer assumes a significant role in miRNA synthesis; thus, monitoring exosomal Dicer may prove more efficient than miRNA for diagnosing esophageal cancer[2]. Small RNAs derived from tRNA (tsRNAs), such as tRNA-GlyGCC-5 and sRESE in exosomes, display potential as pre-operative biomarkers for ESCC, predicting prognosis and response to adjuvant therapy. The combined sensitivity of tRNA-GlyGCC-5 and sRESE reaches 90.5%, with a specificity of 94.2%[5].

lncRNAs are extensively investigated as crucial biomarkers in various cancer types. Exosomal related-lncRNAs (ER-lncRNAs), including AC082651.3, AP000487.1, PLA2G4E-AS1, C8orf49, and AL356056.2, are identified as potential markers for ESCC, diminishing after surgery. ER-lncRNA pairs exhibit superior predictive value compared to traditional clinical indicators for median survival time[3].

ER-lncRNAs, such as PCAT1, UCA1, POU3F3, ESCCAL-1, and PEG10, are promising biomarkers for ESCC diagnosis and prognosis. Specifically, the lncRNA UCA1 serves as a potent diagnostic marker with a sensitivity and specificity of 86.7% and 70.2%, respectively. These lncRNAs, as a diagnostic panel, provide an accurate diagnosis of early esophageal cancer[12]. Candidate lncRNAs, including AC098818.2, RASSF8-AS1, LINC00958, GMDS-DT, and AL591721.1, are significantly overexpressed in ESCC patients' blood serum samples and cancerous tissues compared to healthy donors or other cancer types, indicating increased specificity for screening[13]. The expression levels of lncRNAs NR_039819, NR_036133, NR_003353, ENST00000442416.1, and ENST00000416100.1 increase in patients with ESCC and decrease after surgery. Serum lncRNA RASSF8-AS1 has been identified as the most reliable diagnostic marker for ESCC[14].

CircRNAs also exhibit potential as diagnostic biomarkers. Has-circ-0001946 and has-circ-0043603, secreted by ESCC cells, may serve as diagnostic biomarkers for ESCC, with combined detection showing improved accuracy compared to single detection. Serum exosomal has-circ-0026611 emerges as a novel predictor of ESCC prognosis[15].

Exosomes, as carriers of various types of non-coding RNAs and a multitude of molecules, present a comprehensive landscape for potential biomarkers. Zhu *et al* investigated four exosomal metabolites in ESCC patients, revealing a marker panel, including 3'-UMP, palmitoleic acid, palmitaldehyde, and isobutyl decanoate, with excellent diagnostic performance (area under the curve = 0.98) for predicting ESCC recurrence. Among these, 3'-UMP is identified as the most crucial for diagnosis[4]. Rao *et al*[16] observed that intercellular adhesion molecule-1 (CD54) is upregulated in cancer tissues as well as the exosomal CD54, thus making it a great option for diagnostic biomarker. Utilizing exosomal CD54 as a biomarker, they observed a sensitivity and specificity of 66.13% and 71.31%, respectively. Exosomes exhibit advantages over other cancer biomarkers, such as circulating tumor cells and circulating tumor DNA, owing to their ample quantity, robust stability, and ease of accessibility.

UTILITY OF EXOSOMES IN TREATMENT

Exosomes serve as delivery tools, transporting miRNAs, mRNAs, lncRNAs, and proteins selectively to target cells. They possess unique natural advantages including immune-escape, easy penetration of cell membranes, and specific recognition by receptor cells. Their low immunogenicity, high biocompatibility, long circulating half-life, less toxicity, and the ability to cross biological barriers, render them potent vectors for therapeutic agents[1,6,12].

Exosomes encapsulate various therapeutic agents, such as chemotherapeutic medicines, proteins, siRNAs, lncRNAs, and miRNAs. Exosome-derived miR-154-5p attenuates the invasion of EC cells and inhibits their angiogenic capability *in vitro*, curbing the malignant progression of ESCC. The exosomal lncRNA UCA1 is observed to inhibit EC progression, and miR-339-5p transferred *via* exosomes induces radiosensitivity in ESCC cells. Certain exosomal miRNAs, like miR-339-5p, promote sensitivity to radiation therapy, downregulating Cdc25A. Thus, high levels of miR-339-5p in tumor tissues and serum indicate a good prognosis. Exosomal derived miR-375 promotes apoptosis and suppresses proliferation, invasion, and migration of tumor cells, explaining its potential in tumor treatment[1,17,18]. Exosomal miR-19b-3p transferred to cells reverses the inhibitory effect of PTEN overexpression on cell invasion. PTEN overexpression downregulates MMP-2 and vimentin, and upregulates E-cadherin. Thus, PTEN upregulation through exosomal miRNA seems promising[12,19]. Exosomes play a significant role in gene therapy, and exosomal ECRG4 mRNA is a potential target for EC gene therapy, suppressing cell proliferation *in vitro* and inhibiting tumor growth *in vivo*. Moreover, exosomes can exert antitumor effects from immune cells by displaying specific surface antigens, activating T cells vital for tumor immunity[2]. Specifically, engineered M1 macrophage-derived exosomes inhibit tumor growth and transform M2-type tumor-associated macrophages into M1-like macrophages[1]. Exosomes are also superior drug delivery systems compared to liposomes, polymers, *etc.*, due to their biocompatibility, biodegradability, and better target specificity. Adriamycin- and paclitaxel-loaded exosomes exhibit low immunogenicity and toxicity, improving efficacy in treating

multidrug-resistant cancer cells[1,12]. Engineered exosomes, with specific molecules attached, enhance targeting, increase production, and offer various lipid and protein compositions. The utilization of exosomes in immunotherapy is an important aspect. Dendritic cell-derived exosomes loaded with tumor antigens induce anti-tumor immune responses, showing potential in cancer therapy. Exosomes may play a role in inducing senescence, representing a promising strategy for ESCC treatment.

CONCLUSION

EC is a major worldwide health issue, with high incidence and mortality. At present, the only definitive therapy is surgery, but by the time the tumor is diagnosed, it is at an advanced stage. Early identification and diagnosis may be the key to a better prognosis, and exosomal miRNAs, lncRNAs, and circular RNAs show promise as diagnostic and prognostic indicators for ESCC. Exosomes are an appealing option for enhancing the accuracy of early diagnosis and prognosis prediction due to their ease of extraction, stability, and ability to communicate dynamic information about the tumor status. Furthermore, the utilization of exosomes in the therapy of EC as a therapeutic drug delivery method, is under investigation. Their unique natural advantages, including immune-escape and specific recognition by receptor cells, rendering them as potent vectors for therapeutic interventions. In summary, the potential of exosomes in early diagnosis and as carriers of therapeutic agents opens new avenues for precision medicine in the management of esophageal cancer.

FOOTNOTES

Author contributions: Christodoulidis G, Konstantinos-Eleftherios K, and Marina-Nektaria K contributed to the preparation of this paper; Christodoulidis G designed the overall concept and outline of the manuscript; Christodoulidis G, Konstantinos-Eleftherios K, and Marina-Nektaria K contributed to the discussion and design of the manuscript; Christodoulidis G, Konstantinos-Eleftherios K, and Marina-Nektaria K contributed to the writing and editing the manuscript, and review of the literature.

Conflict-of-interest statement: All the authors declare that they have no conflict of interest to disclose.

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <https://creativecommons.org/licenses/by-nc/4.0/>

Country/Territory of origin: Greece

ORCID number: Grigorios Christodoulidis 0000-0003-3413-0666; Konstantinos Eleftherios Koumarelas 0000-0002-5614-4770; Marina Nektaria Kouliou 0000-0002-2055-2297.

S-Editor: Liu JH

L-Editor: Wang TQ

P-Editor: Yuan YY

REFERENCES

- 1 Ning XY, Ma JH, He W, Ma JT. Role of exosomes in metastasis and therapeutic resistance in esophageal cancer. *World J Gastroenterol* 2023; **29**: 5699-5715 [PMID: 38075847 DOI: 10.3748/wjg.v29.i42.5699]
- 2 Jing Z, Chen K, Gong L. The Significance of Exosomes in Pathogenesis, Diagnosis, and Treatment of Esophageal Cancer. *Int J Nanomedicine* 2021; **16**: 6115-6127 [PMID: 34511909 DOI: 10.2147/IJN.S321555]
- 3 Zhao F, Li Z, Dong Z, Wang Z, Guo P, Zhang D, Li S. Exploring the Potential of Exosome-Related lncRNA Pairs as Predictors for Immune Microenvironment, Survival Outcome, and Microbiota Landscape in Esophageal Squamous Cell Carcinoma. *Front Immunol* 2022; **13**: 918154 [PMID: 35880180 DOI: 10.3389/fimmu.2022.918154]
- 4 Zhu Q, Huang L, Yang Q, Ao Z, Yang R, Krzesniak J, Lou D, Hu L, Dai X, Guo F, Liu F. Metabolomic analysis of exosomal-markers in esophageal squamous cell carcinoma. *Nanoscale* 2021; **13**: 16457-16464 [PMID: 34648610 DOI: 10.1039/d1nr04015d]
- 5 Li Q, Li B, Li Q, Wei S, He Z, Huang X, Wang L, Xia Y, Xu Z, Li Z, Wang W, Yang L, Zhang D. Exosomal miR-21-5p derived from gastric cancer promotes peritoneal metastasis via mesothelial-to-mesenchymal transition. *Cell Death Dis* 2018; **9**: 854 [PMID: 30154401 DOI: 10.1038/s41419-018-0928-8]
- 6 Liu Z, Huang Y, Han Z, Shen Z, Yu S, Wang T, Dong Z, Kang M. Exosome-mediated miR-25/miR-203 as a potential biomarker for esophageal squamous cell carcinoma: improving early diagnosis and revealing malignancy. *Transl Cancer Res* 2021; **10**: 5174-5182 [PMID: 35116367 DOI: 10.21037/tcr-21-1123]
- 7 Chen C, Ma Z, Jiang H. EMT Participates in the Regulation of Exosomes Secretion and Function in Esophageal Cancer Cells. *Technol Cancer Res Treat* 2021; **20**: 15330338211033077 [PMID: 34278849 DOI: 10.1177/15330338211033077]
- 8 Matsumoto Y, Kano M, Murakami K, Toyozumi T, Suito H, Takahashi M, Sekino N, Shiraishi T, Kamata T, Ryuzaki T, Hirasawa S, Kinoshita K, Matsubara H. Tumor-derived exosomes influence the cell cycle and cell migration of human esophageal cancer cell lines. *Cancer*

- Sci* 2020; **111**: 4348-4358 [PMID: [32969511](#) DOI: [10.1111/cas.14660](#)]
- 9 **Warnecke-Eberz U**, Chon SH, Hölscher AH, Drebbler U, Bollschweiler E. Exosomal onco-miRs from serum of patients with adenocarcinoma of the esophagus: comparison of miRNA profiles of exosomes and matching tumor. *Tumour Biol* 2015; **36**: 4643-4653 [PMID: [25631748](#) DOI: [10.1007/s13277-015-3112-0](#)]
 - 10 **Gao J**, Zhao Z, Zhang H, Huang S, Xu M, Pan H. Transcriptomic characterization and construction of M2 macrophage-related prognostic and immunotherapeutic signature in ovarian metastasis of gastric cancer. *Cancer Immunol Immunother* 2023; **72**: 1121-1138 [PMID: [36336725](#) DOI: [10.1007/s00262-022-03316-z](#)]
 - 11 **Qiu ML**, Li X, Lin JB, Luo RG, Liu B, Feng Z. Serum exosomal miR-182 upregulation predicts unfavorable prognosis of esophageal squamous cell carcinoma. *Eur Rev Med Pharmacol Sci* 2020; **24**: 5412-5418 [PMID: [32495876](#) DOI: [10.26355/eurev.202005.21325](#)]
 - 12 **Su XY**, Si MY, Zhu ZK, Jiang Y, Liu YL. The healthy living master plan (HLMP): Singapore's experience and enlightenment. *Zhonghua Yu Fang Yi Xue Za Zhi* 2019; **53**: 1198-1202 [PMID: [31795574](#) DOI: [10.3760/cma.j.issn.0253-9624.2019.12.002](#)]
 - 13 **Xie K**, Zheng C, Gu W, Jiang Z, Luo C, Luo J, Diao Y, Wang G, Cong Z, Yao X, Hu L, Shen Y. A RASSF8-AS1 based exosomal lncRNAs panel used for diagnostic and prognostic biomarkers for esophageal squamous cell carcinoma. *Thorac Cancer* 2022; **13**: 3341-3352 [PMID: [36266257](#) DOI: [10.1111/1759-7714.14690](#)]
 - 14 **Jiao Z**, Yu A, Rong W, He X, Zen K, Shi M, Wang T. Five-lncRNA signature in plasma exosomes serves as diagnostic biomarker for esophageal squamous cell carcinoma. *Aging (Albany NY)* 2020; **12**: 15002-15010 [PMID: [32597791](#) DOI: [10.18632/aging.103559](#)]
 - 15 **Fan L**, Cao Q, Liu J, Zhang J, Li B. Circular RNA profiling and its potential for esophageal squamous cell cancer diagnosis and prognosis. *Mol Cancer* 2019; **18**: 16 [PMID: [30674324](#) DOI: [10.1186/s12943-018-0936-4](#)]
 - 16 **Rao D**, Lu H, Wang X, Lai Z, Zhang J, Tang Z. Tissue-derived exosome proteomics identifies promising diagnostic biomarkers for esophageal cancer. *Elife* 2023; **12** [PMID: [37966470](#) DOI: [10.7554/eLife.86209](#)]
 - 17 **Luo A**, Zhou X, Shi X, Zhao Y, Men Y, Chang X, Chen H, Ding F, Li Y, Su D, Xiao Z, Hui Z, Liu Z. Exosome-derived miR-339-5p mediates radiosensitivity by targeting Cdc25A in locally advanced esophageal squamous cell carcinoma. *Oncogene* 2019; **38**: 4990-5006 [PMID: [30858545](#) DOI: [10.1038/s41388-019-0771-0](#)]
 - 18 **He Z**, Li W, Zheng T, Liu D, Zhao S. Human umbilical cord mesenchymal stem cells-derived exosomes deliver microRNA-375 to downregulate ENAH and thus retard esophageal squamous cell carcinoma progression. *J Exp Clin Cancer Res* 2020; **39**: 140 [PMID: [32698859](#) DOI: [10.1186/s13046-020-01631-w](#)]
 - 19 **Zeng Q**, Zhu Z, Song L, He Z. Transferred by exosomes-derived MiR-19b-3p targets PTEN to regulate esophageal cancer cell apoptosis, migration and invasion. *Biosci Rep* 2020; **40** [PMID: [33146702](#) DOI: [10.1042/BSR20201858](#)]



Published by **Baishideng Publishing Group Inc**
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA

Telephone: +1-925-3991568

E-mail: office@baishideng.com

Help Desk: <https://www.f6publishing.com/helpdesk>

<https://www.wjgnet.com>

