



**PEER-REVIEW REPORT**

**Name of journal:** World Journal of Clinical Cases

**Manuscript NO:** 59450

**Title:** Endoscopic salvage treatment of add-on histoacryl (n-butyl-2-cyanoacrylate) sealing after fully covered metal stent application on the anastomotic leak after proximal gastrectomy

**Reviewer’s code:** 05348259

**Position:** Peer Reviewer

**Academic degree:** MD

**Professional title:** Doctor

**Reviewer’s Country/Territory:** China

**Author’s Country/Territory:** South Korea

**Manuscript submission date:** 2020-09-21

**Reviewer chosen by:** AI Technique

**Reviewer accepted review:** 2020-09-21 04:08

**Reviewer performed review:** 2020-09-21 13:44

**Review time:** 9 Hours

<b>Scientific quality</b>	<input type="checkbox"/> Grade A: Excellent <input checked="" type="checkbox"/> Grade B: Very good <input type="checkbox"/> Grade C: Good <input type="checkbox"/> Grade D: Fair <input type="checkbox"/> Grade E: Do not publish
<b>Language quality</b>	<input type="checkbox"/> Grade A: Priority publishing <input checked="" type="checkbox"/> Grade B: Minor language polishing <input type="checkbox"/> Grade C: A great deal of language polishing <input type="checkbox"/> Grade D: Rejection
<b>Conclusion</b>	<input type="checkbox"/> Accept (High priority) <input checked="" type="checkbox"/> Accept (General priority) <input type="checkbox"/> Minor revision <input type="checkbox"/> Major revision <input type="checkbox"/> Rejection
<b>Re-review</b>	<input checked="" type="checkbox"/> Yes <input type="checkbox"/> No
<b>Peer-reviewer</b>	Peer-Review: <input checked="" type="checkbox"/> Anonymous <input type="checkbox"/> Onymous



**Baishideng  
Publishing  
Group**

7041 Koll Center Parkway, Suite  
160, Pleasanton, CA 94566, USA  
**Telephone:** +1-925-399-1568  
**E-mail:** bpgoffice@wjgnet.com  
**https://**www.wjgnet.com

statements

Conflicts-of-Interest: [ ] Yes [Y] No

### **SPECIFIC COMMENTS TO AUTHORS**

Postoperative anastomotic leakage is the most serious complications after gastrectomy, most patients received conservative treatment, or with percutaneous drainage. For some limited dehiscence, endoscopic placement of metal extent could be effective. Generally, the endoscopic glue injection might have lower successful rate than the extent. This case combined two different treatment was rarely reported, and got a satisfactory result. For some relatively serious anastomotic leakage, this successful case provided a strategy and worth learning from.