

7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA **Telephone:** +1-925-399-1568 **E-mail:** bpgoffice@wjgnet.com https://www.wjgnet.com

PEER-REVIEW REPORT

Name of journal: World Journal of Clinical Cases

Manuscript NO: 69897

Title: Correlation between the Systemic Immune Inflammation Index, LMR, LMR/LDH

and prognosis of DLBCL patients

Reviewer's code: 06140404 Position: Peer Reviewer Academic degree: MD, PhD

Professional title: Associate Professor

Reviewer's Country/Territory: Switzerland

Author's Country/Territory: China

Manuscript submission date: 2021-08-18

Reviewer chosen by: AI Technique

Reviewer accepted review: 2021-08-23 14:12

Reviewer performed review: 2021-09-20 11:03

Review time: 27 Days and 20 Hours

Scientific quality	[] Grade A: Excellent [] Grade B: Very good [Y] Grade C: Good [] Grade D: Fair [] Grade E: Do not publish
Language quality	[] Grade A: Priority publishing [Y] Grade B: Minor language polishing [] Grade C: A great deal of language polishing [] Grade D: Rejection
Conclusion	[] Accept (High priority) [] Accept (General priority) [Y] Minor revision [] Major revision [] Rejection
Re-review	[Y] Yes [] No
Peer-reviewer statements	Peer-Review: [Y] Anonymous [] Onymous Conflicts-of-Interest: [] Yes [Y] No



7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA

Telephone: +1-925-399-1568 **E-mail:** bpgoffice@wjgnet.com

https://www.wjgnet.com

SPECIFIC COMMENTS TO AUTHORS

Most diffuse large B-cell lymphoma patients can achieve a complete response after first-line treatment. And finding effective biomarkers to predict the prognosis and recurrence of diffuse large B-cell lymphoma among patients is still needed. Studies show that the ratio of lymphocytes to monocytes can best reflect the host immune state, thus can be an effective prognostic marker in diffuse large B-cell lymphoma patients. The systemic immune inflammation index is a comprehensive inflammatory marker based on peripheral blood neutrophils, platelets and lymphocytes count, which can be used to predict the clinical prognosis of non-small cell lung cancer, gastric cancer, pancreatic cancer and many other solid tumors. In this study, the authors explored and discussed the association between SII, LMR, LMR/LDH, and prognosis of diffuse large B-cell lymphoma patients. The study is over all very interesting. The manuscript is very well written. Some minor comments: 1. A minor language polishing should be revised. 2. The images are too small, please check and update. 3. The references should edited and updated.



7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA **Telephone:** +1-925-399-1568 **E-mail:** bpgoffice@wjgnet.com https://www.wjgnet.com

PEER-REVIEW REPORT

Name of journal: World Journal of Clinical Cases

Manuscript NO: 69897

Title: Correlation between the Systemic Immune Inflammation Index, LMR, LMR/LDH

and prognosis of DLBCL patients

Reviewer's code: 03360012 Position: Peer Reviewer Academic degree: MD

Professional title: Professor

Reviewer's Country/Territory: Netherlands

Author's Country/Territory: China

Manuscript submission date: 2021-08-18

Reviewer chosen by: Jin-Lei Wang

Reviewer accepted review: 2021-09-07 12:20

Reviewer performed review: 2021-09-20 11:19

Review time: 12 Days and 22 Hours

Scientific quality	[] Grade A: Excellent [] Grade B: Very good [Y] Grade C: Good [] Grade D: Fair [] Grade E: Do not publish
Language quality	[] Grade A: Priority publishing [Y] Grade B: Minor language polishing [] Grade C: A great deal of language polishing [] Grade D: Rejection
Conclusion	[] Accept (High priority) [] Accept (General priority) [Y] Minor revision [] Major revision [] Rejection
Re-review	[]Yes [Y]No
Peer-reviewer	Peer-Review: [Y] Anonymous [] Onymous
statements	Conflicts-of-Interest: [] Yes [Y] No



7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA **Telephone:** +1-925-399-1568 **E-mail:** bpgoffice@wjgnet.com

https://www.wjgnet.com

SPECIFIC COMMENTS TO AUTHORS

This is an interesting study about the association between systemic immune inflammation index, ratio of lymphocytes to monocytes, ratio of LMR to LDH and prognosis of patients with diffuse large B-cell lymphoma. The study is well designed. The results are discussed with updated references. Before publication, a minor editing is required.