



Baishideng Publishing Group Co., Limited

Flat C, 23/F., Lucky Plaza,
315-321 Lockhart Road,
Wan Chai, Hong Kong, China

ESPS Peer-review Report

Name of Journal: World Journal of Gastroenterology

ESPS Manuscript NO: 7595

Title: Two rare gastric hamartomatous inverted polyp cases suggest the pathogenesis of their growth

Reviewer code: 02504712

Science editor: Wen, Ling-Ling

Date sent for review: 2013-11-26 21:43

Date reviewed: 2013-12-03 23:24

CLASSIFICATION	LANGUAGE EVALUATION	RECOMMENDATION	CONCLUSION
<input type="checkbox"/> Grade A (Excellent)	<input type="checkbox"/> Grade A: Priority Publishing	Google Search:	<input type="checkbox"/> Accept
<input type="checkbox"/> Grade B (Very good)	<input checked="" type="checkbox"/> Grade B: minor language polishing	<input type="checkbox"/> Existed	<input type="checkbox"/> High priority for publication
<input checked="" type="checkbox"/> Grade C (Good)	<input type="checkbox"/> Grade C: a great deal of language polishing	<input type="checkbox"/> No records	<input type="checkbox"/> Rejection
<input type="checkbox"/> Grade D (Fair)	<input type="checkbox"/> Grade D: rejected	<input type="checkbox"/> Existed	<input checked="" type="checkbox"/> Minor revision
<input type="checkbox"/> Grade E (Poor)		<input type="checkbox"/> No records	<input type="checkbox"/> Major revision

COMMENTS TO AUTHORS

Manuscript title: Two rare gastric hamartomatous inverted polyp cases suggest the pathogenesis of their growth. Case report Authors: Mori H et al. The manuscript describes two case reports of rare gastric hamartomatous inverted polyps (GHIPs). In addition, the authors hypothesise that after a hamartomatous change occurs in the submucosal layer, some of these components are exposed to the gastric mucosa and, consequently, form a hypertrophic lesion which inverts. The authors also advance the ultrasonographic appearance of these hamartomatous lesions. In general, These cases are rare and a detailed pathological and ultrasonographic characterization is relevant. Although, this is not original, as these lesions have been described previously, the combination of the pathological and ultrasonographic appearance is novel and useful. Bearing in mind the rarity of these lesions, their significance is important when they are diagnosed. The manuscript is generally readable with a few editorial corrections necessary. There are no ethical issues to be addressed in this article. Specifically, the two case reports are satisfactory. The authors hypothesise the formation of these hamartomatous lesions and they just need to stress that these are hypothesis. I am happy with the suggestions made but they remain unproven scientifically.



Baishideng Publishing Group Co., Limited

Flat C, 23/F., Lucky Plaza,
315-321 Lockhart Road,
Wan Chai, Hong Kong, China

ESPS Peer-review Report

Name of Journal: World Journal of Gastroenterology

ESPS Manuscript NO: 7595

Title: Two rare gastric hamartomatous inverted polyp cases suggest the pathogenesis of their growth

Reviewer code: 02563195

Science editor: Wen, Ling-Ling

Date sent for review: 2013-11-26 21:43

Date reviewed: 2013-12-15 20:29

CLASSIFICATION	LANGUAGE EVALUATION	RECOMMENDATION	CONCLUSION
<input checked="" type="checkbox"/> Grade A (Excellent)	<input checked="" type="checkbox"/> Grade A: Priority Publishing	Google Search:	<input type="checkbox"/> Accept
<input type="checkbox"/> Grade B (Very good)	<input type="checkbox"/> Grade B: minor language polishing	<input type="checkbox"/> Existed	<input type="checkbox"/> High priority for publication
<input type="checkbox"/> Grade C (Good)	<input type="checkbox"/> Grade C: a great deal of	<input type="checkbox"/> No records	<input type="checkbox"/> Rejection
<input type="checkbox"/> Grade D (Fair)	language polishing	BPG Search:	<input type="checkbox"/> Rejection
<input type="checkbox"/> Grade E (Poor)	<input type="checkbox"/> Grade D: rejected	<input type="checkbox"/> Existed	<input checked="" type="checkbox"/> Minor revision
		<input type="checkbox"/> No records	<input type="checkbox"/> Major revision

COMMENTS TO AUTHORS

They are interesting cases described in manuscript



Baishideng Publishing Group Co., Limited

Flat C, 23/F., Lucky Plaza,
315-321 Lockhart Road,
Wan Chai, Hong Kong, China

ESPS Peer-review Report

Name of Journal: World Journal of Gastroenterology

ESPS Manuscript NO: 7595

Title: Two rare gastric hamartomatous inverted polyp cases suggest the pathogenesis of their growth

Reviewer code: 02537189

Science editor: Wen, Ling-Ling

Date sent for review: 2013-11-26 21:43

Date reviewed: 2013-12-16 12:31

CLASSIFICATION	LANGUAGE EVALUATION	RECOMMENDATION	CONCLUSION
<input type="checkbox"/> Grade A (Excellent)	<input type="checkbox"/> Grade A: Priority Publishing	Google Search:	<input type="checkbox"/> Accept
<input type="checkbox"/> Grade B (Very good)	<input checked="" type="checkbox"/> Grade B: minor language polishing	<input type="checkbox"/> Existed	<input checked="" type="checkbox"/> High priority for publication
<input checked="" type="checkbox"/> Grade C (Good)	<input type="checkbox"/> Grade C: a great deal of language polishing	<input type="checkbox"/> No records	<input type="checkbox"/> Rejection
<input type="checkbox"/> Grade D (Fair)		BPG Search:	<input type="checkbox"/> Minor revision
<input type="checkbox"/> Grade E (Poor)	<input type="checkbox"/> Grade D: rejected	<input type="checkbox"/> Existed	<input type="checkbox"/> Major revision
		<input type="checkbox"/> No records	

COMMENTS TO AUTHORS

Gastric hamartomatous inverted polyps (GHIP) are difficult to accurately diagnose because of their inversion into the submucosal layer. An en bloc resection by endoscopic submucosal dissection (ESD) is recommended for treatment. This case report will be helpful the treatment of GHIP.