



Baishideng Publishing Group Co., Limited

Flat C, 23/F., Lucky Plaza,
315-321 Lockhart Road, Wan Chai, Hong Kong, China

ESPS Peer-review Report

Name of Journal: World Journal of Gastroenterology

ESPS Manuscript NO: 8933

Title: One fifth of hospitalizations for peptic ulcer-related bleeding are potentially preventable

Reviewer code: 00051367

Science editor: Gou, Su-Xin

Date sent for review: 2014-01-23 10:17

Date reviewed: 2014-01-28 16:00

CLASSIFICATION	LANGUAGE EVALUATION	RECOMMENDATION	CONCLUSION
<input type="checkbox"/> Grade A (Excellent)	<input checked="" type="checkbox"/> Grade A: Priority Publishing	Google Search:	<input checked="" type="checkbox"/> Accept
<input type="checkbox"/> Grade B (Very good)	<input type="checkbox"/> Grade B: minor language polishing	<input type="checkbox"/> Existed	<input type="checkbox"/> High priority for publication
<input checked="" type="checkbox"/> Grade C (Good)	<input type="checkbox"/> Grade C: a great deal of language polishing	<input type="checkbox"/> No records	<input type="checkbox"/> Rejection
<input type="checkbox"/> Grade D (Fair)		BPG Search:	<input type="checkbox"/> Minor revision
<input type="checkbox"/> Grade E (Poor)	<input type="checkbox"/> Grade D: rejected	<input type="checkbox"/> Existed	<input type="checkbox"/> Major revision
		<input type="checkbox"/> No records	

COMMENTS TO AUTHORS

The paper reports a retrospective analysis. By placing data on low-risk patients the authors could, for comparison, highlight the use of gastroprotection in patients at low risk, that constitute the majority of bleeding (2/3) and discuss the use of PPI on the basis of the prevalence estimates obtained. A discussion should be provided about the discrepancy between expert consensus and real world, since only a fourth of bleeding patients could be framed in an assessment based on current guidelines