



## Baishideng Publishing Group Co., Limited

Flat C, 23/F., Lucky Plaza,  
315-321 Lockhart Road,  
Wan Chai, Hong Kong, China

---

### ESPS Peer-review Report

**Name of Journal:** World Journal of Gastrointestinal Oncology

**Ms:** 2995

**Title:** A GIST OF GASTROINTESTINAL STROMAL TUMOURS: A REVIEW

**Reviewer code:** 00039581

**Science editor:** l.l.wen@wjgnet.com

**Date sent for review:** 2013-04-01 10:54

**Date reviewed:** 2013-04-05 14:25

CLASSIFICATION	LANGUAGE EVALUATION	RECOMMENDATION	CONCLUSION
<input type="checkbox"/> Grade A (Excellent)	<input checked="" type="checkbox"/> Grade A: Priority Publishing	Google Search:	<input checked="" type="checkbox"/> Accept
<input checked="" type="checkbox"/> Grade B (Very good)	<input type="checkbox"/> Grade B: minor language polishing	<input type="checkbox"/> Existed	<input type="checkbox"/> High priority for publication
<input type="checkbox"/> Grade C (Good)	<input type="checkbox"/> Grade C: a great deal of language polishing	<input type="checkbox"/> No records	<input type="checkbox"/> Rejection
<input type="checkbox"/> Grade D (Fair)		BPG Search:	<input type="checkbox"/> Minor revision
<input type="checkbox"/> Grade E (Poor)	<input type="checkbox"/> Grade D: rejected	<input type="checkbox"/> Existed	<input type="checkbox"/> Major revision
		<input type="checkbox"/> No records	

### COMMENTS

#### COMMENTS TO AUTHORS:

This article is interesting review regarding GIST. It summarized well about the gist of history about GIST. I read this with great interest and thank you for author' s contribution. Comments 1. Small GIST of stomach could be managed conservatively with regular surveillance; however, endoscopic resection would be another option. There are several reports regarding these results. Please add this statement to the 'Management' section with some references. 2. Also laparoscopic or open resection could be TOC in larger (>2cm) GIST with malignant potential. Recent studies also show that combined modality (laparoscopy + endoscopic resection) would be good treatment modality in some patients, who had tumor at cardia or high gastric body.