

## **Title Page**

### **Combined liver and kidney transplantation in children and long-term outcome**

**Name of Journal:** *World Journal of Transplantation*

**Manuscript No:** 58104

**Manuscript Type:** REVIEW

## **Authors**

**Randula Ranawaka, Kavinda Dayasiri, Manoji Gamage**

Randula Ranawaka, Consultant Paediatric Nephrologist and Senior Lecturer, Department of Paediatrics, Faculty of Medicine, University of Colombo, Sri Lanka

Kavinda Dayasiri, Consultant Paediatrician, Base Hospital Mahaoya, Mahaoya, Sri Lanka

Manoji Gamage, Clinical Nutritionist, Lady Ridgeway Hospital for Children, Colombo, Sri Lanka

Author contributions: RR, KD, MG performed literature survey, RR, KD, MG wrote manuscript. RR edited the final version of the manuscript. All authors read and approved the final version of the manuscript.

## **Acknowledgement**

**All authors sincerely thank Dr. Dominic Kelly of University of Oxford for conducting English language editing of this manuscript.**

Corresponding Author

Dr. Randula Ranawaka

Department of Paediatrics, Faculty of Medicine

University of Colombo

Kynsey Road, Colombo 00800

Sri Lanka

Email: rrandula@yahoo.com

## Introduction

The first successful combined liver-kidney transplantation (CLKT) in an adult was reported in 1984. After that it has become more common in adults but remains a relatively infrequent procedure in children (1). Between 1988 and 2007, there were 2829 CLKT procedures performed in the USA of which only 166 were carried out in children (2). Combined liver-kidney transplantation is a challenging form of surgery in the paediatric age group (1) with only about thirty surgical procedures being performed worldwide over a given year (3). CLKT is a procedure in which a liver (whole or in part) and kidney allografts, from the same deceased **or living donor**, are transplanted during a single surgical procedure. Sequential liver-kidney transplantation (SLKT) is isolated transplantation of the liver (or kidney) followed by the transplantation of the kidney (or liver) after a certain time interval.

Limited numbers of this surgical procedure are due to rarity of clear indications for surgery, long waiting times for the surgery and shortage of suitable donors (4). With advancement of medical practices and innovative surgery, it is crucial that clinicians are insightful about when would CLKT be the right choice for a child, as compared to alternative therapeutic and surgical interventions. In the paediatric age group, CLKT can be considered for: (1) children with profound and irreversible liver and kidney disease eg. autosomal recessive polycystic kidney disease; (2) children with metabolic liver diseases causing end-stage renal disease eg. primary hyperoxaluria and atypical haemolytic uremic syndrome. A minority of children undergo CLKT following combined liver and renal failure. CLKT has been shown to be associated with immunoprotection of the kidney allograft and reduced rejection rates as compared with sequential kidney and liver transplantation (5) or kidney transplantation alone (5,6,7). However, experience of this observation is limited in paediatric studies due to lack of comparative paediatric studies (8).

This review focuses on clinical indications, procedures, and acute, short and long term outcomes of combined liver-kidney transplantation in the paediatric age group. The review also discusses evidence that support the immunoprotective effects of CLKT on renal graft survival.

## Indications

### 1. Autosomal recessive polycystic kidney disease (ARPKD)

ARPKD is a rare renal cystic disease in children with an incidence of 1:20,000 (9). It is also known as fibro-polycystic liver and kidney disease and the most common renal cystic/ciliopathy disease in children. In ARPKD there is a mutation in PKDHD1, a gene located on chromosome 6p12 (10). The defective gene encodes for fibrocystin and leads to defective tubular formation in renal tubules and hepatic bile ducts, leading to the early development of cysts (10). Most severe

variants are associated with pulmonary hypoplasia and high neonatal mortality. Approximately 50% of children develop end stage renal disease (ESRD) during the first decade of life (11). Hepatic fibrosis is also seen early in life leading to recurrent cholangitis and manifestations of chronic liver disease which includes portal hypertension, splenomegaly and variceal haemorrhage (8). However, chronic liver failure has a variable age of onset as opposed to early onset End Stage Renal Disease (ESRD), making management of these children quite a challenge. These children undergo early nephrectomy and renal transplantation followed by sequential liver transplantation (12) or combined liver-kidney transplantation.

## 2. Primary hyperoxaluria type 1 (PH-1)

PH-1 is a rare disorder of oxalate metabolism which has an incidence of 1:120,000 (13). It is an autosomal recessive disease and the most common indication for CLKT in children. Primary hyperoxaluria is characterized by elevated plasma and urinary oxalate levels due to the defective liver specific peroxisomal enzyme alanine/glyoxylate aminotransferase (14). Increased accumulation of glyoxalate leads to increased oxalate and formation and deposition of insoluble calcium salts in kidney (15). Children with PH-1 develop early onset nephrocalcinosis, nephrolithiasis and ESRD (16). With development of ESRD and impaired excretion, oxalate is deposited in other tissues such as retina, blood vessels, nerves and heart (17). PH-1 often has a variable age of onset for ESRD and oxaluria is responsive to pyridoxine in some children. Therefore, it is important that pyridoxine responsiveness is evaluated and DNA analysis is performed to confirm the diagnosis in all children before planning CLKT (14).

The definitive treatment is CKLT to prevent early recurrence of nephrocalcinosis in the transplanted kidney (18). However, in developing countries, isolated kidney transplantation is considered initially unless a child is referred to a centre performing CLKT (19). Hyperhydration is recommended in immediate post-operative period to prevent the surge of plasma oxalate due to mobilization of oxalate from other tissues and subsequent damage to the transplanted kidney (14). Hyperhydration should be supplemented by post-operative haemodialysis and use of crystallization inhibitors (eg. citrate) for the same reason (20).

## 3. Atypical hemolytic uremic syndrome (aHUS)

Atypical HUS is a rare disorder of the alternative complement pathway involving impaired synthesis or function of factor H, a complement control protein. This leads to a triad of microangiopathic haemolytic anaemia, thrombocytopenia and renal dysfunction (21). For aHUS CLKT was previously the treatment of choice as it corrected both the renal failure and the underlying problem with factor H which is produced by the liver. However, this procedure is no longer recommended as the first choice of definitive treatment due to promising effects of Eculizumab (anti-C5 monoclonal antibody) as a medical treatment and significant incidence of

long term post-operative complications of CLKT. Eculizumab has been shown to inhibit complement activation in alternative pathway that leads to microangiopathy (22), thereby avoiding the need for liver transplantation. However, there are certain genetic variants such as **DGKE (diacylglycerol kinase-epsilon)** mutations for which Eculizumab is not effective (23). AN international consensus statement recommends either liver transplant or CLKT as the only treatment of cure for severe aHUS or defective complement factors synthesized in the liver (**CFH – Complement factor H, CFB – Complement factor B, C3 – Complement 3**) although Eculizumab is also given to reduce post-transplant recurrence in those who only had renal transplantation (24). Use of Eculizumab is limited by lack of availability and very high cost.

#### 4. Methylmalonic acidemia

Methylmalonic acidemia is a rare inherited autosomal recessive metabolic disorder mainly due to defective vitamin-B12-dependent enzyme, methylmalonyl-CoA mutase leading to increased formation of methylmalonic acid. Children with methylmalonic acidemia are at high risk of multi-organ complications including heart, kidney, eyes and nervous system (25). These children can also present with acute metabolic crises with profound metabolic acidosis and seizures (26). Methylmalonic acidemia leads to renal tubular interstitial injury and up to 60% of children develop ESRD during adolescence (27). Since methylmalonyl-CoA mutase activity is present in both liver and kidney, transplantation of one organ will lead to only partial recovery with a risk of recurrence (28). CLKT has been shown to improve methylmalonic acidemia, renal dysfunction<sup>2</sup> and overall quality of life (29).

However, even after the CLKT, some systemic disease manifestations (such as neurologic or muscle impairment) may persist despite normal liver and kidney graft function due to abnormal methylmalonic acid metabolism in other tissues, including the muscles and skin (28). Therefore, it is crucial to provide lifelong specific high-calorie diet low in propiogenic amino acid precursors despite organ transplantation, due to on-going production of methylmalonic acid from skeletal muscles (30).

#### 5. Combined liver and kidney failure

Combined liver and kidney failure occurs in certain metabolic diseases such as alpha-1 antitrypsin deficiency, glycogen storage disease type 1A, Boichis syndrome (nephronophthisis with congenital hepatic fibrosis), and medical conditions such as hepatorenal syndrome and liver tumor with nephrotoxicity. CLKT had been variably successful in children with these conditions (12).

Hepatorenal Syndrome (HRS) is one of the main complications of end stage liver disease with high morbidity and mortality. It results from hypoperfusion of kidneys due to combined effects

of intrarenal arteriolar vasoconstriction and peripheral vasodilatation, mainly in the splanchnic circulation. There are two types of HRS, Type 1 (with the worst prognosis) is rapidly progressive with renal failure while type 2 has slowly developing renal dysfunction in patients with liver cirrhosis.

In most instances, HRS resolves with liver transplantation alone thus HRS is not being considered routinely for CLKT (31). However as prolonged HRS can progress to irreversible renal damage some patients with both end-stage liver and kidney failure may be candidates for CLKT (3, 32).

## **Procedure**

In CLKT, both kidney and either whole or part of liver from a donor (usually cadaveric) is transplanted to the same recipient during a single surgical procedure. Sequential liver-kidney transplantation is performed in two stages where the recipient initially undergoes isolated organ transplantation (either kidney or liver) followed by the other organ (liver or kidney) from two different cadaveric donors or a single living donor. It is of the paramount importance to keep the cold ischemic time shorter to avoid delayed graft function.

Whole cadaveric liver transplantation would help to reduce the post-transplant complications (e.g. bleeding, bile leak) by avoiding prolonged cold ischemic time for both liver and kidney, compared to partial liver graft transplantation. The liver graft is transplanted first to reduce the risk of cold ischaemia to liver and cold ischaemic time is usually kept to less than 8 - 10 hours for the liver and 10 – 12 hours for the kidney. After hepatic vascular anastomoses are performed, renal vessels are anastomosed to the common iliac vessels to achieve early re-perfusion. **Uretero-vesical** and biliary anastomoses are performed only after hepatic and renal vascular re-perfusion is achieved.

## **Acute complications and short term outcomes**

CLKT is a complex major surgical procedure and immediate post-operative complications are frequently reported. Analysis of the Scientific Registry of Transplant Recipients (<https://www.srtr.org/>) of 152 primary paediatric CLKTs performed from October 1987 to February 2011, revealed a total of 32 deaths (21.1%) during the first 30 month postoperative period (33). The main causes of death were: infectious (18.7%), cardiovascular (18.7%), respiratory (6.2%), gastro-intestinal and haemorrhage (6.2%). However with the advancement of surgical and medical interventions, the rate of complications has been progressively reducing over the last decade.

Post-operative hyperoxaluria and graft dysfunction is a major problem following **CLKT** for primary hyperoxaluria and ESRD and these patients should be managed with post-operative

renal replacement therapy. Many studies reported that the acute post CLKT complications were higher in primary hyperoxaluria compared to other causes (34).

Further complications including, post-operative bleeding, bile leaks, hepatic artery thrombosis, and acute liver failure can lead to graft loss and even mortality. Due to bleeding and multiple vascular anastomoses, patients are at risk of hypovolaemia and shock or fluid overload due to multiple transfusions. Therefore, fluids and diuretics should be used carefully. Liver graft can suffer cold ischaemia **with longer anastomosis times** and the risk is higher with partial liver transplantations compared to whole liver transplantations (35).

Mortality during initial six months following CLKT remains high. Septicaemia and multi-organ failure is a common cause of mortality while on high doses of immunosuppressive medications (36). Acute organ rejection was reported in 14% of CLKT patients during first post-operative year in one study (37). The combination of basiliximab and daclizumab reduces acute rejection in renal transplantation when used during induction phase of immunosuppression (38). Calinescu et al. examined 152 patients who had CLKT for short and long term outcomes. Overall, the one year patient survival was 86.8% and the survival of the kidney and liver grafts was 83.4% and 81.9% respectively.

### **Immunoprotective effects of CLKT on renal allograft**

In CLKT patients, the liver allograft confers immunoprotection of the kidney allograft. This was first demonstrated in animal model (39). Later clinical studies of CLKT have shown reductions in both acute cellular rejection and chronic rejection of kidney allograft in those who undergo CLKT when compared to cadaveric renal transplantation (37). The molecular basis of immunoprotection is yet to be precisely defined but it is thought that the transplanted liver can modify the immune system of the recipient by neutralizing circulating autoantibodies (40). Further, a liver graft provides HLA-G antigens that inhibit natural killer cells thought to be involved in acute rejection (41). Improved renal outcomes from such immunoprotection in part explain the fact that CLKT has better outcomes as compared with sequential organ transplantation in patients with combined liver and renal impairment. Liver rejection rates were not different in those with CLKT when compared to isolated liver transplantations (42). Rapamycin, a calcineurin inhibitor, is the mainstay of immunosuppression following CKLT. Use of Rapamycin has been associated with reduced rates of hepatic rejection in adult studies (43).

### **Long term outcome**

The result of CLKT is in part dependent on the aetiology of hepatic and kidney dysfunction. The outcome is also dependent on the child's pre-transplant clinical condition. Diagnosis of a metabolic disease and an early transplantation scenario are good prognostic indicators while multi-organ failure leading to CLKT is a poor prognostic indicator. The long term outcome of the transplanted liver is usually good without evidence of chronic rejection. However, a chronic allograft nephropathy of the kidney is apparent in the long term.

Data regarding long term outcomes of CLKT are limited. Bernabeu et al. reported long term outcomes of 14 children who had CLKT (42). The majority of patients in that cohort had either ARPKD or PH-1. One child had renal re-transplantation following branch renal artery thrombosis of the kidney allograft after it was complicated by severe hypertension and renal dysfunction. One patient had chronic rejection 10 years after the transplant. Two children had BK viral infections which were treated with cidofovir whilst one child died following adenoviral infection 8 months after CLKT. Several children developed progressive renal dysfunction and severe tubulopathy indicative of allograft nephropathy and needed renal re-transplantation.

Long term outcome was dependent on the primary disorder and patients with PH-1 have slower improvement of renal functions as compared with ARPKD. Five-year renal graft survival and overall survival were 85.7% and 92.9% respectively. More recently, Ranawaka et al. (6) reported long term renal outcomes in a cohort of 40 children who had CLKT with the majority being diagnosed with ARPKD. The investigators observed a statistically significant greater decline of estimated glomerular filtration rate (e-GFR) in isolated kidney transplant patients compared to those who had CLKT whilst acute rejection was less in those with CLKT. The investigators observed better outcomes in those with ARPKD compared to PH-1 and comparative to observations made by Bernabeu et al. Although there are several studies which reported long term renal outcomes, only a few studies reported long term liver outcomes of which 5-year graft survival varied from 76.5% (33)- 80% (44).

In one large series, Calinescu et al analyzed data using the Scientific Registry of Transplant Recipients to determine long term outcomes of 152 pediatric CLKT. The liver graft survival at five and ten years, were 76.5%, and 72.6 % respectively, and kidney graft survival was 76.5% and 66.8 %, respectively. Patient survival was 82.1% at five years, and 78.9% at ten years, which was much similar to the isolated liver transplant at five and ten years (81.2% and 77.4%). But isolated kidney transplant showed much better results at those points (95.4% and 90%) (33).

The children with liver and kidney failure are shorter than their peers due to underlying chronic conditions but show improved growth after CLKT. Growth is an important determinant of better functional outcomes in employment, education and marital life (45, 46). North American Paediatric Renal Trials and Collaborative Studies data have shown that better catch-up growth is achieved when CLKT is performed at a younger age, especially below six years (47). Human growth hormone has a place in this group of children and a recent Cochrane update demonstrated that children who were treated with growth hormone showed an increased height velocity of 3.88 cm/year (48).

Although life-saving, CLKT may not be curative in all children, and children who had CLKT may suffer from chronic health problems throughout their life. Therefore, the quality of life is an important aspect to assess in long term outcome of transplant recipients. The World Health Organization defines health as a state of complete physical, mental and social well-being and not merely the absence of disease or infirmity. Thus, it is of the paramount importance to have a

holistic approach from the multidisciplinary transplantation team with participation of primary care physicians, community nurse, psychologists, social workers and the school. Such an approach should extend to address the common attentional, behavioural and peer relationship problems of these children to improve their overall school performance. Furthermore, they should be guided through a proper adolescent transition programme to achieve their ultimate goals in life as young adults.

## Conclusion

CLKT is a complex surgical procedure which is increasingly performed for a number of indications, especially PH1 and ARPKD. Given the risks of the procedure, selection of patients for this surgery needs to be done carefully taking into consideration, the clinical indication, the functional status of liver and kidney, the patient's general health, and the skills and expertise of the transplant centre. In CLKT, the early mortality is mostly related to infections and surgical complications of the liver graft. On the other hand, chronic complications with liver graft are fairly rare, and the liver protects the kidney allograft from rejection, which results in stable function and long term survival of the renal allograft.

However, liver survival rates are not different in CLKT compared to sequential liver and kidney transplantation or isolated liver transplantation. Acute complications are not uncommon given the complexity of the surgical procedure. However, these can be reduced by optimal precautions and early detection and management of problems. Long-term outcomes are promising in children who have had CLKT with good overall long-term survival rates when performed in experienced centres with expertise.

## References

1. Margreiter R, Kramar R, Huber C, Steiner E, Niederwieser D, Judmaier G, Vogel W (1984) Combined liver and kidney transplantation. *Lancet* 1: 1077–1078 PMID: 6144007 DOI: 10.1016/s0140-6736(84)91486-7
2. Sutherland SM, Alexander SR, Sarwal MM, Berquist WE, Concepcion W (2008) Combined liver-kidney transplantation in children: Indications and outcome. *Pediatr Transplant* 12: 835–846 PMID: 19000066 DOI: 10.1111/j.1399-3046.2008.01041.x
3. Jalanko H, Pakarinen M (2014) Combined liver and kidney transplantation in children. *Pediatr Nephrol* 29(5):805–814 PMID: 23644898 DOI: 10.1007/s00467-013-2487-7
4. Otte JB (2002) History of pediatric liver transplantation. Where are we coming from? Where do we stand? *Pediatr Transplant* 6(5):378–387 PMID: 12390423 DOI: 10.1034/j.1399-3046.2002.01082.x
5. Opelz G, Margreiter R, Dohler B (2002) Prolongation of long term kidney graft survival by a simultaneous liver transplant: The liver does it, and the heart does it too. *Transplantation*: 74: 1390–1394 PMID: 12451237 DOI: 10.1097/00007890-200211270-00008
6. Ranawaka R, Lloyd C, McKiernan PJ, Hulton SA, Sharif K, Milford DV. Combined liver and kidney transplantation in children: analysis of renal graft outcome. *Pediatr Nephrol*.

2016;31(9):1539-1543. doi:10.1007/s00467-016-3396-3 PMID: 27105881 DOI: 10.1007/s00467-016-3396-3

7. Creput C, Durrbach A, Samuel D, Eschwege P, Amor M, Kriaa F, Kreis H, Benoit G, Bismuth H, Charpentier B (2003) Incidence of renal and liver rejection and patient survival rate following combined liver and kidney transplantation. *Am J Transplant* 3:348–356 PMID: 12614293 DOI: 10.1034/j.1600-6143.2003.00050.x
8. Millan MT, Berquist WE, So SK, Sarwal MM, Wayman KI, Cox KL, Filler G, Salvatierra O Jr, Esquivel CO. (2003) One hundred percent patient and kidney allograft survival with simultaneous liver and kidney transplantation in infants with primary hyperoxaluria: A single-centre experience. *Transplantation* 76:1458–1463 PMID: 14657686 DOI: 10.1097/01.TP.0000084203.76110.AC
9. Zerres K, Mucher G, Becker J, Steinkamm C, Rudnik-Schoneborn S, Heikkila P, Rapola J, Salonen R, Germino G G, Onuchic L, Somlo S, Avner E D, Harman L A, Stockwin J M, Guay-Woodford L M. Prenatal diagnosis of autosomal recessive polycystic kidney disease (ARPKD): molecular genetics, clinical experience, and fetal morphology. *Am J Med Genet.* 1998; 76: 137-144. PMID: 9511976
10. Ward CJ, Yuan D, Masyuk TV, Wang X, Punyashthiti R, Whelan S, Bacallao R, Torra R, LaRusso NF, Torres VE, Harris PC. Cellular and subcellular localization of the ARPKD protein; fibrocystin is expressed on primary cilia. *Hum Mol Genet.* 2003; 12: 2703-2710. PMID: 12925574 DOI: 10.1093/hmg/ddg274
11. Brinkert F, Lehnhardt A, Montoya C, Helmke K, Schaefer H, Fischer L, Nashan B, Bergmann C, Ganschow R, Kemper MJ. Combined liver-kidney transplantation for children with autosomal recessive polycystic kidney disease (ARPKD): indication and outcome. *Transpl Int.* 2013; 26: 640-650. PMID: 23582048 DOI: 10.1111/tri.12098
12. Cha D, Concepcion K, Gallo A, Concepcion W. Combined Liver Kidney Transplantation in Pediatrics: Indications, Special Considerations, and Outcomes. *Clin Surg.* 2017; 2: 1352.
13. van Woerden CS, Groothoff JW, Wanders RJ, Davin JC, Wijburg FA. Primary hyperoxaluria type 1 in The Netherlands: prevalence and outcome. *Nephrol Dial Transplant.* 2003; 18: 273-279. PMID: 12543880 DOI: 10.1093/ndt/18.2.273
14. Ganschow R, Hoppe B. Review of combined liver and kidney transplantation in children. *Pediatr Transplant.* 2015; 19: 820-826. PMID: 26354144 DOI: 10.1111/petr.12593
15. Milliner DS, Harris PC, Cogal AG, Lieske JC. Primary Hyperoxaluria Type 1. 2002 Jun 19 [Updated 2017 Nov 30]. In: Adam MP, Ardinger HH, Pagon RA, et al., editors. *GeneReviews®* [Internet]. Seattle (WA): University of Washington, Seattle; 1993-2020. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK1283/>
16. Khan K, Schwarzenberg SJ, Sharp HL, Matas AJ, Chavers BM. Morbidity from congenital hepatic fibrosis after renal transplantation for autosomal recessive polycystic kidney disease. *Am J Transplant.* 2002; 2: 360-365. PMID: 12118859 DOI: 10.1034/j.1600-6143.2002.20412.x
17. Bacchetta J, Mekahli D, Rivet C, Demède D, Leclerc AL. Pediatric combined liver-kidney transplantation: a 2015 update. *Curr Opin Organ Transplant.* 2015; 20: 543-549. PMID: 26270957 DOI: 10.1097/MOT.0000000000000225
18. Bergstralh EJ, Monico CG, Lieske JC, Herges RM, Langman CB, Hoppe B, Milliner DS, IPHR Investigators. Transplantation outcomes in primary hyperoxaluria. *Am J*

- Transplant. 2010; 10: 2493-2501. PMID: 20849551 DOI: 10.1111/j.1600-6143.2010.03271.x
19. Harambat J, van Stralen KJ, Espinosa L, Groothoff JW, Hulton SA, Cerkauskiene R, Schaefer F, Verrina E, Jager KJ, Cochat P, European Society for Pediatric Nephrology/European Renal Association-European Dialysis and Transplant Association (ESPN/ERA-EDTA) Registry. Characteristics and outcomes of children with primary oxalosis requiring renal replacement therapy. *Clin J Am Soc Nephrol*. 2012; 7: 458-465. PMID: 22223608 DOI: 10.2215/CJN.07430711
  20. Ganschow R, Hoppe B. Review of combined liver and kidney transplantation in children. *Pediatr Transplant*. 2015; 19: 820-826. PMID: 26354144 DOI: 10.1111/petr.12593
  21. Noris M, Caprioli J, Bresin E, Mossali C, Pianetti G, Gamba S, Daina E, Fenili C, Castelletti F, Sorosina A, Piras R, Donadelli R, Maranta R, van der Meer I, Conway EM, Zipfel PF, Goodship TH, Remuzzi G. Relative role of genetic complement abnormalities in sporadic and familial aHUS and their impact on clinical phenotype. *Clin J Am Soc Nephrol*. 2010; 5:1844-1859. PMID: 20595690 DOI: 10.2215/CJN.02210310
  22. Licht C, Greenbaum LA, Muus P, Babu S, Bedrosian CL. Efficacy and safety of eculizumab in atypical hemolytic uremic syndrome from 2-year extensions of phase 2 studies. *Kidney Int*. 2015; 87: 1061-1073. PMID: 25651368 DOI: 10.1038/ki.2014.423
  23. Lemaire M, Frémeaux-Bacchi V, Schaefer F, Choi M, Tang WH. Recessive mutations in DGKE cause atypical hemolytic-uremic syndrome. *Nat Genet*. 2013; 45: 531-536. PMID: 23542698 DOI: 10.1038/ng.2590
  24. Lieske JC, Monico CG, Holmes WS, et al. International registry for primary hyperoxaluria. *Am J Nephrol*. 2005;25(3):290-296. PMID: 15961949 DOI: 10.1159/000086360
  25. Fraser JL, Venditti CP. Methylmalonic and propionic acidemias: clinical management update. *Curr Opin Pediatr*. 2016; 28: 682-693 PMID: 27653704 DOI: 10.1097/MOP.0000000000000422
  26. Kasahara M, Horikawa R, Tagawa M, Uemoto S, Yokoyama S. Current role of liver transplantation for methylmalonic acidemia: a review of the literature. *Pediatr Transplant*. 2006; 10: 943-947. PMID: 17096763 DOI: 10.1111/j.1399-3046.2006.00585.x
  27. van 't Hoff WG, Dixon M, Taylor J, Mistry P, Rolles K. Combined liver-kidney transplantation in methylmalonic acidemia. *J Pediatr*. 1998; 132: 1043-1044. PMID: 9627602 DOI: 10.1016/s0022-3476(98)70407-x
  28. Kaplan P, Ficicioglu C, Mazur AT, Palmieri MJ, Berry GT. Liver transplantation is not curative for methylmalonic acidopathy caused by methylmalonyl-CoA mutase deficiency. *Mol Genet Metab*. 2006; 88: 322- 326. PMID: 16750411 DOI: 10.1016/j.ymgme.2006.04.003
  29. Niemi AK, Kim IK, Krueger CE, Cowan TM, Baugh N. Treatment of methylmalonic acidemia by liver or combined liver-kidney transplantation. *J Pediatr*. 2015; 166: 1455-1461. PMID: 25771389 DOI: 10.1016/j.jpeds.2015.01.051
  30. Sloan JL, Manoli I, Venditti CP (2015) Liver or combined liver– kidney transplantation for patients with isolated methylmalonic acidemia: who and when? *J Pediatr* 166:1346–1350 PMID: 25882873 DOI: 10.1016/j.jpeds.2015.03.026
  31. Wood RP, Ellis D, Starzl TE. The reversal of hepatorenal syndrome in four pediatric patients following successful orthotopic liver transplantation. *Ann Surg*. 1987;205:415–9. PMID: 3551858 DOI: 10.1097/00000658-198704000-00012

32. Davis CL. Impact of pretransplant renal failure: when is listing for kidney-liver indicated? *Liver Transpl.* 2005; (11 Suppl 2): S35-44. PMID: 16237696 DOI: 10.1002/lt.20617
33. Calinescu AM, Wildhaber BE, Poncet A, Toso C, McLin VA. Outcomes of combined liver-kidney transplantation in children: analysis of the scientific registry of transplant recipients. *Am J Transplant.* 2014;14(12):2861-2868. PMID: 25274400 DOI: 10.1111/ajt.12935
34. Herden U, Kemper M, Ganschow R, Klaassen I, Grabhorn E, Brinkert F, Nashan B, Fischer L. Surgical aspects and outcome of combined liver and kidney transplantation in children. *Transpl Int.* 2011; 24: 805-811. PMID: 21615550 DOI: 10.1111/j.1432-2277.2011.01278.x
35. Grenda R, Kaliciński P. Combined and sequential liver-kidney transplantation in children. *Pediatr Nephrol.* 2018;33(12):2227-2237. PMID: 29322327 DOI:10.1007/s00467-017-3880-4
36. Fong TL, Bunnapradist S, Jordan SC, Selby RR, Cho YW. Analysis of the United Network for Organ Sharing database comparing renal allografts and patient survival in combined liver-kidney transplantation with the contralateral allografts in kidney alone or kidney-pancreas transplantation. *Transplantation.* 2003;76: 348-353. PMID: 12883191 DOI: 10.1097/01.TP.0000071204.03720.BB
37. Ruiz R, Kunitake H, Wilkinson AH, Danovitch GM, Farmer DG, Ghobrial RM, Yersiz H, Hiatt JR, Busuttil RW. Long-term analysis of combined liver and kidney transplantation at a single center. *Arch Surg.* 2006;141(8):735-742. PMID: 16924080 DOI: 10.1001/archsurg.141.8.735
38. Bumgardner GL, Hardie I, Johnson RW, Lin A, Nashan B, Pescovitz M D, Ramos E, Vincenti F, Phase III Daclizumab Study Group. Results of 3-year phase III clinical trials with daclizumab prophylaxis for prevention of acute rejection after renal transplantation. *Transplantation.* 2001;72(5):839-845. PMID: 11571447 DOI: 10.1097/00007890-200109150-00017
39. Calne RY, Davis DR, Hadjiyannakis E, Sells R A, White D, Herbertson B M, Millard P R, Joysey V C, Davies D A, Binns R M, Festenstein H. Immunosuppressive effects of soluble cell membrane fractions, donor blood and serum on renal allograft survival. *Nature.* 1970;227:903-906. PMID: 4988655 DOI: 10.1038/227903a0
40. Knechtle SJ, Kwun J (2009) Unique aspects of rejection and tolerance in liver transplantation. *Semin Liver Dis* 29:91–101 PMID: 19235662 DOI: 10.1055/s-0029-1192058
41. Key T, Watson C, Clarworthy O’RCM, Goodman RS, Taylor CJ, Butler AJ (2010) The kinetics of donor HLA class I-specific antibody absorption following split liver and kidney transplant. *Nephrol Dial Transplant* 3:579–581 PMID: 25949473 DOI: 10.1093/ndtplus/sfq160
42. Bernabeu QJ, Juamperez J, Muñoz M, Rodríguez O, Vilalta R, A Molino JA, Asensio M, Bilbao I, Ariceta G, Rodrigo C, Charco R. Successful long-term outcome of pediatric liver–kidney transplantation: a single-center study. *Pediatr Nephrol* **33**, 351–358 (2018). PMID: 28842757 DOI: 10.1007/s00467-017-3782-5
43. Kniepiess D, Iberer F, Grasser B, Schaffellner S, Tscheliessnigg KH (2003) Sirolimus and mycophenolate mofetil after liver transplantation. *Transpl Int* 16:504–509 PMID: 12687325 DOI: 10.1007/s00147-003-0579-1

44. Herden U, Kemper M, Ganschow R, Klaassen I, Grabhorn E, Brinkert F, Nashan B, Fischer L. Surgical aspects and outcome of combined liver and kidney transplantation in children. *Transpl Int* 2011; 24: 805–811. PMID: 21615550 DOI: 10.1111/j.1432-2277.2011.01278.x
45. Broyer M, Le Bihan C, Charbit M, Guest G, Tete MJ, Gagnadoux MF, Niaudet P. Long-term social outcome of children after kidney transplantation. *Transplantation*. 2004;77(7):1033-7, PMID: 15087767 DOI: 10.1097/01.tp.0000120947.75697.8b
46. Alonso EM, Shepherd R, Martz KL, Yin W, Anand R. Linear growth patterns in prepubertal children following liver transplantation. *Am J Transplant*. 2009;9(6):1389-97. PMID: 19459820 DOI: 10.1111/j.1600-6143.2009.02634.x
47. North American Pediatric Renal Trials and Collaborative Studies. 2011 Annual Transplant Report.[30 October 2013]; Available from: <https://web.emmes.com/study/ped/annlrept/annualrept2011.pdf>.
48. Hodson EM, Willis NS, Craig JC. Growth hormone for children with chronic kidney disease. *Cochrane Database Syst Rev*. 2012;2012(2):CD003264. Published 2012 Feb 15. PMID: 22336787 DOI:10.1002/14651858.CD003264.pub3